wwPDB X-ray Structure Validation Summary Report (i)

Aug 21, 2023 – 08:08 PM EDT

PDB ID	:	2OYQ
Title	:	Crystal structure of RB69 gp43 in complex with DNA with 5-NIMP opposite an abasic site analog
Authors	:	Zahn, K.E.; Belrhali, H.; Wallace, S.S.; Doublie, S.
Deposited on	:	2007-02-22
Resolution	:	2.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	21	24% 52% 24%
1	G	21	19% 38% 14% 48%
1	I	21	10% 57% 29% 10% 5%
1	K	21	38% 19% 43%
2	F	15	47% 40% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	15	47% 27% 33% 33% 7%
2	J	15	33% 47% 20%
2	L	15	7% 7% 67% 13% 13%
3	A	903	% 73% 19% • 6%
3	B	903	3% 62% 20% • 16%
3	C	903	3% 77% 19% • •
3	D	903	11% 74% 14% • 11%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 28889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	E	16	314	150	56	93	15	0	0	0
1	G	11	223	106	44	63	10	0	0	0
1	I	20	395	188	72	116	19	0	0	0
1	K	12	244	116	49	68	11	0	0	0

- Molecule 2 is a DNA chain called Primer DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	F	15	310	150	57	89	14	0	0	0
2	H	14	287	137	55	82	13	0	0	0
2	J	15	310	150	57	89	14	0	0	0
2	L	13	265	127	50	76	12	0	0	0

- Molecule 3 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	848	6877	4423	1143	1280	31	0	0	0
3	B	756	6045	3883	1001	1134	27	0	0	0
3	C	891	7150	4583	1182	1352	33	0	0	0
3	D	807	5688	3586	953	1125	24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	ALA	ASP	engineered mutation	UNP Q38087
A	327	ALA	ASP	engineered mutation	UNP Q38087
B	222	ALA	ASP	engineered mutation	UNP Q38087
B	327	ALA	ASP	engineered mutation	UNP Q38087
C	222	ALA	ASP	engineered mutation	UNP Q38087
C	327	ALA	ASP	engineered mutation	UNP Q38087
D	222	ALA	ASP	engineered mutation	UNP Q38087
D	327	ALA	ASP	engineered mutation	UNP Q38087

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Mg			
4	I	1	1	1		0	0
			Total	Mg			
4	K	1	1	1		0	0

- Molecule 5 is 1-{2-DEOXY-5-O-[(R)-HYDROXY{[(R)-HYDROXY(PHOSPHONOOXY)PHOSPHORYL]OXY}PHOSPHORYL]-BETA-D-ERYTHRO-PENTOFURANOSYL}-5-NITRO -1H-INDOLE (three-letter code: N5P) (formula: $C_{13}H_{17}N_2O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	I	1	32	13	2	14	3	0	0
			Total	C	N	O	P		
5	A	1	32	13	2	14	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	32	13	2	14	3	0	0
5	C	1	32	13	2	14	3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	E	5	5	5	0	0
6	F	8	8	8	0	0
6	G	2	2	2	0	0
6	H	3	3	3	0	0
6	I	22	22	22	0	0
6	J	19	19	19	0	0
6	K	3	3	3	0	0
6	L	3	3	3	0	0
6	A	235	235	235	0	0
6	B	125	125	125	0	0
6	C	185	185	185	0	0
6	D	41	41	41	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Template DNA

Chain E:





- Molecule 1: Template DNA

Chain G:





- Molecule 1: Template DNA

Chain I:



- Molecule 1: Template DNA

Chain K:





- Molecule 2: Primer DNA

Chain F:

- Molecule 2: Primer DNA

Chain H:



• Molecule 2: Primer DNA

Chain J:





• Molecule 2: Primer DNA

Chain L:



• Molecule 3: DNA polymerase

Chain A:



• Molecule 3: DNA polymerase

Chain B:



- Molecule 3: DNA polymerase

Chain C:





• Molecule 3: DNA polymerase

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	132.59Å 123.48Å 164.33Å 90.00° 96.22° 90.00°	Depositor
Resolution (Å)	30.00 – 2.86 49.82 – 2.85	Depositor EDS
% Data completeness (in resolution range)	91.2 (30.00-2.86) 96.3 (49.82-2.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.85 (at 2.86Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.228 , 0.294 0.231 , 0.289	Depositor DCC
R_{free} test set	23125 reflections (9.68%)	wwPDB-VP
Wilson B-factor (Å^2)	62.5	Xtriage
Anisotropy	0.210	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 48.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	28889	wwPDB-VP
Average B, all atoms (Å^2)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: N5I, N5P, MG, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	E	0.55	0/342	1.04	0/525
1	G	0.76	0/250	1.23	2/384 (0.5%)
1	I	0.77	0/429	1.45	5/657 (0.8%)
1	K	0.68	0/274	1.31	3/421 (0.7%)
2	F	0.69	0/322	1.42	4/496 (0.8%)
2	H	0.75	0/322	1.44	7/496 (1.4%)
2	J	0.66	0/322	1.41	9/496 (1.8%)
2	L	0.74	0/297	1.41	4/457 (0.9%)
3	A	0.42	0/7049	0.55	0/9536
3	B	0.40	0/6193	0.52	0/8393
3	C	0.40	0/7324	0.53	0/9912
3	D	0.34	0/5812	0.47	0/7955
All	All	0.43	0/28936	0.65	34/39728 (0.1%)

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	I	7	DC	O4'-C1'-N1	9.80	114.86	108.00
2	H	105	DC	O4'-C1'-N1	7.96	113.57	108.00
1	I	20	DC	O4'-C1'-N1	7.69	113.38	108.00
2	F	113	DA	O4'-C1'-N9	7.59	113.32	108.00
1	K	20	DC	O4'-C1'-N1	7.59	113.32	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	314	0	177	20	0
1	G	223	0	124	2	0
1	I	395	0	222	4	0
1	K	244	0	135	3	0
2	F	310	0	171	6	0
2	H	287	0	159	11	0
2	J	310	0	171	13	0
2	L	265	0	148	6	0
3	A	6877	0	6731	115	0
3	B	6045	0	5799	124	0
3	C	7150	0	6884	116	0
3	D	5688	0	4707	64	0
4	I	1	0	0	0	0
4	K	1	0	0	0	0
5	A	64	0	26	1	0
5	C	32	0	13	0	0
5	I	32	0	13	0	0
6	A	235	0	0	4	0
6	B	125	0	0	7	0
6	C	185	0	0	2	0
6	D	41	0	0	1	0
6	E	5	0	0	2	0
6	F	8	0	0	1	0
6	G	2	0	0	0	0
6	H	3	0	0	1	0
6	I	22	0	0	1	0
6	J	19	0	0	0	0
6	K	3	0	0	0	0
6	L	3	0	0	0	0
All	All	28889	0	25480	469	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 469 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:13:DA:H2"	1:E:14:DC:H5"	1.29	1.07
3:A:356:GLN:H	3:A:356:GLN:HE21	1.02	0.98
1:E:8:DT:O4	2:F:113:DA:N6	2.03	0.91
1:E:13:DA:C2'	1:E:14:DC:H5"	2.00	0.91
1:E:8:DT:H4'	1:E:9:DT:OP1	1.70	0.91

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	844/903 (94%)	794 (94%)	45 (5%)	5 (1%)	25	53
3	B	750/903 (83%)	697 (93%)	49 (6%)	4 (0%)	29	57
3	C	885/903 (98%)	817 (92%)	61 (7%)	7 (1%)	19	46
3	D	797/903 (88%)	701 (88%)	83 (10%)	13 (2%)	9	28
All	All	3276/3612 (91%)	3009 (92%)	238 (7%)	29 (1%)	17	43

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	622	THR
3	B	736	SER
3	C	622	THR
3	D	622	THR
3	A	549	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	738/800 (92%)	708 (96%)	30 (4%)	30	61
3	B	634/800 (79%)	605 (95%)	29 (5%)	27	56
3	C	755/800 (94%)	714 (95%)	41 (5%)	22	49
3	D	483/800 (60%)	453 (94%)	30 (6%)	18	43
All	All	2610/3200 (82%)	2480 (95%)	130 (5%)	24	53

5 of 130 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	373	LEU
3	D	602	ASN
3	B	403	ARG
3	B	326	ILE
3	D	618	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	591	GLN
3	D	678	GLN
3	C	595	GLN
3	D	572	ASN
3	D	754	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	N5I	F	115	2	20,25,26	0.60	0	25,36,39	0.98	1 (4%)
1	3DR	E	6	1	8,8,12	0.41	0	9,10,17	0.77	0
2	N5I	J	115	2	20,25,26	0.56	0	25,36,39	1.09	1 (4%)
1	3DR	I	6	1	8,11,12	0.50	0	9,14,17	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N5I	F	115	2	-	0/5/25/26	0/3/3/3
1	3DR	E	6	1	-	2/2/12/16	0/1/1/1
2	N5I	J	115	2	-	0/5/25/26	0/3/3/3
1	3DR	I	6	1	-	2/3/15/16	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	J	115	N5I	CD1-NE1-C1'	-2.86	122.57	125.40
2	F	115	N5I	CD1-NE1-C1'	-2.26	123.16	125.40

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	E	6	3DR	C3'-C4'-C5'-O5'
1	E	6	3DR	O4'-C4'-C5'-O5'
1	I	6	3DR	C3'-C4'-C5'-O5'
1	I	6	3DR	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	115	N5I	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	N5P	A	905	4	28,34,34	0.53	0	38,53,53	1.16	2 (5%)
5	N5P	C	904	-	28,34,34	0.56	0	38,53,53	1.15	3 (7%)
5	N5P	I	704	4	28,34,34	0.54	0	38,53,53	1.17	2 (5%)
5	N5P	A	904	-	28,34,34	0.59	0	38,53,53	1.16	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	N5P	A	905	4	-	4/20/38/38	0/3/3/3
5	N5P	C	904	-	-	0/20/38/38	0/3/3/3
5	N5P	I	704	4	-	4/20/38/38	0/3/3/3
5	N5P	A	904	-	-	4/20/38/38	0/3/3/3

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	904	N5P	PB-O3A-PA	-3.43	121.04	132.83
5	I	704	N5P	PB-O3B-PG	-3.02	122.48	132.83
5	I	704	N5P	PB-O3A-PA	-3.00	122.54	132.83
5	A	905	N5P	PB-O3B-PG	-2.92	122.82	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	C	904	N5P	PB-O3A-PA	-2.85	123.06	132.83

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	904	N5P	O4'-C4'-C5'-O5'
5	A	904	N5P	C3'-C4'-C5'-O5'
5	I	704	N5P	PA-O3A-PB-O1B
5	A	905	N5P	PA-O3A-PB-O1B
5	A	905	N5P	PB-O3A-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	905	N5P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand N5P A 905



Ligand N5P C 904



Ligand N5P I 704



Ligand N5P A 904



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	E	15/21 (71%)	0.12	0 100 100	58, 78, 115, 119	0
1	G	11/21 (52%)	1.39	4 (36%) 0 0	62, 115, 126, 132	0
1	I	19/21 (90%)	0.01	2 (10%) 6 4	43, 56, 138, 144	0
1	K	12/21 (57%)	-0.18	0 100 100	42, 94, 107, 115	0
2	F	14/15 (93%)	-0.02	0 100 100	73, 106, 124, 126	0
2	H	14/15 (93%)	1.92	7 (50%) 0 0	102, 129, 134, 137	0
2	J	14/15 (93%)	-0.39	0 100 100	44, 68, 85, 88	0
2	L	13/15 (86%)	0.64	1 (7%) 13 10	85, 108, 113, 113	0
3	A	848/903 (93%)	0.04	6 (0%) 87 87	36, 53, 79, 96	0
3	B	756/903 (83%)	0.21	29 (3%) 40 35	44, 68, 105, 116	0
3	C	891/903 (98%)	0.25	29 (3%) 46 41	36, 63, 110, 135	0
3	D	807/903 (89%)	0.71	96 (11%) 4 3	105, 121, 145, 147	0
All	All	3414/3756 (90%)	0.30	174 (5%) 28 23	36, 69, 138, 147	0

The worst 5 of 174 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	257	TYR	8.0
3	D	395	PHE	6.8
3	D	162	TRP	6.5
3	C	252	VAL	6.4
3	B	259	SER	5.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	N5I	F	115	23/24	0.80	0.26	127,128,128,128	0
1	3DR	E	6	8/12	0.81	0.31	123,124,124,124	0
1	3DR	I	6	11/12	0.89	0.20	104,108,115,115	0
2	N5I	J	115	23/24	0.93	0.18	83,89,89,89	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MG	I	802	1/1	0.73	0.10	82,82,82,82	1
4	MG	K	801	1/1	0.80	0.24	67,67,67,67	1
5	N5P	C	904	32/32	0.80	0.29	79,81,83,83	32
5	N5P	A	904	32/32	0.86	0.20	74,76,88,88	32
5	N5P	I	704	32/32	0.87	0.28	78,80,84,84	32
5	N5P	A	905	32/32	0.92	0.22	61,65,75,76	32

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around N5P C 904:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around N5P A 904:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around N5P I 704:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around N5P A 905:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.