wwPDB X-ray Structure Validation Summary Report (i)

Dec 16, 2023 – 10:20 AM EST

PDB ID	:	4P4X
Title	:	Dodecamer formed by a macrocyclic peptide derived from beta-2-microglobulin (63-69) - (ORN)YLL(PHI)YTE(ORN)KVA(MLE)AVK
Authors	:	Spencer, R.K.; Nowick, J.S.
Deposited on	:	2014-03-13
Resolution	:	1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	16	81% 12% 6%
1	B	16	75% 19% 6%
1	C	16	75% 19% 6%
1	D	16	81% 12% 6%
1	E	16	88% 6% 6%
1	F	16	81% 12% 6%
1	G	16	69% 31%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	16	88% 12%
1	I	16	88% 6% 6%
1	J	16	75% 25%
1	K	16	75% 25%
1	L	16	75% 19% 6%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3507 atoms, of which 1775 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	I	N	O			
1	A	16	283	93	148	1	20	21	0	0	0
1	B	16	283	93	148	1	20	21	0	0	0
1	C	16	283	93	148	1	20	21	0	0	0
1	D	16	283	93	148	1	20	21	0	0	0
1	E	16	283	93	148	1	20	21	0	0	0
1	F	16	283	93	148	1	20	21	0	0	0
1	G	16	283	93	148	1	20	21	0	0	0
1	H	16	282	93	147	1	20	21	0	0	0
1	I	16	283	93	148	1	20	21	0	0	0
1	J	16	283	93	148	1	20	21	0	0	0
1	K	16	283	93	148	1	20	21	0	0	0
1	L	16	283	93	148	1	20	21	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	G	1	Total O S 5 4 1	0	0
2	G	1	Total O S 5 4 1	0	0
2	K	1	Total O S 5 4 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	K	1	Total Cl 1 1	0	0
3	L	1	Total Cl 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	10	10	10	0	0
			Total	O		
4	B	8	8	8	0	0
			Total	O		
4	C	10	10	10	0	0
			Total	O		
4	D	4	4	4	0	0
			Total	O		
4	E	8	8	8	0	0
			Total	O		
4	F	4	4	4	0	0
			Total	O		
4	G	7	7	7	0	0
			Total	O		
4	H	4	4	4	0	0
			Total	O		
4	I	3	3	3	0	0
			Total	O		
4	J	7	7	7	0	0
			Total	O		
4	K	3	3	3	0	0
			Total	O		
4	L	2	2	2	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain A: 81% 12% 6%



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain B: 75% 19% 6%



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain C: 75% 19% 6%



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain D: 81% 12% 6%



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain E: 88% 6% 6%



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain F:

• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain G:



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain H:

• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain I:

• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain J:



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain K:



• Molecule 1: CYCLIC HEXADECAPEPTIDE (ORN)YLL(PH7)YTE(ORN)KVA(MLE)AVK

Chain L:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α, β, γ	61.34Å 61.34Å 113.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.30 – 1.90 32.30 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (32.30-1.90) 94.6 (32.30-1.90)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.08 (at 1.89Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.209 , 0.235 0.263 , 0.272	Depositor DCC
R_{free} test set	1793 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å^2)	28.5	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 51.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	3507	wwPDB-VP
Average B, all atoms (Å^2)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.85% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ORN, PHI, CL, MLE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.51	0/96	0.63	0/124
1	B	0.64	0/96	0.66	0/124
1	C	0.62	0/96	0.62	0/124
1	D	0.54	0/96	0.56	0/124
1	E	0.50	0/96	0.78	0/124
1	F	0.48	0/96	0.69	0/124
1	G	0.57	0/96	0.61	0/124
1	H	0.51	0/96	0.70	0/124
1	I	0.45	0/96	0.75	0/124
1	J	0.46	0/96	0.61	0/124
1	K	0.40	0/96	0.74	0/124
1	L	0.42	0/96	0.78	0/124
All	All	0.51	0/1152	0.68	0/1488

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	135	148	145	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	135	148	147	3	0
1	C	135	148	146	4	0
1	D	135	148	146	2	0
1	E	135	148	145	1	0
1	F	135	148	147	1	0
1	G	135	148	147	6	0
1	H	135	147	146	0	0
1	I	135	148	145	1	0
1	J	135	148	145	5	0
1	K	135	148	145	3	0
1	L	135	148	145	2	0
2	A	15	0	0	0	0
2	C	5	0	0	0	0
2	E	5	0	0	0	0
2	G	10	0	0	2	0
2	K	5	0	0	1	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
4	A	10	0	0	0	0
4	B	8	0	0	0	0
4	C	10	0	0	1	0
4	D	4	0	0	1	0
4	E	8	0	0	0	0
4	F	4	0	0	0	0
4	G	7	0	0	1	0
4	H	4	0	0	0	0
4	I	3	0	0	0	0
4	J	7	0	0	1	0
4	K	3	0	0	0	0
4	L	2	0	0	1	0
All	All	1732	1775	1749	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:16:LYS:NZ	4:L:201:HOH:O	2.20	0.75
1:B:1:ORN:N	1:C:7:THR:O	2.21	0.72
1:G:16:LYS:NZ	2:G:102:SO4:O1	2.35	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:7:THR:OG1	2:K:101:SO4:O3	2.22	0.58
2:G:101:SO4:O3	4:G:201:HOH:O	2.18	0.53

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	10/16 (62%)	10 (100%)	0	0	100	100
1	B	10/16 (62%)	10 (100%)	0	0	100	100
1	C	10/16 (62%)	10 (100%)	0	0	100	100
1	D	10/16 (62%)	10 (100%)	0	0	100	100
1	E	10/16 (62%)	10 (100%)	0	0	100	100
1	F	10/16 (62%)	10 (100%)	0	0	100	100
1	G	10/16 (62%)	10 (100%)	0	0	100	100
1	H	10/16 (62%)	10 (100%)	0	0	100	100
1	I	10/16 (62%)	10 (100%)	0	0	100	100
1	J	10/16 (62%)	10 (100%)	0	0	100	100
1	K	10/16 (62%)	10 (100%)	0	0	100	100
1	L	10/16 (62%)	10 (100%)	0	0	100	100
All	All	120/192 (62%)	120 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/10 (100%)	10 (100%)	0	100	100
1	B	10/10 (100%)	10 (100%)	0	100	100
1	C	10/10 (100%)	10 (100%)	0	100	100
1	D	10/10 (100%)	10 (100%)	0	100	100
1	E	10/10 (100%)	10 (100%)	0	100	100
1	F	10/10 (100%)	10 (100%)	0	100	100
1	G	10/10 (100%)	10 (100%)	0	100	100
1	H	10/10 (100%)	10 (100%)	0	100	100
1	I	10/10 (100%)	10 (100%)	0	100	100
1	J	10/10 (100%)	10 (100%)	0	100	100
1	K	10/10 (100%)	10 (100%)	0	100	100
1	L	10/10 (100%)	9 (90%)	1 (10%)	7	3
All	All	120/120 (100%)	119 (99%)	1 (1%)	81	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	4	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

48 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	ORN	J	1	1	6,7,8	0.42	0	2,7,9	0.32	0
1	ORN	A	9	1	6,7,8	0.48	0	2,7,9	0.45	0
1	MLE	G	13	1	7,8,9	1.20	1 (14%)	6,9,11	0.82	0
1	MLE	D	13	1	7,8,9	1.01	0	6,9,11	0.63	0
1	MLE	E	13	1	7,8,9	0.98	0	6,9,11	1.13	1 (16%)
1	ORN	H	9	1	6,7,8	0.47	0	2,7,9	0.44	0
1	MLE	K	13	1	7,8,9	0.85	0	6,9,11	0.84	0
1	ORN	B	1	1	6,7,8	1.09	1 (16%)	2,7,9	0.14	0
1	ORN	B	9	1	6,7,8	0.49	0	2,7,9	0.39	0
1	PHI	D	5	1	11,12,13	1.67	2 (18%)	12,15,17	1.31	1 (8%)
1	ORN	E	1	1	6,7,8	0.72	0	2,7,9	0.27	0
1	ORN	C	1	1	6,7,8	0.64	0	2,7,9	0.09	0
1	ORN	C	9	1	6,7,8	0.79	0	2,7,9	0.50	0
1	PHI	F	5	1	11,12,13	1.83	4 (36%)	12,15,17	1.41	2 (16%)
1	PHI	G	5	1	11,12,13	2.00	3 (27%)	12,15,17	1.37	1 (8%)
1	ORN	L	9	1	6,7,8	0.65	0	2,7,9	0.46	0
1	PHI	J	5	1	11,12,13	1.50	3 (27%)	12,15,17	1.39	2 (16%)
1	MLE	B	13	1	7,8,9	0.86	0	6,9,11	0.93	0
1	ORN	H	1	1	6,7,8	0.99	0	2,7,9	0.14	0
1	PHI	K	5	1	11,12,13	1.63	3 (27%)	12,15,17	1.22	2 (16%)
1	ORN	A	1	1	6,7,8	0.95	0	2,7,9	0.27	0
1	MLE	L	13	1	7,8,9	1.08	0	6,9,11	1.03	0
1	PHI	A	5	1	11,12,13	1.63	2 (18%)	12,15,17	0.95	0
1	ORN	G	9	1	6,7,8	0.44	0	2,7,9	0.77	0
1	MLE	C	13	1	7,8,9	0.70	0	6,9,11	0.78	0
1	MLE	F	13	1	7,8,9	1.23	1 (14%)	6,9,11	0.81	0
1	MLE	J	13	1	7,8,9	0.81	0	6,9,11	0.94	0
1	PHI	H	5	1	11,12,13	1.72	3 (27%)	12,15,17	1.51	1 (8%)
1	ORN	I	1	1	6,7,8	0.64	0	2,7,9	0.15	0
1	PHI	B	5	1	11,12,13	1.86	4 (36%)	12,15,17	1.35	1 (8%)
1	PHI	C	5	1	11,12,13	2.02	5 (45%)	12,15,17	1.66	3 (25%)
1	ORN	D	1	1	6,7,8	0.56	0	2,7,9	0.16	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	ORN	I	9	1	6,7,8	0.55	0	2,7,9	0.39	0
1	ORN	D	9	1	6,7,8	0.50	0	2,7,9	0.50	0
1	ORN	F	1	1	6,7,8	1.19	1 (16%)	2,7,9	0.32	0
1	PHI	E	5	1	11,12,13	1.55	3 (27%)	12,15,17	1.60	3 (25%)
1	PHI	I	5	1	11,12,13	1.46	3 (27%)	12,15,17	1.47	2 (16%)
1	ORN	G	1	1	6,7,8	0.57	0	2,7,9	0.22	0
1	MLE	A	13	1	7,8,9	0.90	0	6,9,11	1.29	1 (16%)
1	ORN	E	9	1	6,7,8	0.69	0	2,7,9	0.54	0
1	PHI	L	5	1	11,12,13	1.67	3 (27%)	12,15,17	0.96	1 (8%)
1	ORN	L	1	1	6,7,8	0.78	0	2,7,9	0.33	0
1	ORN	F	9	1	6,7,8	0.38	0	2,7,9	0.45	0
1	ORN	K	9	1	6,7,8	0.72	0	2,7,9	0.43	0
1	MLE	H	13	1	7,8,9	1.15	1 (14%)	6,9,11	1.10	0
1	ORN	J	9	1	6,7,8	0.58	0	2,7,9	0.33	0
1	MLE	I	13	1	7,8,9	0.87	0	6,9,11	0.94	1 (16%)
1	ORN	K	1	1	6,7,8	0.41	0	2,7,9	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ORN	J	1	1	-	0/5/6/8	-
1	ORN	A	9	1	-	0/5/6/8	-
1	MLE	G	13	1	-	0/5/8/10	-
1	MLE	D	13	1	-	1/5/8/10	-
1	MLE	E	13	1	-	1/5/8/10	-
1	ORN	H	9	1	-	0/5/6/8	-
1	MLE	K	13	1	-	1/5/8/10	-
1	ORN	B	1	1	-	0/5/6/8	-
1	ORN	B	9	1	-	0/5/6/8	-
1	PHI	D	5	1	-	2/5/6/8	0/1/1/1
1	ORN	E	1	1	-	0/5/6/8	-
1	ORN	C	1	1	-	1/5/6/8	-
1	ORN	C	9	1	-	1/5/6/8	-
1	PHI	F	5	1	-	0/5/6/8	0/1/1/1
1	PHI	G	5	1	-	1/5/6/8	0/1/1/1
1	ORN	L	9	1	-	1/5/6/8	-
1	PHI	J	5	1	-	0/5/6/8	0/1/1/1
1	MLE	B	13	1	-	0/5/8/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ORN	H	1	1	-	1/5/6/8	-
1	PHI	K	5	1	-	0/5/6/8	0/1/1/1
1	ORN	A	1	1	-	0/5/6/8	-
1	MLE	L	13	1	-	5/5/8/10	-
1	PHI	A	5	1	-	0/5/6/8	0/1/1/1
1	ORN	G	9	1	-	1/5/6/8	-
1	MLE	C	13	1	-	0/5/8/10	-
1	MLE	F	13	1	-	0/5/8/10	-
1	MLE	J	13	1	-	0/5/8/10	-
1	PHI	H	5	1	-	1/5/6/8	0/1/1/1
1	ORN	I	1	1	-	0/5/6/8	-
1	PHI	B	5	1	-	0/5/6/8	0/1/1/1
1	PHI	C	5	1	-	0/5/6/8	0/1/1/1
1	ORN	D	1	1	-	0/5/6/8	-
1	ORN	I	9	1	-	4/5/6/8	-
1	ORN	D	9	1	-	1/5/6/8	-
1	ORN	F	1	1	-	1/5/6/8	-
1	PHI	E	5	1	-	0/5/6/8	0/1/1/1
1	PHI	I	5	1	-	0/5/6/8	0/1/1/1
1	ORN	G	1	1	-	0/5/6/8	-
1	MLE	A	13	1	-	2/5/8/10	-
1	ORN	E	9	1	-	4/5/6/8	-
1	PHI	L	5	1	-	1/5/6/8	0/1/1/1
1	ORN	L	1	1	-	1/5/6/8	-
1	ORN	F	9	1	-	0/5/6/8	-
1	ORN	K	9	1	-	0/5/6/8	-
1	MLE	H	13	1	-	0/5/8/10	-
1	ORN	J	9	1	-	1/5/6/8	-
1	MLE	I	13	1	-	3/5/8/10	-
1	ORN	K	1	1	-	0/5/6/8	-

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	5	PHI	CD1-CG	3.96	1.47	1.38
1	H	5	PHI	CD1-CG	3.81	1.47	1.38
1	A	5	PHI	CD1-CG	3.70	1.46	1.38
1	C	5	PHI	CD1-CG	3.65	1.46	1.38
1	B	5	PHI	CE2-CZ	3.62	1.46	1.38

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	H	5	PHI	CG-CB-CA	-4.25	105.50	114.10
1	B	5	PHI	CG-CB-CA	-3.99	106.03	114.10
1	E	5	PHI	CG-CB-CA	-3.46	107.09	114.10
1	F	5	PHI	CG-CB-CA	-3.44	107.13	114.10
1	C	5	PHI	CG-CB-CA	-3.44	107.13	114.10

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	1	ORN	O-C-CA-CB
1	F	1	ORN	O-C-CA-CB
1	H	5	PHI	O-C-CA-CB
1	L	5	PHI	O-C-CA-CB
1	D	9	ORN	O-C-CA-CB

There are no ring outliers.

11 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	J	1	ORN	1	0
1	D	13	MLE	1	0
1	K	13	MLE	1	0
1	B	1	ORN	1	0
1	D	5	PHI	1	0
1	A	5	PHI	1	0
1	G	9	ORN	1	0
1	F	13	MLE	1	0
1	C	5	PHI	1	0
1	E	5	PHI	1	0
1	I	13	MLE	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	SO4	E	101	-	4,4,4	0.20	0	6,6,6	0.39	0
2	SO4	G	102	-	4,4,4	0.13	0	6,6,6	0.14	0
2	SO4	K	101	-	4,4,4	0.19	0	6,6,6	0.37	0
2	SO4	C	101	-	4,4,4	0.14	0	6,6,6	0.08	0
2	SO4	G	101	-	4,4,4	0.13	0	6,6,6	0.18	0
2	SO4	A	101	-	4,4,4	0.16	0	6,6,6	0.15	0
2	SO4	A	102	-	4,4,4	0.15	0	6,6,6	0.11	0
2	SO4	A	103	-	4,4,4	0.14	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	102	SO4	1	0
2	K	101	SO4	1	0
2	G	101	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.