Full wwPDB X-ray Structure Validation Report (i)

Jun 23, 2024 – 01:04 AM EDT

PDB ID	:	6P61
Title	:	Structure of a Glycosyltransferase from Leptospira borgpetersenii serovar Hardjo-bovis (strain JB197)
Authors	:	Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2019-05-31
Resolution	:	1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	2%, 82%, •, 15%
1	B	228	5%, 80%, •, 16%
1	C	228	6%, 75%, 8%, 17%
1	D	228	2%, 81%, •, 15%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	193	1611	1034	269	301	7	0	11	0
1	B	191	1572	1006	263	296	7	0	8	0
1	C	190	1563	1001	259	296	7	0	11	0
1	D	193	1600	1027	267	299	7	0	9	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q04TP4
A	2	ALA	-	expression tag	UNP Q04TP4
A	3	HIS	-	expression tag	UNP Q04TP4
A	4	HIS	-	expression tag	UNP Q04TP4
A	5	HIS	-	expression tag	UNP Q04TP4
A	6	HIS	-	expression tag	UNP Q04TP4
A	7	HIS	-	expression tag	UNP Q04TP4
A	8	HIS	-	expression tag	UNP Q04TP4
B	1	MET	-	initiating methionine	UNP Q04TP4
B	2	ALA	-	expression tag	UNP Q04TP4
B	3	HIS	-	expression tag	UNP Q04TP4
B	4	HIS	-	expression tag	UNP Q04TP4
B	5	HIS	-	expression tag	UNP Q04TP4
B	6	HIS	-	expression tag	UNP Q04TP4
B	7	HIS	-	expression tag	UNP Q04TP4
B	8	HIS	-	expression tag	UNP Q04TP4
C	1	MET	-	initiating methionine	UNP Q04TP4
C	2	ALA	-	expression tag	UNP Q04TP4
C	3	HIS	-	expression tag	UNP Q04TP4
C	4	HIS	-	expression tag	UNP Q04TP4
C	5	HIS	-	expression tag	UNP Q04TP4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	HIS	-	expression tag	UNP Q04TP4
C	7	HIS	-	expression tag	UNP Q04TP4
C	8	HIS	-	expression tag	UNP Q04TP4
D	1	MET	-	initiating methionine	UNP Q04TP4
D	2	ALA	-	expression tag	UNP Q04TP4
D	3	HIS	-	expression tag	UNP Q04TP4
D	4	HIS	-	expression tag	UNP Q04TP4
D	5	HIS	-	expression tag	UNP Q04TP4
D	6	HIS	-	expression tag	UNP Q04TP4
D	7	HIS	-	expression tag	UNP Q04TP4
D	8	HIS	-	expression tag	UNP Q04TP4

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
2	A	1	4	2	2		0	0
2	A	1	4	2	2		0	0
2	A	1	4	2	2		0	0
2	B	1	4	2	2		0	0
2	B	1	4	2	2		0	0
2	B	1	4	2	2		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 6 7	0	0
3	A	1	Total C O 13 6 7	0	0
3	B	1	Total C O 13 6 7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C O 13 6 7	0	0
3	D	1	Total C O 13 6 7	0	0
3	D	1	Total C O 13 6 7	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	204	Total O 206 206	0	2
5	B	155	Total O 158 158	0	3
5	C	133	Total O 133 133	0	0
5	D	222	Total O 226 226	0	4

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glycosyltransferase

Chain A:

• Molecule 1: Glycosyltransferase

Chain B:



• Molecule 1: Glycosyltransferase

Chain C:



• Molecule 1: Glycosyltransferase

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	56.25Å 123.17Å 76.77Å 90.00° 92.44° 90.00°	Depositor
Resolution (Å)	33.15 – 1.95 46.28 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (33.15-1.95) 99.1 (46.28-1.95)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.06 (at 1.95Å)	Xtriage
Refinement program	PHENIX (dev_3500)	Depositor
R, R_{free}	0.151 , 0.183 0.153 , 0.185	Depositor DCC
R_{free} test set	2078 reflections (2.76%)	wwPDB-VP
Wilson B-factor (Å^2)	26.7	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 54.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.035 for h,-k,-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	7215	wwPDB-VP
Average B, all atoms (Å^2)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, EDO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.37	0/1675	0.50	0/2263
1	B	0.35	0/1625	0.52	0/2193
1	C	0.36	0/1623	0.50	0/2192
1	D	0.38	0/1655	0.52	0/2236
All	All	0.37	0/6578	0.51	0/8884

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1611	0	1603	5	0
1	B	1572	0	1557	7	0
1	C	1563	0	1550	15	0
1	D	1600	0	1588	6	0
2	A	12	0	18	1	0
2	B	12	0	18	0	0
2	C	16	0	24	2	0
2	D	16	0	24	1	0
3	A	26	0	10	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	13	0	5	0	0
3	C	13	0	5	0	0
3	D	26	0	10	0	0
4	A	6	0	8	1	0
4	D	6	0	8	1	0
5	A	206	0	0	2	0
5	B	158	0	0	1	0
5	C	133	0	0	1	0
5	D	226	0	0	2	0
All	All	7215	0	6428	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:27:TYR:OH	1:C:135:ASP:OD2	2.14	0.61
1:A:22:PHE:HB3	3:A:304:CIT:H22	1.83	0.60
1:B:42:LYS:HE3	5:B:411:HOH:O	2.06	0.55
4:A:305:GOL:O3	5:A:401:HOH:O	2.16	0.54
1:B:109:TYR:OH	1:B:153:GLY:HA2	2.08	0.54
1:D:19:VAL:HB	1:D:49:ILE:HD13	1.89	0.53
3:A:303:CIT:H21	5:A:431:HOH:O	2.10	0.52
1:C:153:GLY:HA3	2:C:301:EDO:H22	1.91	0.52
1:B:68:GLU:OE2	1:C:33[B]:ARG:HD2	2.11	0.50
4:D:306:GOL:O1	5:D:401:HOH:O	2.11	0.50
1:C:109:TYR:OH	1:C:153:GLY:HA2	2.12	0.49
1:A:41:PRO:HD3	2:A:306:EDO:H12	1.95	0.48
1:B:33[B]:ARG:HD2	1:C:68:GLU:OE2	2.13	0.48
1:D:40:PHE:CD1	2:D:307:EDO:H12	2.50	0.47
1:C:14[B]:THR:HG23	5:C:413:HOH:O	2.14	0.47
1:B:16:SER:HB2	1:B:96:ILE:HD13	1.96	0.46
1:D:184:GLU:HA	1:D:187:TYR:O	2.17	0.45
1:A:14[B]:THR:HG22	1:A:95:PHE:HD2	1.81	0.45
1:B:65:PHE:HA	1:C:33[A]:ARG:HD2	1.99	0.44
1:C:16:SER:HB2	1:C:96:ILE:HD13	2.00	0.44
1:C:167:TYR:CZ	1:C:179:LEU:HD22	2.53	0.44
1:A:184:GLU:HA	1:A:187:TYR:O	2.19	0.43
1:D:27:TYR:OH	1:D:135:ASP:OD1	2.16	0.43
1:B:33[A]:ARG:HD2	1:C:65:PHE:HA	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:LYS:HB2	1:A:115:LYS:HE2	1.89	0.42
1:C:184:GLU:HA	1:C:187:TYR:O	2.19	0.42
1:D:14[A]:THR:HG22	1:D:95:PHE:HD2	1.85	0.41
1:C:167:TYR:CE1	1:C:179:LEU:HD22	2.56	0.41
1:C:40:PHE:HA	2:C:305:EDO:H21	2.03	0.40
1:C:98:ARG:NE	1:C:152[A]:CYS:SG	2.94	0.40
1:D:14[B]:THR:HG23	5:D:453:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/228 (89%)	199 (98%)	3 (2%)	0	100	100
1	B	197/228 (86%)	192 (98%)	5 (2%)	0	100	100
1	C	197/228 (86%)	193 (98%)	4 (2%)	0	100	100
1	D	200/228 (88%)	198 (99%)	2 (1%)	0	100	100
All	All	796/912 (87%)	782 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/208 (86%)	180 (100%)	0	100	100
1	B	174/208 (84%)	174 (100%)	0	100	100
1	C	174/208 (84%)	174 (100%)	0	100	100
1	D	178/208 (86%)	178 (100%)	0	100	100
All	All	706/832 (85%)	706 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	EDO	D	301	-	3,3,3	0.59	0	2,2,2	0.45	0
2	EDO	D	302	-	3,3,3	0.45	0	2,2,2	0.50	0
2	EDO	D	307	-	3,3,3	0.46	0	2,2,2	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	EDO	C	301	-	3,3,3	0.49	0	2,2,2	0.39	0
2	EDO	C	305	-	3,3,3	0.46	0	2,2,2	0.32	0
4	GOL	D	306	-	5,5,5	0.94	0	5,5,5	1.07	1 (20%)
2	EDO	C	302	-	3,3,3	0.41	0	2,2,2	0.53	0
3	CIT	B	303	-	12,12,12	1.05	0	17,17,17	1.41	3 (17%)
2	EDO	A	302	-	3,3,3	0.46	0	2,2,2	0.47	0
2	EDO	B	304	-	3,3,3	0.46	0	2,2,2	0.53	0
3	CIT	A	303	-	12,12,12	1.03	0	17,17,17	1.51	3 (17%)
3	CIT	A	304	-	12,12,12	1.01	0	17,17,17	1.57	3 (17%)
3	CIT	D	304	-	12,12,12	1.07	0	17,17,17	1.50	2 (11%)
2	EDO	B	301	-	3,3,3	0.52	0	2,2,2	0.43	0
2	EDO	C	304	-	3,3,3	0.44	0	2,2,2	0.49	0
2	EDO	A	301	-	3,3,3	0.53	0	2,2,2	0.39	0
4	GOL	A	305	-	5,5,5	0.81	0	5,5,5	1.09	0
3	CIT	D	303	-	12,12,12	1.03	0	17,17,17	1.56	3 (17%)
2	EDO	D	305	-	3,3,3	0.55	0	2,2,2	0.07	0
3	CIT	C	303	-	12,12,12	1.08	0	17,17,17	1.48	3 (17%)
2	EDO	A	306	-	3,3,3	0.47	0	2,2,2	0.35	0
2	EDO	B	302	-	3,3,3	0.45	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	301	-	-	0/1/1/1	-
2	EDO	D	302	-	-	0/1/1/1	-
2	EDO	D	307	-	-	1/1/1/1	-
2	EDO	C	301	-	-	0/1/1/1	-
2	EDO	C	305	-	-	1/1/1/1	-
4	GOL	D	306	-	-	2/4/4/4	-
2	EDO	C	302	-	-	0/1/1/1	-
3	CIT	B	303	-	-	7/16/16/16	-
2	EDO	A	302	-	-	0/1/1/1	-
2	EDO	B	304	-	-	1/1/1/1	-
3	CIT	A	303	-	-	6/16/16/16	-
3	CIT	A	304	-	-	3/16/16/16	-
3	CIT	D	304	-	-	5/16/16/16	-
2	EDO	B	301	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	C	304	-	-	0/1/1/1	-
2	EDO	A	301	-	-	0/1/1/1	-
4	GOL	A	305	-	-	4/4/4/4	-
3	CIT	D	303	-	-	6/16/16/16	-
2	EDO	D	305	-	-	0/1/1/1	-
3	CIT	C	303	-	-	8/16/16/16	-
2	EDO	A	306	-	-	0/1/1/1	-
2	EDO	B	302	-	-	0/1/1/1	-

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	304	CIT	O6-C6-C3	4.18	121.15	113.14
3	D	303	CIT	O6-C6-C3	3.90	120.62	113.14
3	A	303	CIT	O6-C6-C3	3.84	120.51	113.14
3	A	304	CIT	O6-C6-C3	3.59	120.03	113.14
3	A	304	CIT	O4-C5-C4	3.25	124.64	114.35
3	B	303	CIT	O6-C6-C3	3.12	119.12	113.14
3	C	303	CIT	O6-C6-C3	2.99	118.87	113.14
3	D	303	CIT	O4-C5-O3	-2.58	116.69	123.33
3	C	303	CIT	O4-C5-C4	2.56	122.45	114.35
3	A	303	CIT	O4-C5-C4	2.33	121.72	114.35
3	C	303	CIT	O4-C5-O3	-2.28	117.46	123.33
3	B	303	CIT	O4-C5-C4	2.27	121.54	114.35
3	A	303	CIT	O4-C5-O3	-2.24	117.58	123.33
3	D	304	CIT	O4-C5-O3	-2.17	117.76	123.33
3	D	303	CIT	O4-C5-C4	2.07	120.90	114.35
4	D	306	GOL	C3-C2-C1	-2.07	104.21	111.80
3	A	304	CIT	O3-C5-C4	-2.06	117.12	122.95
3	B	303	CIT	O4-C5-O3	-2.01	118.17	123.33

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	303	CIT	O7-C3-C6-O5
3	A	303	CIT	O7-C3-C6-O6
3	A	303	CIT	C4-C3-C6-O5
3	A	303	CIT	C4-C3-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	303	CIT	O7-C3-C6-O5
3	B	303	CIT	O7-C3-C6-O6
3	B	303	CIT	C4-C3-C6-O5
3	B	303	CIT	C4-C3-C6-O6
3	C	303	CIT	O7-C3-C6-O5
3	C	303	CIT	O7-C3-C6-O6
3	C	303	CIT	C4-C3-C6-O5
3	C	303	CIT	C4-C3-C6-O6
3	D	303	CIT	O7-C3-C6-O5
3	D	303	CIT	O7-C3-C6-O6
3	D	303	CIT	C4-C3-C6-O5
3	D	303	CIT	C4-C3-C6-O6
3	D	304	CIT	C1-C2-C3-O7
3	D	304	CIT	C1-C2-C3-C6
4	A	305	GOL	O1-C1-C2-C3
4	D	306	GOL	O1-C1-C2-C3
4	A	305	GOL	O1-C1-C2-O2
4	D	306	GOL	O1-C1-C2-O2
3	D	304	CIT	C1-C2-C3-C4
4	A	305	GOL	C1-C2-C3-O3
4	A	305	GOL	O2-C2-C3-O3
2	C	305	EDO	O1-C1-C2-O2
2	B	304	EDO	O1-C1-C2-O2
2	D	307	EDO	O1-C1-C2-O2
3	B	303	CIT	C3-C4-C5-O4
3	C	303	CIT	C3-C4-C5-O4
3	A	303	CIT	C3-C4-C5-O4
3	B	303	CIT	C3-C4-C5-O3
3	C	303	CIT	C3-C4-C5-O3
3	A	303	CIT	C3-C4-C5-O3
3	D	304	CIT	C2-C3-C4-C5
3	D	303	CIT	C2-C3-C6-O5
3	D	303	CIT	C2-C3-C6-O6
3	A	304	CIT	C3-C4-C5-O4
3	A	304	CIT	C2-C3-C4-C5
3	A	304	CIT	C3-C4-C5-O3
3	D	304	CIT	C6-C3-C4-C5
3	C	303	CIT	O1-C1-C2-C3
3	B	303	CIT	O1-C1-C2-C3
3	C	303	CIT	O2-C1-C2-C3

There are no ring outliers.

8 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	307	EDO	1	0
2	C	301	EDO	1	0
2	C	305	EDO	1	0
4	D	306	GOL	1	0
3	A	303	CIT	1	0
3	A	304	CIT	1	0
4	A	305	GOL	1	0
2	A	306	EDO	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	193/228 (84%)	-0.10	5 (2%)	56	65	18, 27, 47, 97	0
1	B	191/228 (83%)	0.19	11 (5%)	23	31	20, 30, 66, 106	0
1	C	190/228 (83%)	0.28	14 (7%)	14	22	21, 34, 68, 95	0
1	D	193/228 (84%)	-0.03	5 (2%)	56	65	18, 27, 48, 87	0
All	All	767/912 (84%)	0.08	35 (4%)	32	42	18, 29, 61, 106	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	171	PHE	9.5
1	B	170	GLY	5.7
1	C	173	SER	5.4
1	B	172	LEU	5.3
1	B	173	SER	4.9
1	C	170	GLY	4.8
1	A	136	HIS	4.0
1	B	174	HIS	3.9
1	C	168	ASP	3.5
1	C	174	HIS	3.4
1	C	167	TYR	3.3
1	D	10	LYS	3.3
1	C	176	ASP	3.3
1	A	139	PHE	3.2
1	D	139	PHE	3.2
1	C	175	GLU	3.1
1	B	176	ASP	3.1
1	B	169	ASP	3.0
1	A	138	ASP	3.0
1	D	136	HIS	3.0
1	A	10	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	165	GLY	2.7
1	C	166	LEU	2.7
1	B	168	ASP	2.6
1	B	166	LEU	2.5
1	B	167	TYR	2.5
1	C	169	ASP	2.5
1	D	138	ASP	2.5
1	B	175	GLU	2.5
1	A	11	LYS	2.4
1	C	77	ASN	2.4
1	C	76	SER	2.3
1	C	11	LYS	2.3
1	C	178	ASP	2.2
1	D	202	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	EDO	C	305	4/4	0.69	0.20	72,73,73,75	0
2	EDO	C	304	4/4	0.81	0.16	51,54,58,66	0
2	EDO	D	307	4/4	0.81	0.26	60,70,70,74	0
2	EDO	A	306	4/4	0.82	0.21	63,65,66,69	0
3	CIT	A	303	13/13	0.84	0.29	44,57,103,105	0
2	EDO	D	305	4/4	0.85	0.17	37,53,56,57	0
3	CIT	B	303	13/13	0.86	0.19	36,57,109,109	0
2	EDO	B	304	4/4	0.87	0.14	46,50,58,67	0
3	CIT	C	303	13/13	0.87	0.26	39,62,107,108	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CIT	A	304	13/13	0.88	0.27	41,89,104,107	0
3	CIT	D	303	13/13	0.90	0.30	33,53,111,112	0
3	CIT	D	304	13/13	0.90	0.21	39,92,105,105	0
4	GOL	A	305	6/6	0.90	0.15	38,43,50,51	0
2	EDO	A	302	4/4	0.92	0.20	33,35,36,42	0
4	GOL	D	306	6/6	0.92	0.13	38,43,46,50	0
2	EDO	C	302	4/4	0.93	0.17	33,43,44,47	0
2	EDO	C	301	4/4	0.94	0.26	34,35,37,39	0
2	EDO	D	302	4/4	0.94	0.10	29,35,35,36	0
2	EDO	B	302	4/4	0.94	0.16	32,35,37,42	0
2	EDO	B	301	4/4	0.94	0.18	24,34,34,43	0
2	EDO	A	301	4/4	0.97	0.12	22,24,25,28	0
2	EDO	D	301	4/4	0.98	0.14	22,23,23,27	0

6.5 Other polymers (i)

There are no such residues in this entry.