Full wwPDB X-ray Structure Validation Report (i)

Dec 16, 2023 – 04:22 PM EST

PDB ID	:	4P8S
Title	:	Crystal structure of Nogo-receptor-2
Authors	:	Semavina, M.; Saha, N.; Kolev, M.V.; Giger, R.J.; Himanen, J.P.; Nikolov, D.B.
Deposited on	:	2014-04-01
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	2%, 78%, 17%, • •
2	B	2	50%, 50%
3	C	2	50%, 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	405	X	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticulon-4 receptor-like 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	282	2249	1425	412	402	10	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	365	365	365	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reticulon-4 receptor-like 2

Chain A:



• Molecule 2: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	56.82Å 56.52Å 128.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.80 25.30 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.3 (25.00-1.80) 98.5 (25.30-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ [1]	2.17 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.171 , 0.205 0.169 , 0.203	Depositor DCC
R_{free} test set	1959 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	20.2	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 46.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2684	wwPDB-VP
Average B, all atoms (Å^2)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.42	13/2305 (0.6%)	1.21	18/3137 (0.6%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	47	GLN	CB-CG	13.46	1.89	1.52
1	A	245	GLU	CB-CG	6.84	1.65	1.52
1	A	35	CYS	CB-SG	6.66	1.93	1.82
1	A	265	CYS	CB-SG	6.37	1.93	1.82
1	A	53	SER	CB-OG	6.06	1.50	1.42
1	A	255	PHE	CD1-CE1	5.87	1.50	1.39
1	A	102	PHE	CE1-CZ	5.81	1.48	1.37
1	A	141	CYS	CB-SG	5.46	1.91	1.82
1	A	282	SER	CB-OG	-5.36	1.35	1.42
1	A	288	CYS	CB-SG	-5.33	1.73	1.81
1	A	220	ARG	CG-CD	5.25	1.65	1.51
1	A	284	SER	CB-OG	5.20	1.49	1.42
1	A	254	GLU	CG-CD	-5.09	1.44	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	73	ARG	NE-CZ-NH1	7.74	124.17	120.30
1	A	73	ARG	NE-CZ-NH2	-6.55	117.03	120.30
1	A	209	LEU	CB-CG-CD1	6.49	122.04	111.00
1	A	207	ARG	CG-CD-NE	-6.31	98.55	111.80
1	A	303	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	A	172	ASP	CB-CG-OD1	5.97	123.67	118.30
1	A	253	LEU	CB-CG-CD2	-5.73	101.26	111.00
1	A	152	ARG	NE-CZ-NH1	5.72	123.16	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	A	167	LEU	CB-CG-CD1	-5.64	101.41	111.00
1	A	250	LEU	CB-CG-CD2	-5.53	101.61	111.00
1	A	152	ARG	NE-CZ-NH2	-5.51	117.54	120.30
1	A	119	LEU	CB-CG-CD2	-5.42	101.79	111.00
1	A	242	LEU	CA-CB-CG	5.40	127.73	115.30
1	A	207	ARG	NE-CZ-NH1	-5.28	117.66	120.30
1	A	37	CYS	CA-CB-SG	-5.16	104.71	114.00
1	A	268	ARG	CG-CD-NE	5.12	122.55	111.80
1	A	34	LEU	CB-CG-CD1	5.08	119.64	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2249	0	2222	59	0
2	B	28	0	24	1	0
3	C	28	0	25	1	0
4	A	14	0	13	1	0
5	A	365	0	0	14	4
All	All	2684	0	2284	62	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:GLN:CB	1:A:47:GLN:CG	1.88	1.49
1:A:169:HIS:HB3	5:A:569:HOH:O	1.27	1.31
1:A:117:ARG:HG3	5:A:732:HOH:O	1.52	1.08
1:A:117:ARG:CG	5:A:732:HOH:O	2.06	1.00
5:A:779:HOH:O	2:B:1:NAG:H82	1.64	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:ARG:HH11	1:A:268:ARG:HG3	1.35	0.92
1:A:303:ARG:HH11	1:A:303:ARG:HG2	1.34	0.90
1:A:47:GLN:CG	1:A:47:GLN:H	1.87	0.87
1:A:68:GLN:H	1:A:68:GLN:HE21	1.17	0.87
1:A:220:ARG:CD	1:A:220:ARG:H	1.95	0.78
1:A:70:ASN:HB2	1:A:92:ASN:HD21	1.49	0.77
1:A:72:ILE:H	1:A:92:ASN:HD22	1.33	0.75
1:A:239:LEU:H	1:A:261:ASN:HD22	1.37	0.73
1:A:237:ASN:HB2	1:A:261:ASN:HD21	1.55	0.72
1:A:215:LEU:H	1:A:237:ASN:HD22	1.39	0.71
1:A:47:GLN:H	1:A:47:GLN:HG3	1.56	0.69
1:A:47:GLN:CG	1:A:47:GLN:N	2.56	0.68
1:A:213:ASN:HB2	1:A:237:ASN:HD21	1.59	0.66
1:A:266:ASP:OD2	1:A:268:ARG:NH1	2.29	0.66
1:A:137:HIS:HD2	5:A:838:HOH:O	1.78	0.65
1:A:266:ASP:OD1	1:A:268:ARG:NH1	2.30	0.65
1:A:141:CYS:H	1:A:165:ASN:HD21	1.45	0.64
1:A:115:ASP:HB2	5:A:755:HOH:O	1.97	0.64
1:A:303:ARG:HH11	1:A:303:ARG:CG	2.08	0.64
1:A:68:GLN:H	1:A:68:GLN:NE2	1.94	0.61
1:A:51:PHE:H	1:A:70:ASN:HD22	1.49	0.61
1:A:207:ARG:NH1	5:A:698:HOH:O	2.13	0.60
1:A:137:HIS:CD2	5:A:838:HOH:O	2.52	0.59
1:A:268:ARG:HH11	1:A:268:ARG:CG	2.09	0.59
1:A:268:ARG:HG3	1:A:268:ARG:NH1	2.13	0.57
1:A:49:ASN:H	1:A:70:ASN:HD21	1.53	0.57
1:A:94:LEU:H	1:A:116:ASN:HD22	1.52	0.57
1:A:215:LEU:H	1:A:237:ASN:ND2	2.04	0.56
1:A:141:CYS:H	1:A:165:ASN:ND2	2.03	0.56
1:A:245:GLU:O	1:A:245:GLU:HG2	2.06	0.55
1:A:220:ARG:H	1:A:220:ARG:HD2	1.69	0.54
1:A:239:LEU:H	1:A:261:ASN:ND2	2.05	0.54
1:A:47:GLN:CG	1:A:47:GLN:CA	2.84	0.54
1:A:143:LEU:H	1:A:165:ASN:HD22	1.56	0.53
1:A:266:ASP:CG	1:A:268:ARG:NH1	2.63	0.52
1:A:295:GLN:NE2	1:A:295:GLN:C	2.63	0.52
1:A:49:ASN:H	1:A:70:ASN:ND2	2.08	0.51
1:A:94:LEU:H	1:A:116:ASN:ND2	2.10	0.50
4:A:405:NAG:H82	5:A:596:HOH:O	2.10	0.50
1:A:266:ASP:CG	1:A:268:ARG:HH12	2.13	0.49
1:A:47:GLN:CB	1:A:47:GLN:CD	2.72	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASP:HA	1:A:137:HIS:HB2	1.94	0.49
1:A:119:LEU:O	5:A:748:HOH:O	2.20	0.49
1:A:51:PHE:H	1:A:70:ASN:ND2	2.12	0.47
1:A:303:ARG:CG	1:A:303:ARG:NH1	2.75	0.47
1:A:92:ASN:HB2	1:A:116:ASN:HD21	1.80	0.47
1:A:237:ASN:CB	1:A:261:ASN:HD21	2.24	0.45
1:A:265:CYS:HB3	1:A:302:LEU:HD11	1.99	0.45
1:A:245:GLU:CG	5:A:721:HOH:O	2.66	0.44
1:A:33:MET:HG3	1:A:34:LEU:HD13	1.99	0.43
1:A:117:ARG:HG2	5:A:732:HOH:O	1.99	0.43
1:A:295:GLN:NE2	1:A:295:GLN:O	2.52	0.43
5:A:855:HOH:O	3:C:2:NAG:H61	2.17	0.42
1:A:32:PRO:HG2	1:A:58:LEU:HD23	2.00	0.42
1:A:226:LEU:HD23	1:A:226:LEU:HA	1.88	0.41
1:A:294:ARG:NH2	1:A:306:ASP:HB3	2.35	0.41
1:A:245:GLU:HG3	5:A:721:HOH:O	2.20	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:505:HOH:O	5:A:568:HOH:O[4_455]	1.45	0.75
5:A:556:HOH:O	5:A:575:HOH:O[3_645]	1.83	0.37
5:A:522:HOH:O	5:A:579:HOH:O[3_655]	2.12	0.08
5:A:553:HOH:O	5:A:559:HOH:O[3_655]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	280/282 (99%)	267 (95%)	13 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	250/250 (100%)	238 (95%)	12 (5%)	25 11

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	SER
1	A	34	LEU
1	A	47	GLN
1	A	68	GLN
1	A	209	LEU
1	A	214	ARG
1	A	220	ARG
1	A	268	ARG
1	A	270	ARG
1	A	295	GLN
1	A	303	ARG
1	A	304	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	70	ASN
1	A	92	ASN
1	A	116	ASN
1	A	137	HIS
1	A	165	ASN
1	A	180	ASN
1	A	237	ASN
1	A	261	ASN
1	A	295	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	B	1	2,1	14,14,15	0.76	0	17,19,21	2.19	5 (29%)
2	NDG	B	2	2	14,14,15	0.91	0	17,19,21	2.55	5 (29%)
3	NAG	C	1	3,1	14,14,15	1.26	2 (14%)	17,19,21	1.58	2 (11%)
3	NAG	C	2	3	14,14,15	0.71	0	17,19,21	1.63	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NDG	B	2	2	-	1/6/23/26	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	NAG	C1-C2	2.49	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	NAG	C4-C5	2.14	1.57	1.53

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	2	NDG	O5-C5-C6	6.43	117.28	107.20
2	B	2	NDG	C1-C2-N2	-5.10	101.78	110.49
2	B	1	NAG	O5-C1-C2	-4.49	104.20	111.29
3	C	1	NAG	O5-C1-C2	-4.22	104.62	111.29
2	B	1	NAG	C1-O5-C5	4.16	117.82	112.19
2	B	2	NDG	C4-C3-C2	3.79	116.57	111.02
2	B	1	NAG	O6-C6-C5	-3.74	98.46	111.29
2	B	1	NAG	C2-N2-C7	-3.38	118.09	122.90
3	C	2	NAG	C3-C4-C5	3.22	115.99	110.24
3	C	2	NAG	O3-C3-C2	-3.07	103.11	109.47
2	B	2	NDG	C3-C4-C5	3.03	115.65	110.24
3	C	2	NAG	O5-C1-C2	-2.78	106.90	111.29
2	B	2	NDG	C1-O5-C5	2.78	115.96	112.19
3	C	1	NAG	C4-C3-C2	-2.49	107.37	111.02
3	C	2	NAG	O6-C6-C5	-2.07	104.19	111.29
2	B	1	NAG	O3-C3-C2	-2.05	105.22	109.47

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	2	NDG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0
3	C	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain B



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain C



5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	A	405	1	14,14,15	1.28	2 (14%)	17,19,21	3.32	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	405	1	1/1/5/7	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	NAG	C1-C2	3.01	1.56	1.52
4	A	405	NAG	O5-C1	2.31	1.47	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	405	NAG	O5-C5-C6	7.51	118.98	107.20
4	A	405	NAG	O5-C1-C2	6.00	120.77	111.29
4	A	405	NAG	C2-N2-C7	5.31	130.47	122.90
4	A	405	NAG	C3-C4-C5	-5.03	101.27	110.24
4	A	405	NAG	O4-C4-C5	3.09	116.98	109.30
4	A	405	NAG	C1-C2-N2	-3.00	105.36	110.49
4	A	405	NAG	C6-C5-C4	2.62	119.13	113.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	405	NAG	C1

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	282/282 (100%)	-0.18	7 (2%) 57 52	10, 19, 37, 54	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	29	PRO	5.6
1	A	278	ARG	3.5
1	A	280	ARG	3.1
1	A	30	SER	3.0
1	A	304	ASP	2.6
1	A	303	ARG	2.5
1	A	117	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NDG	B	2	14/15	0.68	0.36	68,70,72,72	0
3	NAG	C	2	14/15	0.84	0.31	39,43,50,54	0
2	NAG	B	1	14/15	0.85	0.27	36,45,54,62	0
3	NAG	C	1	14/15	0.89	0.21	27,32,36,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NAG	A	405	14/15	0.30	0.46	59,69,71,73	0

6.5 Other polymers (i)

There are no such residues in this entry.