



## wwPDB EM Validation Summary Report ⓘ

Aug 28, 2025 – 02:53 PM EDT

PDB ID : 9PDY / pdb\_00009pdy  
EMDB ID : EMD-71548  
Title : Human antibody Fab MPV510 bound to hMPV DsCav-ES2-IPDS F protein  
Authors : Ghazi Esfahani, N.; Mousa, J.; Khalil, A.M.  
Deposited on : 2025-07-01  
Resolution : 2.85 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev126  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : **FAILED**  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.85 Å.

There are no overall percentile quality scores available for this entry.

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 25087 atoms, of which 9898 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F0.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	431	Total	C	H	N	O	S	0	0
			6543	2047	3273	561	631	31		
1	B	431	Total	C	H	N	O	S	0	0
			6543	2047	3273	561	631	31		
1	C	431	Total	C	H	N	O	S	0	0
			6538	2047	3268	561	631	31		

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	84	CYS	VAL	conflict	UNP H6X1Z0
A	110	CYS	LEU	conflict	UNP H6X1Z0
A	127	CYS	THR	conflict	UNP H6X1Z0
A	140	CYS	ALA	conflict	UNP H6X1Z0
A	147	CYS	ALA	conflict	UNP H6X1Z0
A	153	CYS	ASN	conflict	UNP H6X1Z0
A	185	PRO	ALA	conflict	UNP H6X1Z0
A	219	LYS	LEU	conflict	UNP H6X1Z0
A	231	ILE	VAL	conflict	UNP H6X1Z0
A	249	CYS	ALA	conflict	UNP H6X1Z0
A	322	CYS	ASN	conflict	UNP H6X1Z0
A	365	CYS	THR	conflict	UNP H6X1Z0
A	453	GLN	GLU	conflict	UNP H6X1Z0
A	463	CYS	VAL	conflict	UNP H6X1Z0
B	84	CYS	VAL	conflict	UNP H6X1Z0
B	110	CYS	LEU	conflict	UNP H6X1Z0
B	127	CYS	THR	conflict	UNP H6X1Z0
B	140	CYS	ALA	conflict	UNP H6X1Z0
B	147	CYS	ALA	conflict	UNP H6X1Z0
B	153	CYS	ASN	conflict	UNP H6X1Z0
B	185	PRO	ALA	conflict	UNP H6X1Z0
B	219	LYS	LEU	conflict	UNP H6X1Z0
B	231	ILE	VAL	conflict	UNP H6X1Z0
B	249	CYS	ALA	conflict	UNP H6X1Z0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	322	CYS	ASN	conflict	UNP H6X1Z0
B	365	CYS	THR	conflict	UNP H6X1Z0
B	453	GLN	GLU	conflict	UNP H6X1Z0
B	463	CYS	VAL	conflict	UNP H6X1Z0
C	84	CYS	VAL	conflict	UNP H6X1Z0
C	110	CYS	LEU	conflict	UNP H6X1Z0
C	127	CYS	THR	conflict	UNP H6X1Z0
C	140	CYS	ALA	conflict	UNP H6X1Z0
C	147	CYS	ALA	conflict	UNP H6X1Z0
C	153	CYS	ASN	conflict	UNP H6X1Z0
C	185	PRO	ALA	conflict	UNP H6X1Z0
C	219	LYS	LEU	conflict	UNP H6X1Z0
C	231	ILE	VAL	conflict	UNP H6X1Z0
C	249	CYS	ALA	conflict	UNP H6X1Z0
C	322	CYS	ASN	conflict	UNP H6X1Z0
C	365	CYS	THR	conflict	UNP H6X1Z0
C	453	GLN	GLU	conflict	UNP H6X1Z0
C	463	CYS	VAL	conflict	UNP H6X1Z0


- Molecule 2 is a protein called MPV510 Fab heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	121	Total	C	N	O	S	0	0
			937	588	157	190	2		
2	E	121	Total	C	N	O	S	0	0
			937	588	157	190	2		
2	F	121	Total	C	N	O	S	0	0
			937	588	157	190	2		

- Molecule 3 is a protein called MPV510 Fab light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	107	Total	C	N	O	S	0	0
			828	518	144	162	4		
3	H	107	Total	C	N	O	S	0	0
			828	518	144	162	4		
3	I	107	Total	C	N	O	S	0	0
			828	518	144	162	4		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	H	N	O	0
			28	8	14	1	5	
4	A	1	Total	C	H	N	O	0
			28	8	14	1	5	
4	B	1	Total	C	H	N	O	0
			28	8	14	1	5	
4	B	1	Total	C	H	N	O	0
			28	8	14	1	5	
4	C	1	Total	C	H	N	O	0
			28	8	14	1	5	
4	C	1	Total	C	H	N	O	0
			28	8	14	1	5	

MolProbity failed to run properly - this section is therefore empty.

### 3 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	287000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	60	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	Not provided	
Image detector	DIRECT ELECTRON APOLLO (4k x 4k)	Depositor
Maximum map value	0.728	Depositor
Minimum map value	-0.501	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.03	Depositor
Map size ( $\text{\AA}$ )	221.0, 221.0, 221.0	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	0.7892857, 0.7892857, 0.7892857	Depositor

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	B	601	1	14,14,15	0.73	0	17,19,21	0.88	0
4	NAG	C	601	1	14,14,15	0.76	0	17,19,21	0.89	0
4	NAG	A	601	1	14,14,15	0.74	0	17,19,21	0.88	0
4	NAG	A	602	1	14,14,15	0.70	0	17,19,21	0.84	0
4	NAG	C	602	1	14,14,15	0.71	0	17,19,21	0.84	0
4	NAG	B	602	1	14,14,15	0.72	0	17,19,21	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	601	1	-	1/6/23/26	0/1/1/1
4	NAG	C	601	1	-	1/6/23/26	0/1/1/1
4	NAG	A	601	1	-	1/6/23/26	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
4	NAG	C	602	1	-	0/6/23/26	0/1/1/1
4	NAG	B	602	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	NAG	O5-C5-C6-O6
4	B	601	NAG	O5-C5-C6-O6
4	C	601	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 4.7 Other polymers

There are no such residues in this entry.



## 4.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


## 5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-71548. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 5.1 Orthogonal projections [i](#)


#### 5.1.1 Primary map



X




Y




Z


#### 5.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 5.2 Central slices [i](#)


### 5.2.1 Primary map



X Index: 140




Y Index: 140




Z Index: 140


### 5.2.2 Raw map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

## 5.3 Largest variance slices [i](#)


### 5.3.1 Primary map



X Index: 152




Y Index: 149




Z Index: 150


### 5.3.2 Raw map



X Index: 152



Y Index: 149




Z Index: 150


The images above show the largest variance slices of the map in three orthogonal directions.

## 5.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 5.4.1 Primary map



X




Y




Z


### 5.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 5.5 Orthogonal surface views [i](#)

### 5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.03. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 5.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 6.2 Volume estimate [i](#)




The volume at the recommended contour level is 316 nm<sup>3</sup>; this corresponds to an approximate mass of 285 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 6.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.351  $\text{\AA}^{-1}$

## 7 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 7.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.351 \text{ \AA}^{-1}$

## 7.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.85	-	-
Author-provided FSC curve	2.85	3.31	2.96
Unmasked-calculated*	3.34	3.82	3.38

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.34 differs from the reported value 2.85 by more than 10 %

## 8 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-71548 and PDB model 9PDY. Per-residue inclusion information can be found in section ?? on page ??.

### 8.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).

## 8.4 Atom inclusion [i](#)



At the recommended contour level, 89% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8870	<div></div> 0.4300
A	<div></div> 0.9410	<div></div> 0.4760
B	<div></div> 0.9390	<div></div> 0.4750
C	<div></div> 0.9410	<div></div> 0.4800
D	<div></div> 0.8320	<div></div> 0.3720
E	<div></div> 0.8240	<div></div> 0.3480
F	<div></div> 0.8230	<div></div> 0.3360
G	<div></div> 0.7810	<div></div> 0.3430
H	<div></div> 0.7740	<div></div> 0.3390
I	<div></div> 0.7560	<div></div> 0.3110

1.0

0.0

<0.0