Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 06:19 AM EDT

PDB ID	:	6PUA
Title	:	The 2.0 A Crystal Structure of the Type B Chloramphenicol Acetyltransferase from Vibrio cholerae
Authors	:	Kim, Y.; Maltseva, N.; Stam, J.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on	:	2019-07-18
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5475 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chloramphenicol acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	210	1693	1080	288	314	11	0	3	0
1	B	210	1665	1064	280	310	11	0	0	0
1	C	206	1650	1055	275	309	11	0	2	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9KMN1
A	-1	ASN	-	expression tag	UNP Q9KMN1
A	0	ALA	-	expression tag	UNP Q9KMN1
B	-2	SER	-	expression tag	UNP Q9KMN1
B	-1	ASN	-	expression tag	UNP Q9KMN1
B	0	ALA	-	expression tag	UNP Q9KMN1
C	-2	SER	-	expression tag	UNP Q9KMN1
C	-1	ASN	-	expression tag	UNP Q9KMN1
C	0	ALA	-	expression tag	UNP Q9KMN1

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	8	6	2	0	0
			Total	C	O		
2	A	1	8	6	2	0	0
			Total	C	O		
2	B	1	8	6	2	0	0
			Total	C	O		
2	B	1	8	6	2	0	0
			Total	C	O		
2	C	1	8	6	2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
3	A	1	4	2	2		0	0
			Total	C	O			
3	A	1	4	2	2		0	0
			Total	C	O			
3	A	1	4	2	2		0	0
			Total	C	O			
3	A	1	4	2	2		0	0
			Total	C	O			
3	A	1	4	2	2		0	0
			Total	C	O			
3	B	1	4	2	2		0	0
			Total	C	O			
3	B	1	4	2	2		0	0
			Total	C	O			
3	B	1	4	2	2		0	0
			Total	C	O			
3	C	1	4	2	2		0	0
			Total	C	O			
3	C	1	4	2	2		0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Cl			
4	A	1	1	1		0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
5	A	1	5	4	1	0	0
			Total	O	P		
5	B	1	5	4	1	0	0
			Total	O	P		
5	B	1	5	4	1	0	0
			Total	O	P		
5	C	1	5	4	1	0	0
			Total	O	P		
5	C	1	5	4	1	0	0
			Total	O	P		
5	C	1	5	4	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	145	145	145	0	0
			Total	O		
6	B	103	103	103	0	0
			Total	O		
6	C	108	108	108	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α, β, γ	100.06Å 100.06Å 128.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.42 – 2.00	Depositor
% Data completeness (in resolution range)	99.5 (39.42-2.00)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.51 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.163 , 0.188	Depositor
Wilson B-factor (Å^2)	37.4	Xtriage
Anisotropy	0.190	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtriage
Total number of atoms	5475	wwPDB-VP
Average B, all atoms (Å^2)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.78% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 1 is monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	EDO	A	302	-	3,3,3	0.51	0	2,2,2	0.15	0
3	EDO	B	302	-	3,3,3	0.49	0	2,2,2	0.25	0
2	MPD	C	304	-	7,7,7	0.57	0	9,10,10	0.75	0
3	EDO	B	301	-	3,3,3	0.45	0	2,2,2	0.37	0
3	EDO	B	303	-	3,3,3	0.50	0	2,2,2	0.32	0
2	MPD	A	307	-	7,7,7	0.37	0	9,10,10	1.17	1 (11%)
5	PO4	B	306	-	4,4,4	0.86	0	6,6,6	0.53	0
5	PO4	C	302	-	4,4,4	0.92	0	6,6,6	0.39	0
3	EDO	A	303	-	3,3,3	0.55	0	2,2,2	0.16	0
5	PO4	A	309	-	4,4,4	0.84	0	6,6,6	0.48	0
3	EDO	A	305	-	3,3,3	0.48	0	2,2,2	0.31	0
2	MPD	A	301	-	7,7,7	0.23	0	9,10,10	0.40	0
5	PO4	C	306	-	4,4,4	0.87	0	6,6,6	0.48	0
3	EDO	A	304	-	3,3,3	0.51	0	2,2,2	0.20	0
5	PO4	B	304	-	4,4,4	0.88	0	6,6,6	0.35	0
2	MPD	B	305	-	7,7,7	0.30	0	9,10,10	0.53	0
3	EDO	C	303	-	3,3,3	0.45	0	2,2,2	0.43	0
3	EDO	A	306	-	3,3,3	0.48	0	2,2,2	0.32	0
2	MPD	B	307	-	7,7,7	0.27	0	9,10,10	0.23	0
5	PO4	C	301	-	4,4,4	0.89	0	6,6,6	0.49	0
3	EDO	C	305	-	3,3,3	0.47	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	B	305	-	-	3/5/5/5	-
3	EDO	A	304	-	-	0/1/1/1	-
3	EDO	B	303	-	-	0/1/1/1	-
3	EDO	C	303	-	-	0/1/1/1	-
3	EDO	A	302	-	-	0/1/1/1	-
3	EDO	A	306	-	-	0/1/1/1	-
3	EDO	C	305	-	-	1/1/1/1	-
3	EDO	B	302	-	-	1/1/1/1	-
2	MPD	C	304	-	-	2/5/5/5	-
2	MPD	A	307	-	-	3/5/5/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	B	307	-	-	0/5/5/5	-
3	EDO	A	303	-	-	0/1/1/1	-
3	EDO	B	301	-	-	1/1/1/1	-
3	EDO	A	305	-	-	1/1/1/1	-
2	MPD	A	301	-	-	0/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	307	MPD	CM-C2-C1	-2.73	104.89	110.57

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	EDO	O1-C1-C2-O2
2	A	307	MPD	O2-C2-C3-C4
2	B	305	MPD	O2-C2-C3-C4
2	C	304	MPD	C2-C3-C4-O4
2	A	307	MPD	C1-C2-C3-C4
2	A	307	MPD	CM-C2-C3-C4
2	B	305	MPD	C1-C2-C3-C4
2	B	305	MPD	CM-C2-C3-C4
3	A	305	EDO	O1-C1-C2-O2
3	B	302	EDO	O1-C1-C2-O2
3	C	305	EDO	O1-C1-C2-O2
2	C	304	MPD	C2-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.