



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 12:19 PM EDT

PDB ID : 4PYS
Title : The crystal structure of beta-N-acetylhexosaminidase from Bacteroides fragilis NCTC 9343
Authors : Tan, K.; Hatzos-Skintges, C.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2014-03-27
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	506	<div> <div>8%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>
1	B	506	<div> <div>9%</div> <div>86%</div> <div>9%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	A	607	-	-	X	-
4	FMT	B	608	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-N-acetylhexosaminidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	492	Total	C	N	O	S	Se	0	1	0
			3972	2564	675	713	8	12			
1	B	483	Total	C	N	O	S	Se	0	4	0
			3909	2517	674	698	8	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	ASN	TYR	CONFLICT	UNP Q5LAT3
A	15	ALA	SER	CONFLICT	UNP Q5LAT3
B	14	ASN	TYR	CONFLICT	UNP Q5LAT3
B	15	ALA	SER	CONFLICT	UNP Q5LAT3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 3 1 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	B	1	Total C O 3 1 2	0	0
4	B	1	Total C O 3 1 2	0	0
4	B	1	Total C O 3 1 2	0	0
4	B	1	Total C O 3 1 2	0	0
4	B	1	Total C O 3 1 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	280	Total O 280 280	0	0
5	B	324	Total O 324 324	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: beta-N-acetylhexosaminidase



- Molecule 1: beta-N-acetylhexosaminidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.82Å 96.82Å 315.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.03 – 1.82 37.97 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.6 (34.03-1.82) 99.6 (37.97-1.82)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 1.82Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.165 , 0.190 0.164 , 0.188	Depositor DCC
R_{free} test set	6739 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8553	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9169e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/4078	0.53	0/5512
1	B	0.35	0/4016	0.54	2/5423 (0.0%)
All	All	0.35	0/8094	0.54	2/10935 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	119	ARG	NE-CZ-NH1	-6.35	117.12	120.30
1	B	119	ARG	NE-CZ-NH2	5.75	123.18	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3972	0	3798	29	0
1	B	3909	0	3766	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	24	0	32	1	0
3	B	12	0	16	0	0
4	A	15	0	5	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	15	0	5	2	0
5	A	280	0	0	2	0
5	B	324	0	0	2	0
All	All	8553	0	7622	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:LEU:HD21	1:A:263:GLN:HA	1.69	0.74
1:B:388[A]:ARG:NH2	5:B:1022:HOH:O	2.23	0.71
1:A:21:GLN:HG3	1:A:22:ILE:H	1.61	0.64
1:B:203:VAL:HB	1:B:261:VAL:HG21	1.82	0.60
1:A:275:LYS:NZ	1:A:315:ASP:OD2	2.37	0.57
1:A:192:ARG:NH2	5:A:780:HOH:O	2.39	0.55
1:A:104:GLU:OE2	3:A:605:GOL:H12	2.06	0.54
1:A:44:ARG:HB3	1:A:71:CYS:SG	2.47	0.54
1:A:338:ARG:NH2	1:A:362:TYR:OH	2.40	0.54
1:B:192[A]:ARG:HD3	1:B:193:TYR:CZ	2.43	0.53
1:B:148:MSE:HG3	1:B:177:HIS:CG	2.44	0.53
1:A:20:GLY:N	1:A:121:TRP:O	2.41	0.52
1:A:148:MSE:HG3	1:A:177:HIS:CG	2.44	0.52
1:A:307:LYS:HD2	1:A:310:TRP:CE3	2.44	0.52
1:B:356:VAL:HG23	1:B:357:ILE:HG23	1.93	0.50
1:A:364:LEU:HB3	1:A:368:VAL:HG21	1.93	0.50
1:A:383:LEU:O	1:A:387:VAL:HG13	2.13	0.49
1:A:505:PRO:HD3	1:A:515:TYR:CZ	2.48	0.49
1:B:247:ASN:ND2	1:B:259:VAL:O	2.45	0.49
1:B:36:ARG:NH2	1:B:125:GLU:OE1	2.46	0.49
1:A:250:PRO:HB3	1:A:257:VAL:O	2.15	0.46
1:A:252:LEU:HD11	1:A:281:PHE:CZ	2.51	0.46
1:A:148:MSE:HE1	1:A:448:TRP:CD2	2.50	0.46
1:B:388[A]:ARG:NH1	5:B:920:HOH:O	2.47	0.46
1:A:252:LEU:HD21	1:A:281:PHE:CD2	2.51	0.46
1:B:444:SER:HB2	4:B:608:FMT:C	2.47	0.45
1:B:360:ASP:OD1	1:B:362:TYR:N	2.50	0.45
1:B:148:MSE:HE1	1:B:448:TRP:CD2	2.53	0.44
1:A:432:ARG:NH2	1:A:436:GLU:OE1	2.39	0.44
1:A:59:TYR:HA	1:A:63:VAL:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:307:LYS:HA	1:A:307:LYS:HD3	1.62	0.44
1:A:435:GLU:O	1:A:436:GLU:HB3	2.17	0.43
1:A:306:PRO:HD3	5:A:970:HOH:O	2.18	0.43
1:A:382:ALA:CB	4:A:607:FMT:H	2.49	0.43
1:B:177:HIS:ND1	4:B:608:FMT:O2	2.52	0.42
1:B:505:PRO:HD3	1:B:515:TYR:CZ	2.55	0.42
1:B:43:ILE:HD12	1:B:63:VAL:HG21	2.02	0.42
1:B:44:ARG:HB3	1:B:71:CYS:SG	2.60	0.41
1:B:59:TYR:HA	1:B:63:VAL:O	2.21	0.41
1:B:240:GLY:HA2	1:B:270:ILE:HG22	2.02	0.41
1:B:364:LEU:HB3	1:B:368:VAL:HG21	2.03	0.41
1:A:342:TYR:O	1:A:345:GLN:HG2	2.21	0.41
1:A:382:ALA:HB2	4:A:607:FMT:H	2.01	0.41
1:B:92:TYR:CE2	1:B:138:ASP:HB3	2.55	0.41
1:A:150:ASP:HA	1:A:179:HIS:HB3	2.03	0.41
1:A:192:ARG:NH2	1:B:503:PHE:H	2.19	0.41
1:A:255:PHE:HB2	1:A:257:VAL:HG13	2.02	0.41
1:B:82:LYS:O	1:B:104:GLU:HA	2.21	0.41
1:B:218:VAL:O	1:B:222:ILE:HG12	2.20	0.41
1:B:301:GLY:HA2	1:B:353:TRP:CD1	2.56	0.41
1:B:195:PHE:HB2	1:B:249:TYR:CE2	2.56	0.40
1:B:511:ALA:O	1:B:515:TYR:HB2	2.21	0.40
1:B:195:PHE:HB2	1:B:249:TYR:CZ	2.57	0.40
1:A:38:ARG:HH21	1:A:130:ILE:HD13	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	487/506 (96%)	471 (97%)	15 (3%)	1 (0%)	47 33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	475/506 (94%)	460 (97%)	14 (3%)	1 (0%)	47 33
All	All	962/1012 (95%)	931 (97%)	29 (3%)	2 (0%)	47 33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	435	GLU
1	B	314	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	405/425 (95%)	400 (99%)	5 (1%)	71 64
1	B	403/425 (95%)	396 (98%)	7 (2%)	60 50
All	All	808/850 (95%)	796 (98%)	12 (2%)	67 55

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	318	SER
1	A	338	ARG
1	A	341	ASP
1	A	450	ASP
1	A	513	THR
1	B	75	VAL
1	B	192[A]	ARG
1	B	192[B]	ARG
1	B	228	ARG
1	B	333	LEU
1	B	363	SER
1	B	450	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	478	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FMT	A	606	-	2,2,2	0.64	0	1,1,1	0.29	0
4	FMT	A	607	-	2,2,2	0.78	0	1,1,1	0.28	0
4	FMT	A	610	-	2,2,2	0.67	0	1,1,1	0.14	0
3	GOL	A	602	-	5,5,5	0.36	0	5,5,5	0.26	0
4	FMT	A	608	-	2,2,2	0.67	0	1,1,1	0.26	0
4	FMT	B	604	-	2,2,2	0.78	0	1,1,1	0.28	0
3	GOL	A	604	-	5,5,5	0.32	0	5,5,5	0.46	0
3	GOL	A	603	-	5,5,5	0.37	0	5,5,5	0.56	0
4	FMT	B	608	-	2,2,2	0.62	0	1,1,1	0.14	0
4	FMT	B	605	-	2,2,2	0.84	0	1,1,1	0.21	0
3	GOL	A	605	-	5,5,5	0.33	0	5,5,5	0.30	0
4	FMT	A	609	-	2,2,2	0.66	0	1,1,1	0.16	0
4	FMT	B	607	-	2,2,2	0.67	0	1,1,1	0.24	0
4	FMT	B	606	-	2,2,2	0.65	0	1,1,1	0.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	602	-	5,5,5	0.38	0	5,5,5	0.62	0
3	GOL	B	603	-	5,5,5	0.39	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	A	604	-	-	0/4/4/4	-
3	GOL	A	603	-	-	0/4/4/4	-
3	GOL	A	605	-	-	2/4/4/4	-
3	GOL	B	602	-	-	0/4/4/4	-
3	GOL	B	603	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	605	GOL	O1-C1-C2-C3
3	A	605	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	607	FMT	2	0
4	B	608	FMT	2	0
3	A	605	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	480/506 (94%)	0.17	39 (8%) 12 9	16, 34, 89, 118	0
1	B	471/506 (93%)	0.14	46 (9%) 7 5	16, 32, 81, 104	0
All	All	951/1012 (93%)	0.15	85 (8%) 9 7	16, 33, 85, 118	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	316	CYS	11.1
1	B	320	ILE	9.1
1	B	313	CYS	9.0
1	B	325	LEU	8.9
1	A	310	TRP	8.2
1	A	262	PRO	7.7
1	A	309	ASN	6.9
1	A	255	PHE	6.9
1	B	259	VAL	6.6
1	B	317	ARG	6.3
1	B	261	VAL	6.2
1	A	259	VAL	6.1
1	A	261	VAL	6.0
1	A	308	GLY	5.9
1	B	314	PRO	5.9
1	B	306	PRO	5.8
1	B	258	ALA	5.7
1	A	258	ALA	5.5
1	B	318	SER	5.4
1	A	513	THR	5.0
1	A	311	ASP	5.0
1	A	321	GLU	5.0
1	B	260	LYS	4.9
1	B	257	VAL	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	20	GLY	4.7
1	B	315	ASP	4.6
1	A	316	CYS	4.6
1	B	324	LYS	4.6
1	B	321	GLU	4.5
1	B	274	GLY	4.5
1	B	323	GLU	4.4
1	B	329	HIS	4.4
1	B	360	ASP	4.3
1	A	325	LEU	4.2
1	A	320	ILE	4.2
1	B	334	TRP	4.1
1	B	319	ARG	4.0
1	B	254	CYS	3.9
1	A	260	LYS	3.9
1	B	270	ILE	3.9
1	B	322	LYS	3.9
1	A	314	PRO	3.7
1	B	513	THR	3.6
1	A	263	GLN	3.6
1	B	273	ALA	3.5
1	A	254	CYS	3.5
1	B	328	SER	3.4
1	B	269	ASN	3.4
1	A	322	LYS	3.4
1	A	318	SER	3.4
1	B	330	ASP	3.3
1	A	313	CYS	3.3
1	B	307	LYS	3.2
1	A	312	LYS	3.2
1	A	327	ASP	3.1
1	B	331	LEU	3.1
1	A	326	LYS	3.1
1	B	479	PRO	3.0
1	B	256	ASN	3.0
1	A	362	TYR	2.9
1	B	127	ARG	2.7
1	A	330	ASP	2.7
1	B	125	GLU	2.7
1	A	257	VAL	2.6
1	B	272	CYS	2.6
1	A	125	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	273	ALA	2.6
1	B	304	GLU	2.5
1	B	34	ASP	2.5
1	A	329	HIS	2.5
1	B	333	LEU	2.4
1	B	512	GLY	2.4
1	B	305	ALA	2.4
1	A	361	GLY	2.4
1	B	126	GLU	2.3
1	B	478	ASN	2.3
1	A	34	ASP	2.2
1	A	126	GLU	2.2
1	A	466	ILE	2.2
1	A	128	GLY	2.1
1	A	302	GLY	2.1
1	B	38	ARG	2.1
1	B	446	ALA	2.1
1	A	274	GLY	2.1
1	A	323	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	FMT	A	610	3/3	0.77	0.16	32,32,48,50	0
3	GOL	A	605	6/6	0.80	0.29	44,51,62,64	0
4	FMT	A	609	3/3	0.86	0.20	39,39,41,46	0
3	GOL	A	603	6/6	0.87	0.14	32,42,46,49	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	FMT	B	606	3/3	0.87	0.20	43,43,44,52	0
4	FMT	A	608	3/3	0.89	0.16	47,47,53,66	0
4	FMT	B	608	3/3	0.90	0.21	40,40,44,45	0
4	FMT	B	607	3/3	0.91	0.21	38,38,44,52	0
3	GOL	B	602	6/6	0.91	0.16	29,46,52,52	0
2	ZN	A	601	1/1	0.94	0.11	66,66,66,66	1
2	ZN	B	601	1/1	0.94	0.14	91,91,91,91	1
3	GOL	A	602	6/6	0.94	0.09	28,29,33,35	0
3	GOL	B	603	6/6	0.96	0.09	29,33,36,36	0
3	GOL	A	604	6/6	0.97	0.07	26,31,33,34	0
4	FMT	A	607	3/3	0.97	0.11	21,21,35,36	0
4	FMT	B	605	3/3	0.98	0.10	22,22,30,35	0
4	FMT	B	604	3/3	0.98	0.08	24,24,31,33	0
4	FMT	A	606	3/3	0.99	0.07	22,22,33,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.