wwPDB X-ray Structure Validation Summary Report (i)

Jun 23, 2024 – 11:59 PM EDT

PDB ID	:	6QAV
Title	:	Crystal structure of ULK2 in complexed with MRT68921
Authors	:	Chaikuad, A.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Knapp, S.; Structural Genomics Consortium; Structural Genomics Consortium (SGC)
Deposited on	:	2018-12-19
Resolution	:	2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	2%, 87%, 9%
1	B	280	4%, 89%, 7%
1	C	280	7%, 91%, 6%
1	D	280	5%, 83%, 8%, 9%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase ULK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	273	2192	1402	385	391	14	0	1	0
1	B	269	2176	1397	375	389	15	0	4	0
1	C	275	2217	1417	390	396	14	0	2	0
1	D	254	2069	1328	362	367	12	0	5	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q8IYT8
A	-2	GLY	-	expression tag	UNP Q8IYT8
A	-1	GLY	-	expression tag	UNP Q8IYT8
A	0	SER	-	expression tag	UNP Q8IYT8
A	173	ASP	THR	conflict	UNP Q8IYT8
B	-3	GLY	-	expression tag	UNP Q8IYT8
B	-2	GLY	-	expression tag	UNP Q8IYT8
B	-1	GLY	-	expression tag	UNP Q8IYT8
B	0	SER	-	expression tag	UNP Q8IYT8
B	173	ASP	THR	conflict	UNP Q8IYT8
C	-3	GLY	-	expression tag	UNP Q8IYT8
C	-2	GLY	-	expression tag	UNP Q8IYT8
C	-1	GLY	-	expression tag	UNP Q8IYT8
C	0	SER	-	expression tag	UNP Q8IYT8
C	173	ASP	THR	conflict	UNP Q8IYT8
D	-3	GLY	-	expression tag	UNP Q8IYT8
D	-2	GLY	-	expression tag	UNP Q8IYT8
D	-1	GLY	-	expression tag	UNP Q8IYT8
D	0	SER	-	expression tag	UNP Q8IYT8
D	173	ASP	THR	conflict	UNP Q8IYT8

- Molecule 2 is {N}-[3-[[5-cyclopropyl-2-[(2-methyl-3,4-dihydro-1 {H}-isoquinolin-6-yl)amino]pyrimidin-4-yl]amino]propyl]cyclobutanecarboxamide (three-letter code: HVH) (formula: $C_{25}H_{34}N_6O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	32	25	6	1	0	0
			Total	C	N	O		
2	B	1	32	25	6	1	0	0
			Total	C	N	O		
2	C	1	32	25	6	1	0	0
			Total	C	N	O		
2	D	1	32	25	6	1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0
3	A	1	4	2	2	0	0
3	A	1	4	2	2	0	0
3	A	1	4	2	2	0	0
3	B	1	4	2	2	0	0
3	B	1	4	2	2	0	0
3	B	1	4	2	2	0	0
3	B	1	4	2	2	0	0
3	C	1	4	2	2	0	0
3	C	1	4	2	2	0	0
3	C	1	4	2	2	0	0
3	D	1	4	2	2	0	0
3	D	1	4	2	2	0	0
3	D	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	D	1	4	2	2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	B	1	1	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	6	3	3	0	0
5	C	1	6	3	3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	90	90	90	0	0
6	B	91	91	91	0	0
6	C	112	112	112	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	D	64	64	64	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serine/threonine-protein kinase ULK2

Chain A:



• Molecule 1: Serine/threonine-protein kinase ULK2

Chain B:



• Molecule 1: Serine/threonine-protein kinase ULK2

Chain C:



• Molecule 1: Serine/threonine-protein kinase ULK2

Chain D:


GLY
GLY
GLY
SER
M1
E8
R12
A19
R25
R29
L46
L53
L63
E66
N67
I68
Y96
K100
R120
A142
K146
R153
D158
A172
D173
L174
C175
G176
S177
M181
A182
P183
E184
Q189
H190
Y191
P213
F214
Q215
ALA

ASN
SER
PRO
GLN
ASP
LEU
ARG
MET
PHE
TYR
GLU
LYS
ASN
ARG
SER
LEU
MET
P234
R238
P242
Y243
R254
P269
E272
GLN
GLY
PRO
VAL

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	74.65Å 69.05Å 107.71Å 90.00° 97.93° 90.00°	Depositor
Resolution (Å)	57.97 – 2.05 57.97 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (57.97-2.05) 100.0 (57.97-2.05)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.34 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.195 , 0.231 0.205 , 0.235	Depositor DCC
R_{free} test set	3331 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å^2)	32.9	Xtriage
Anisotropy	0.315	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 52.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9212	wwPDB-VP
Average B, all atoms (Å^2)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HVH, GOL, EDO, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.69	1/2242 (0.0%)	0.66	0/3021
1	B	0.64	0/2235	0.65	0/3012
1	C	0.69	1/2272 (0.0%)	0.67	0/3062
1	D	0.67	1/2116 (0.0%)	0.66	0/2849
All	All	0.67	3/8865 (0.0%)	0.66	0/11944

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	3
1	C	0	4
1	D	0	6
All	All	0	19

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	8	GLU	CD-OE1	-6.33	1.18	1.25
1	D	8	GLU	CD-OE1	-5.31	1.19	1.25
1	C	62	GLU	CD-OE1	5.11	1.31	1.25

There are no bond angle outliers.

There are no chirality outliers.

5 of 19 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	120	ARG	Sidechain
1	A	130	ARG	Sidechain
1	A	238	ARG	Sidechain
1	A	25	ARG	Sidechain
1	A	29	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2192	0	2214	18	0
1	B	2176	0	2198	16	1
1	C	2217	0	2239	12	0
1	D	2069	0	2098	14	1
2	A	32	0	0	0	0
2	B	32	0	0	0	0
2	C	32	0	0	0	0
2	D	32	0	0	1	0
3	A	16	0	24	0	0
3	B	16	0	24	1	0
3	C	12	0	18	2	0
3	D	16	0	24	0	0
4	B	1	0	0	0	0
5	B	6	0	8	0	0
5	C	6	0	8	0	0
6	A	90	0	0	1	0
6	B	91	0	0	3	0
6	C	112	0	0	2	0
6	D	64	0	0	2	0
All	All	9212	0	8855	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:ASP:OD2	6:A:401:HOH:O	2.09	0.69
1:C:158:ASP:OD2	6:C:401:HOH:O	2.13	0.65
1:D:63:LEU:HD13	1:D:68:ILE:HD13	1.79	0.63
1:D:46:LEU:HD21	1:D:53[B]:LEU:HA	1.80	0.63
1:A:174:LEU:HD12	1:B:51:ILE:CD1	2.29	0.62

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33[B]:ASP:OD2	1:D:29:ARG:NH2[1_554]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/280 (97%)	266 (98%)	6 (2%)	0	100	100
1	B	269/280 (96%)	261 (97%)	8 (3%)	0	100	100
1	C	275/280 (98%)	267 (97%)	8 (3%)	0	100	100
1	D	255/280 (91%)	246 (96%)	9 (4%)	0	100	100
All	All	1071/1120 (96%)	1040 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	237/241 (98%)	235 (99%)	2 (1%)	81 82
1	B	237/241 (98%)	237 (100%)	0	100 100
1	C	241/241 (100%)	240 (100%)	1 (0%)	91 91
1	D	223/241 (92%)	223 (100%)	0	100 100
All	All	938/964 (97%)	935 (100%)	3 (0%)	92 93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	ARG
1	A	254	ARG
1	C	12	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 1 is monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	EDO	D	305	-	3,3,3	0.50	0	2,2,2	0.32	0
3	EDO	B	303	-	3,3,3	0.45	0	2,2,2	0.29	0
3	EDO	D	304	-	3,3,3	0.38	0	2,2,2	0.41	0
3	EDO	B	306	-	3,3,3	0.43	0	2,2,2	0.18	0
2	HVH	A	301	-	36,36,36	0.59	0	44,50,50	0.44	0
3	EDO	D	303	-	3,3,3	0.44	0	2,2,2	0.44	0
5	GOL	B	307	-	5,5,5	0.33	0	5,5,5	0.25	0
5	GOL	C	305	-	5,5,5	0.56	0	5,5,5	0.79	0
2	HVH	C	301	-	36,36,36	0.39	0	44,50,50	0.39	0
3	EDO	A	305	-	3,3,3	0.45	0	2,2,2	0.21	0
2	HVH	D	301	-	36,36,36	0.45	0	44,50,50	0.29	0
3	EDO	B	305	-	3,3,3	0.38	0	2,2,2	0.61	0
3	EDO	C	304	-	3,3,3	0.52	0	2,2,2	0.46	0
3	EDO	A	302	-	3,3,3	1.00	0	2,2,2	0.42	0
3	EDO	C	303	-	3,3,3	0.50	0	2,2,2	0.39	0
3	EDO	B	304	-	3,3,3	0.29	0	2,2,2	0.57	0
3	EDO	A	303	-	3,3,3	0.46	0	2,2,2	0.36	0
3	EDO	A	304	-	3,3,3	0.49	0	2,2,2	0.27	0
3	EDO	C	302	-	3,3,3	0.88	0	2,2,2	0.19	0
2	HVH	B	301	-	36,36,36	0.37	0	44,50,50	0.44	0
3	EDO	D	302	-	3,3,3	0.33	0	2,2,2	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	305	-	-	1/1/1/1	-
3	EDO	B	303	-	-	1/1/1/1	-
3	EDO	D	304	-	-	1/1/1/1	-
3	EDO	B	306	-	-	1/1/1/1	-
2	HVH	A	301	-	-	2/20/37/37	0/5/5/5
3	EDO	D	303	-	-	0/1/1/1	-
5	GOL	B	307	-	-	4/4/4/4	-
5	GOL	C	305	-	-	1/4/4/4	-
2	HVH	C	301	-	-	2/20/37/37	0/5/5/5
3	EDO	A	305	-	-	1/1/1/1	-
2	HVH	D	301	-	-	0/20/37/37	0/5/5/5
3	EDO	B	305	-	-	1/1/1/1	-
3	EDO	C	304	-	-	1/1/1/1	-
3	EDO	A	302	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	C	303	-	-	0/1/1/1	-
3	EDO	B	304	-	-	0/1/1/1	-
3	EDO	A	303	-	-	1/1/1/1	-
3	EDO	A	304	-	-	0/1/1/1	-
3	EDO	C	302	-	-	0/1/1/1	-
2	HVH	B	301	-	-	0/20/37/37	0/5/5/5
3	EDO	D	302	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	307	GOL	O1-C1-C2-C3
5	B	307	GOL	C1-C2-C3-O3
5	B	307	GOL	O1-C1-C2-O2
5	B	307	GOL	O2-C2-C3-O3
3	A	303	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	HVH	1	0
3	B	305	EDO	1	0
3	C	303	EDO	1	0
3	C	302	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	273/280 (97%)	-0.03	7 (2%)	56	60	27, 42, 86, 100	0
1	B	269/280 (96%)	-0.05	10 (3%)	41	45	24, 38, 81, 97	0
1	C	275/280 (98%)	0.05	19 (6%)	16	18	24, 37, 89, 113	0
1	D	254/280 (90%)	0.12	15 (5%)	22	24	25, 43, 85, 105	0
All	All	1071/1120 (95%)	0.02	51 (4%)	30	33	24, 40, 85, 113	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	176	GLY	5.7
1	C	176	GLY	5.5
1	D	215	GLN	5.0
1	D	272	GLU	4.8
1	C	174	LEU	4.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	EDO	A	302	4/4	0.62	0.26	41,48,49,51	0
3	EDO	C	302	4/4	0.65	0.21	48,50,52,55	0
3	EDO	B	303	4/4	0.70	0.17	63,65,66,67	0
5	GOL	C	305	6/6	0.77	0.14	50,55,57,58	0
3	EDO	C	304	4/4	0.80	0.20	59,62,64,67	0
3	EDO	A	305	4/4	0.82	0.16	61,63,63,65	0
3	EDO	D	304	4/4	0.83	0.22	59,64,69,69	0
3	EDO	D	305	4/4	0.83	0.12	59,60,60,62	0
3	EDO	A	303	4/4	0.83	0.16	57,59,61,68	0
3	EDO	D	303	4/4	0.88	0.14	59,60,62,64	0
3	EDO	C	303	4/4	0.89	0.12	45,47,49,56	0
3	EDO	D	302	4/4	0.89	0.12	55,57,60,64	0
3	EDO	B	304	4/4	0.90	0.18	64,64,65,75	0
3	EDO	B	305	4/4	0.90	0.14	41,44,44,47	0
3	EDO	A	304	4/4	0.94	0.11	57,58,60,61	0
5	GOL	B	307	6/6	0.95	0.19	45,47,52,57	0
2	HVH	A	301	32/32	0.96	0.11	21,26,33,39	0
2	HVH	B	301	32/32	0.96	0.10	20,26,35,37	0
2	HVH	D	301	32/32	0.97	0.10	23,25,38,42	0
2	HVH	C	301	32/32	0.97	0.10	20,25,30,31	0
3	EDO	B	306	4/4	0.97	0.13	30,30,34,38	0
4	NA	B	302	1/1	0.99	0.05	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HVH A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HVH C 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.