



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 6, 2026 – 03:12 pm BST

PDB ID : 9RTF / pdb_00009rtf
Title : Crystal structure of the human serum transferrin with Fe(III) bound at the C-lobe only
Authors : Paolillo, M.; Ferraro, G.; Banneville, A.S.; Cornaciu, I.; Pica, A.; Merlino, A.
Deposited on : 2025-07-02
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	679	10% 82% 15% ..

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	670	Total	C	N	O	S	0	5	0
			5243	3289	911	996	47			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total Fe 1 1	0	0

- Molecule 3 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is BICARBONATE ION (CCD ID: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	C	O	0	0
			4	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	354	Total	O	0	1
			355	355		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serotransferrin



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	138.15Å 158.03Å 107.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	104.01 – 2.16 104.01 – 2.16	Depositor EDS
% Data completeness (in resolution range)	74.6 (104.01-2.16) 74.9 (104.01-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.185 , 0.232 0.191 , 0.234	Depositor DCC
R_{free} test set	2373 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5636	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FE, BCT, NAG, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.04	1/5360 (0.0%)	1.38	9/7241 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	589	THR	C-O	6.35	1.30	1.23

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	79	PRO	CB-CA-C	-6.53	103.56	111.46
1	AAA	572	GLU	CB-CG-CD	5.52	121.98	112.60
1	AAA	551	ALA	CA-C-N	5.40	127.83	120.54
1	AAA	551	ALA	C-N-CA	5.40	127.83	120.54
1	AAA	652	GLY	CA-C-O	-5.21	118.40	122.52
1	AAA	206	LYS	CB-CA-C	-5.15	101.19	110.36
1	AAA	662	LEU	CA-C-N	5.09	127.36	120.38
1	AAA	662	LEU	C-N-CA	5.09	127.36	120.38
1	AAA	406	PRO	CA-C-O	-5.03	115.45	121.48

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	157	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	5243	0	5049	57	0
2	AAA	1	0	0	0	0
3	AAA	7	0	2	1	0
4	AAA	14	0	13	0	0
5	AAA	12	0	16	1	0
6	AAA	4	0	0	0	0
7	AAA	355	0	0	6	0
All	All	5636	0	5080	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:188:TYR:CE1	1:AAA:206:LYS:HG3	1.99	0.98
1:AAA:425:GLY:HA2	1:AAA:584:ASN:OD1	1.69	0.92
1:AAA:182:LEU:HD12	7:AAA:865:HOH:O	1.91	0.71
1:AAA:468:TYR:OH	1:AAA:664:LYS:HE3	1.97	0.65
1:AAA:3:ASP:CG	1:AAA:5:THR:HG22	2.23	0.63
1:AAA:69:ASP:OD1	1:AAA:327:ARG:NH2	2.31	0.63
1:AAA:663:ARG:NH2	1:AAA:675[A]:THR:HG21	2.14	0.63
1:AAA:76:ASN:HB3	7:AAA:1074:HOH:O	2.00	0.61
1:AAA:188:TYR:CZ	1:AAA:206:LYS:HG3	2.35	0.60
1:AAA:109:MET:HE1	1:AAA:243:LEU:HD21	1.83	0.60
1:AAA:23:ARG:HG3	1:AAA:37:VAL:O	2.02	0.59
1:AAA:646:THR:HG21	7:AAA:1079:HOH:O	2.02	0.59
1:AAA:272:GLU:O	1:AAA:278:LYS:HE2	2.05	0.57
1:AAA:26:MET:CE	1:AAA:274:PHE:CD2	2.89	0.56
1:AAA:63[B]:ASP:OD1	1:AAA:63[B]:ASP:C	2.50	0.54
1:AAA:646:THR:HG22	1:AAA:649:LYS:H	1.74	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:26:MET:HE2	1:AAA:274:PHE:CE2	2.43	0.53
1:AAA:163:ASP:OD2	1:AAA:166:ASP:OD2	2.27	0.52
1:AAA:418:CYS:O	1:AAA:419:GLU:C	2.54	0.51
1:AAA:457:THR:HG21	1:AAA:582:ALA:HB1	1.92	0.51
1:AAA:26:MET:HE1	1:AAA:274:PHE:HD2	1.77	0.50
1:AAA:26:MET:CE	1:AAA:274:PHE:HD2	2.24	0.50
1:AAA:538:VAL:HB	1:AAA:539:PRO:HD3	1.94	0.49
1:AAA:116:LYS:HA	1:AAA:156:GLY:O	2.12	0.49
1:AAA:26:MET:HE1	1:AAA:274:PHE:CD2	2.49	0.48
1:AAA:623:ARG:NH1	7:AAA:814:HOH:O	2.45	0.48
1:AAA:646:THR:HB	1:AAA:649:LYS:HE3	1.97	0.47
1:AAA:354:LYS:HD3	1:AAA:630:LEU:HD22	1.96	0.47
1:AAA:26:MET:HE2	1:AAA:274:PHE:HE2	1.80	0.47
1:AAA:90:ASP:OD1	1:AAA:92[A]:GLN:NE2	2.49	0.44
1:AAA:281:GLU:HG3	1:AAA:282:PHE:N	2.33	0.44
1:AAA:355:CYS:C	1:AAA:368:CYS:SG	3.00	0.44
1:AAA:377:CYS:HB3	1:AAA:389:MET:SD	2.58	0.44
1:AAA:397:TYR:CE2	1:AAA:659:VAL:HG11	2.53	0.43
1:AAA:254:ARG:NH1	1:AAA:257:GLY:O	2.51	0.43
1:AAA:243:LEU:O	1:AAA:318:GLU:HB3	2.19	0.43
1:AAA:460:TRP:O	1:AAA:464:MET:HB2	2.19	0.43
1:AAA:109:MET:HE2	1:AAA:109:MET:HB2	1.92	0.42
1:AAA:135:LEU:O	1:AAA:136:TYR:C	2.62	0.42
1:AAA:407:VAL:HG12	1:AAA:594:GLU:HG3	2.02	0.42
1:AAA:26:MET:CE	1:AAA:274:PHE:CE2	3.02	0.42
1:AAA:471:ILE:HD12	1:AAA:473:HIS:CE1	2.55	0.41
1:AAA:3:ASP:CG	1:AAA:5:THR:CG2	2.92	0.41
1:AAA:71:TYR:HB2	1:AAA:311:ALA:CB	2.50	0.41
1:AAA:411:ASN:HD21	1:AAA:421:THR:CB	2.33	0.41
1:AAA:421:THR:HA	1:AAA:422:PRO:HD3	1.85	0.41
1:AAA:423:GLU:CD	1:AAA:581:ARG:HG2	2.45	0.41
1:AAA:448:LYS:HA	1:AAA:482:GLU:O	2.21	0.41
1:AAA:538:VAL:HG11	1:AAA:571:VAL:HG21	2.01	0.41
1:AAA:183:ASN:HA	5:AAA:705:GOL:O3	2.20	0.41
1:AAA:36:SER:HA	7:AAA:853:HOH:O	2.21	0.41
1:AAA:375:GLU:HG3	7:AAA:1093:HOH:O	2.20	0.41
1:AAA:624:SER:HB2	1:AAA:629:LEU:HD22	2.02	0.41
1:AAA:356:ASP:O	1:AAA:359:SER:HB3	2.20	0.41
1:AAA:585:HIS:CE1	3:AAA:702:MLI:O9	2.74	0.41
1:AAA:538:VAL:O	1:AAA:539:PRO:C	2.64	0.40
1:AAA:421:THR:HG23	1:AAA:424:ALA:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	669/679 (98%)	626 (94%)	35 (5%)	8 (1%)	10 5

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	419	GLU
1	AAA	422	PRO
1	AAA	626	THR
1	AAA	86	GLY
1	AAA	423	GLU
1	AAA	436	ALA
1	AAA	614	ASP
1	AAA	332	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	569/572 (100%)	539 (95%)	30 (5%)	20 17

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	4	LYS
1	AAA	5	THR
1	AAA	29	VAL
1	AAA	71	TYR
1	AAA	89	GLU
1	AAA	90	ASP
1	AAA	122	LEU
1	AAA	249	HIS
1	AAA	300	HIS
1	AAA	327	ARG
1	AAA	338	GLU
1	AAA	365	LYS
1	AAA	369	VAL
1	AAA	370	SER
1	AAA	421	THR
1	AAA	437	SER
1	AAA	447	LYS
1	AAA	496	LYS
1	AAA	501	SER
1	AAA	555	ASN
1	AAA	611	ASN
1	AAA	618	ASN
1	AAA	625	GLU
1	AAA	627	LYS
1	AAA	629	LEU
1	AAA	630	LEU
1	AAA	637	CYS
1	AAA	646	THR
1	AAA	657	LYS
1	AAA	663	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MLI	AAA	702	2	6,6,6	0.93	0	7,7,7	1.47	0
5	GOL	AAA	705	-	5,5,5	0.36	0	5,5,5	0.38	0
4	NAG	AAA	703	1	14,14,15	0.67	0	17,19,21	1.38	3 (17%)
5	GOL	AAA	704	-	5,5,5	0.41	0	5,5,5	0.69	0
6	BCT	AAA	706	-	2,3,3	0.71	0	2,3,3	1.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	AAA	705	-	-	2/4/4/4	-
4	NAG	AAA	703	1	-	0/6/23/26	0/1/1/1
3	MLI	AAA	702	2	-	1/4/4/4	-
5	GOL	AAA	704	-	-	4/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	AAA	703	NAG	O5-C5-C6	3.00	111.91	107.20
4	AAA	703	NAG	O3-C3-C2	2.96	115.59	109.47
4	AAA	703	NAG	C2-N2-C7	2.21	126.05	122.90

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	AAA	704	GOL	O1-C1-C2-O2
5	AAA	704	GOL	O1-C1-C2-C3
5	AAA	704	GOL	C1-C2-C3-O3
5	AAA	705	GOL	C1-C2-C3-O3
5	AAA	705	GOL	O2-C2-C3-O3
5	AAA	704	GOL	O2-C2-C3-O3
3	AAA	702	MLI	C2-C1-C3-O9

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	702	MLI	1	0
5	AAA	705	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	670/679 (98%)	0.46	67 (10%) 12 14	11, 34, 77, 126	5 (0%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	424	ALA	7.4
1	AAA	422	PRO	7.4
1	AAA	606[A]	HIS	7.4
1	AAA	421	THR	6.3
1	AAA	412	TYR	5.7
1	AAA	88	LYS	5.7
1	AAA	613	THR	5.2
1	AAA	626	THR	4.9
1	AAA	86	GLY	4.8
1	AAA	423	GLU	4.7
1	AAA	419	GLU	4.6
1	AAA	612	VAL	4.5
1	AAA	418	CYS	4.1
1	AAA	91	PRO	4.1
1	AAA	87	SER	4.0
1	AAA	339	CYS	3.8
1	AAA	623	ARG	3.5
1	AAA	601	LEU	3.5
1	AAA	340	LYS	3.5
1	AAA	615	CYS	3.4
1	AAA	349	HIS	3.3
1	AAA	332	PRO	3.2
1	AAA	338	GLU	3.2
1	AAA	89	GLU	3.1
1	AAA	417	ASN	3.1
1	AAA	611	ASN	3.0
1	AAA	32	SER	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	347	LEU	3.0
1	AAA	619	PHE	3.0
1	AAA	360	VAL	2.9
1	AAA	333	GLU	2.9
1	AAA	438	ASP	2.8
1	AAA	667	THR	2.8
1	AAA	420	ASP	2.8
1	AAA	627	LYS	2.8
1	AAA	33	ASP	2.7
1	AAA	256	MET	2.7
1	AAA	625	GLU	2.6
1	AAA	620	CYS	2.6
1	AAA	350	HIS	2.6
1	AAA	607	LEU	2.6
1	AAA	603	GLN	2.6
1	AAA	643	ASP	2.6
1	AAA	436	ALA	2.5
1	AAA	572	GLU	2.5
1	AAA	610	SER	2.5
1	AAA	3	ASP	2.4
1	AAA	90	ASP	2.4
1	AAA	637	CYS	2.4
1	AAA	639	ALA	2.4
1	AAA	363	VAL	2.4
1	AAA	365	LYS	2.4
1	AAA	665	CYS	2.4
1	AAA	622	PHE	2.3
1	AAA	617	GLY	2.3
1	AAA	372	GLU	2.3
1	AAA	281	GLU	2.2
1	AAA	602	ARG	2.2
1	AAA	471	ILE	2.1
1	AAA	503	LEU	2.1
1	AAA	437	SER	2.1
1	AAA	618	ASN	2.1
1	AAA	663	ARG	2.1
1	AAA	614	ASP	2.0
1	AAA	427	PHE	2.0
1	AAA	277	ASP	2.0
1	AAA	425	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	AAA	703	14/15	0.64	0.21	95,102,111,121	0
5	GOL	AAA	704	6/6	0.85	0.23	33,47,54,64	0
5	GOL	AAA	705	6/6	0.90	0.20	34,46,53,63	0
6	BCT	AAA	706	4/4	0.91	0.12	43,47,52,61	0
3	MLI	AAA	702	7/7	0.95	0.08	25,27,28,31	0
2	FE	AAA	701	1/1	1.00	0.02	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around MLI AAA 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE AAA 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.