



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 07:56 AM EDT

PDB ID : 1RXQ
Title : YfiT from *Bacillus subtilis* is a probable metal-dependent hydrolase with an unusual four-helix bundle topology
Authors : Rajan, S.S.; Yang, X.; Anderson, W.F.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2003-12-18
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	<div> <div>15%</div> <div>78%</div> <div>19%</div> <div>..</div> </div>
1	B	178	<div> <div>8%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>
1	C	178	<div> <div>10%</div> <div>83%</div> <div>13%</div> <div>..</div> </div>
1	D	178	<div> <div>9%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called yfiT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	Se	0	5	0
			1449	923	253	269	4			
1	B	175	Total	C	N	O	Se	0	7	0
			1465	937	256	268	4			
1	C	174	Total	C	N	O	Se	0	4	0
			1446	920	254	268	4			
1	D	169	Total	C	N	O	Se	0	6	0
			1418	907	248	259	4			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	GB 16077906
A	45	MSE	MET	MODIFIED RESIDUE	GB 16077906
A	73	MSE	MET	MODIFIED RESIDUE	GB 16077906
A	175	MSE	MET	MODIFIED RESIDUE	GB 16077906
B	1	MSE	MET	MODIFIED RESIDUE	GB 16077906
B	45	MSE	MET	MODIFIED RESIDUE	GB 16077906
B	73	MSE	MET	MODIFIED RESIDUE	GB 16077906
B	175	MSE	MET	MODIFIED RESIDUE	GB 16077906
C	1	MSE	MET	MODIFIED RESIDUE	GB 16077906
C	45	MSE	MET	MODIFIED RESIDUE	GB 16077906
C	73	MSE	MET	MODIFIED RESIDUE	GB 16077906
C	175	MSE	MET	MODIFIED RESIDUE	GB 16077906
D	1	MSE	MET	MODIFIED RESIDUE	GB 16077906
D	45	MSE	MET	MODIFIED RESIDUE	GB 16077906
D	73	MSE	MET	MODIFIED RESIDUE	GB 16077906
D	175	MSE	MET	MODIFIED RESIDUE	GB 16077906

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0
2	B	1	Total Ni 1 1	0	0
2	C	1	Total Ni 1 1	0	0
2	D	1	Total Ni 1 1	0	0

- Molecule 3 is GLUTAMIC ACID (three-letter code: GLU) (formula: $C_5H_9NO_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total C N O 9 5 1 3	0	0

- Molecule 4 is SERINE (three-letter code: SER) (formula: $C_3H_7NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			6	3	1	2		
4	D	1	Total	C	N	O	0	0
			6	3	1	2		

- Molecule 5 is ALANINE (three-letter code: ALA) (formula: $C_3H_7NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 6 is THREONINE (three-letter code: THR) (formula: $C_4H_9NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	N	O	0	0
			7	4	1	2		

- Molecule 7 is GLYCINE (three-letter code: GLY) (formula: C₂H₅NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	N	O	0	0
			4	2	1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	114	Total 116	O 116	0	2
8	B	109	Total 111	O 111	0	1
8	C	124	Total 128	O 128	0	6
8	D	122	Total 123	O 123	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: yfiT




- Molecule 1: yfiT




- Molecule 1: yfiT



- Molecule 1: yfiT





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.37Å 50.42Å 89.41Å 104.38° 90.61° 112.05°	Depositor
Resolution (Å)	30.00 – 1.70 27.56 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.6 (30.00-1.70) 93.6 (27.56-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.186 , 0.217 0.198 , 0.225	Depositor DCC
R_{free} test set	7451 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6297	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1505	0.70	4/2036 (0.2%)
1	B	0.46	0/1529	0.68	4/2068 (0.2%)
1	C	0.51	0/1498	0.75	4/2026 (0.2%)
1	D	0.51	0/1478	0.76	3/1999 (0.2%)
All	All	0.48	0/6010	0.72	15/8129 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	25	ASP	CB-CG-OD2	6.66	124.30	118.30
1	B	25	ASP	CB-CG-OD2	6.01	123.71	118.30
1	C	108	ASP	CB-CG-OD2	5.83	123.55	118.30
1	A	47	ASP	CB-CG-OD2	5.82	123.53	118.30
1	C	56	ASP	CB-CG-OD2	5.74	123.47	118.30
1	A	103	ASP	CB-CG-OD2	5.68	123.41	118.30
1	A	143	ASP	CB-CG-OD2	5.66	123.39	118.30
1	B	103	ASP	CB-CG-OD2	5.51	123.26	118.30
1	D	56	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	56	ASP	CB-CG-OD2	5.37	123.13	118.30
1	C	143	ASP	CB-CG-OD2	5.37	123.13	118.30
1	B	132	ASP	CB-CG-OD2	5.27	123.04	118.30
1	C	25	ASP	CB-CG-OD2	5.18	122.97	118.30
1	D	143	ASP	CB-CG-OD2	5.18	122.96	118.30
1	B	56	ASP	CB-CG-OD2	5.14	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1449	0	1422	24	0
1	B	1465	0	1454	16	0
1	C	1446	0	1419	15	1
1	D	1418	0	1397	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	D	9	0	5	2	0
4	D	12	0	10	1	0
5	D	5	0	5	2	0
6	D	7	0	7	0	0
7	D	4	0	3	0	1
8	A	116	0	0	0	0
8	B	111	0	0	0	0
8	C	128	0	0	2	0
8	D	123	0	0	3	0
All	All	6297	0	5722	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73[A]:MSE:HE3	1:D:73[A]:MSE:CG	1.83	1.09
1:A:73[A]:MSE:HE3	1:D:73[A]:MSE:HG3	1.11	1.08
3:D:501:GLU:HB3	8:D:625:HOH:O	1.57	0.96
1:A:73[A]:MSE:CE	1:D:73[A]:MSE:HG3	1.99	0.91
1:A:72:HIS:HE1	1:A:160:HIS:HD2	1.20	0.90
1:D:156[A]:LEU:HD21	5:D:503:ALA:HA	1.55	0.89
1:C:46:THR:H	1:C:49:GLN:HE21	1.18	0.86
1:B:131:THR:H	1:B:134:GLN:HE21	1.30	0.80
1:C:131:THR:H	1:C:134:GLN:HE21	1.32	0.76
3:D:501:GLU:CB	8:D:625:HOH:O	2.21	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:60:THR:H	1:D:63:GLN:HE21	1.36	0.73
1:B:119:LEU:HD13	1:C:73[B]:MSE:HE1	1.69	0.73
1:D:10:ILE:HG23	1:D:10:ILE:O	1.88	0.73
1:A:72:HIS:CE1	1:A:160:HIS:HD2	2.08	0.70
1:A:72:HIS:HD2	1:A:120:HIS:NE2	1.95	0.65
1:C:46:THR:H	1:C:49:GLN:NE2	1.95	0.63
1:A:66:HIS:HD2	1:A:112:SER:OG	1.83	0.62
1:A:42[B]:VAL:HG21	1:A:113:LEU:HD11	1.80	0.62
1:C:39:LYS:HE3	8:C:374:HOH:O	2.01	0.61
1:A:77:ILE:HD12	1:D:77:ILE:CD1	2.30	0.61
1:C:164:HIS:HD1	1:C:167:HIS:HD2	1.49	0.60
1:A:131:THR:H	1:A:134:GLN:HE21	1.47	0.60
1:D:156[A]:LEU:HD11	4:D:502:SER:O	2.02	0.60
1:D:13:TYR:CD2	1:D:156[B]:LEU:HG	2.38	0.58
1:A:32:GLU:OE1	1:A:128:ARG:NH1	2.36	0.58
1:A:77:ILE:CD1	1:D:77:ILE:HD12	2.35	0.57
1:D:102:LYS:NZ	8:D:602:HOH:O	2.37	0.57
1:B:73[B]:MSE:HE2	1:B:115[B]:LEU:HD21	1.87	0.55
1:B:116[B]:LEU:HD22	1:B:120:HIS:CD2	2.41	0.55
1:A:77:ILE:HD12	1:D:77:ILE:HD12	1.90	0.53
1:A:72:HIS:HE1	1:A:160:HIS:CD2	2.12	0.53
1:B:73[B]:MSE:CE	1:B:115[B]:LEU:HD21	2.40	0.52
1:A:42[B]:VAL:CG2	1:A:113:LEU:HD11	2.39	0.52
1:B:90:ILE:HD12	1:B:90:ILE:C	2.30	0.52
1:C:5:ASN:ND2	1:C:8:TYR:H	2.08	0.52
1:D:141:HIS:HB3	1:D:144:THR:HG22	1.91	0.52
1:D:156[A]:LEU:HD21	5:D:503:ALA:CA	2.36	0.52
1:B:29:GLN:O	1:B:33:GLU:HG2	2.10	0.51
1:D:54:TYR:CE1	1:D:167:HIS:ND1	2.78	0.51
1:B:100:GLU:OE2	1:B:105:LYS:CE	2.58	0.51
1:D:90:ILE:HD11	1:D:141:HIS:ND1	2.26	0.51
1:D:90:ILE:C	1:D:90:ILE:HD12	2.32	0.50
1:D:10:ILE:O	1:D:10:ILE:CG2	2.59	0.50
1:D:54:TYR:CE1	1:D:167:HIS:CE1	2.99	0.50
1:B:42[A]:VAL:HG21	1:B:113:LEU:HD11	1.94	0.50
1:B:73[B]:MSE:HG2	1:C:73[B]:MSE:HE2	1.93	0.49
1:B:131:THR:H	1:B:134:GLN:NE2	2.06	0.49
1:B:100:GLU:HA	1:B:105:LYS:HE3	1.95	0.49
1:A:69:ALA:O	1:A:73[B]:MSE:HG3	2.12	0.49
1:A:38:LEU:O	1:A:42[B]:VAL:HG13	2.11	0.49
1:A:163:HIS:CE1	1:A:167:HIS:CE1	3.03	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:54:TYR:HA	1:D:171:LEU:HD22	1.96	0.46
1:C:89:ALA:HB2	1:C:140:TYR:CZ	2.50	0.46
1:C:133:GLN:NE2	1:C:133:GLN:H	2.14	0.45
1:A:119:LEU:HD13	1:D:73[A]:MSE:HE1	1.99	0.45
1:C:45:MSE:HA	1:C:49:GLN:NE2	2.32	0.45
1:A:160:HIS:CE1	1:A:164:HIS:CD2	3.06	0.43
1:B:42[A]:VAL:CG2	1:B:113:LEU:HD11	2.48	0.43
1:B:49:GLN:HG2	1:B:177:TRP:CZ2	2.53	0.43
1:A:174:ARG:HG2	1:A:175:MSE:HE2	2.01	0.42
1:C:39:LYS:CE	8:C:374:HOH:O	2.61	0.42
1:D:60:THR:H	1:D:63:GLN:NE2	2.11	0.42
1:A:7:SER:O	1:A:163:HIS:HE1	2.02	0.42
1:A:90:ILE:HD12	1:A:90:ILE:C	2.41	0.41
1:A:54:TYR:CE2	1:A:55:ARG:HG2	2.55	0.41
1:B:115[B]:LEU:HD12	1:C:115:LEU:HD12	2.03	0.41
1:B:73[A]:MSE:HE1	1:C:76:TYR:CE2	2.57	0.40
1:C:90:ILE:C	1:C:90:ILE:HD12	2.41	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:5:ASN:N	7:D:506:GLY:C[1_456]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/178 (99%)	175 (99%)	2 (1%)	0	100	100
1	B	180/178 (101%)	178 (99%)	2 (1%)	0	100	100
1	C	176/178 (99%)	173 (98%)	2 (1%)	1 (1%)	25	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	173/178 (97%)	172 (99%)	1 (1%)	0	100	100
All	All	706/712 (99%)	698 (99%)	7 (1%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	145	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/156 (103%)	152 (94%)	9 (6%)	21	7
1	B	164/156 (105%)	159 (97%)	5 (3%)	41	22
1	C	160/156 (103%)	152 (95%)	8 (5%)	24	8
1	D	157/156 (101%)	153 (98%)	4 (2%)	47	29
All	All	642/624 (103%)	616 (96%)	26 (4%)	31	13

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ARG
1	A	21	LYS
1	A	22[A]	GLU
1	A	22[B]	GLU
1	A	55	ARG
1	A	72	HIS
1	A	145	LYS
1	A	170	GLU
1	A	178	SER
1	B	4	VAL
1	B	17	GLU
1	B	18	SER
1	B	26	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	40	GLN
1	C	5	ASN
1	C	16	ARG
1	C	32	GLU
1	C	133	GLN
1	C	141	HIS
1	C	145	LYS
1	C	146	GLU
1	C	173	ARG
1	D	10	ILE
1	D	21	LYS
1	D	40	GLN
1	D	145	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	HIS
1	A	72	HIS
1	A	134	GLN
1	A	160	HIS
1	B	40	GLN
1	B	134	GLN
1	C	5	ASN
1	C	49	GLN
1	C	133	GLN
1	C	134	GLN
1	C	167	HIS
1	D	49	GLN
1	D	63	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ALA	D	503	4,6	3,4,5	0.72	0	2,4,6	0.79	0
3	GLU	D	501	4,2	7,8,9	1.04	0	4,9,11	0.70	0
4	SER	D	502	5,3	4,5,6	0.72	0	1,5,7	1.42	0
7	GLY	D	506	4	3,3,4	0.60	0	1,2,4	0.05	0
4	SER	D	505	7,6	4,5,6	0.52	0	1,5,7	0.76	0
6	THR	D	504	5,4	5,6,7	0.52	0	5,7,9	1.26	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ALA	D	503	4,6	-	0/1/2/4	-
3	GLU	D	501	4,2	-	1/6/7/9	-
4	SER	D	502	5,3	-	2/2/4/6	-
7	GLY	D	506	4	-	0/0/1/2	-
4	SER	D	505	7,6	-	0/2/4/6	-
6	THR	D	504	5,4	-	2/5/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	504	THR	OG1-CB-CG2	-2.00	103.69	109.68

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	502	SER	N-CA-CB-OG
4	D	502	SER	C-CA-CB-OG
6	D	504	THR	C-CA-CB-CG2
3	D	501	GLU	CA-CB-CG-CD
6	D	504	THR	N-CA-CB-CG2

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	503	ALA	2	0
3	D	501	GLU	2	0
4	D	502	SER	1	0
7	D	506	GLY	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	171/178 (96%)	0.87	27 (15%) 2 2	14, 22, 47, 58	0
1	B	172/178 (96%)	0.52	14 (8%) 12 14	14, 21, 36, 55	0
1	C	171/178 (96%)	0.47	18 (10%) 6 7	12, 19, 37, 60	0
1	D	166/178 (93%)	0.48	16 (9%) 8 9	12, 19, 36, 52	0
All	All	680/712 (95%)	0.59	75 (11%) 5 6	12, 21, 40, 60	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	SER	8.3
1	A	6	LEU	7.8
1	B	178	SER	7.7
1	B	4	VAL	7.5
1	D	143	ASP	7.2
1	D	178	SER	6.8
1	D	10	ILE	6.5
1	A	5	ASN	6.2
1	C	178	SER	5.6
1	C	143	ASP	5.5
1	A	142	PRO	5.2
1	A	11	GLY	4.7
1	A	145	LYS	4.6
1	A	143	ASP	4.4
1	D	56	ASP	4.4
1	D	11	GLY	4.3
1	A	56	ASP	4.2
1	B	44	VAL	4.1
1	A	8	TYR	3.9
1	C	176	GLY	3.8
1	A	173	ARG	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	145	LYS	3.7
1	D	54	TYR	3.7
1	C	17	GLU	3.5
1	A	18	SER	3.5
1	B	56	ASP	3.4
1	A	9	PRO	3.4
1	A	144	THR	3.4
1	D	144	THR	3.3
1	C	142	PRO	3.2
1	B	5	ASN	3.2
1	B	17	GLU	3.1
1	C	140	TYR	3.1
1	B	6	LEU	3.0
1	B	176	GLY	3.0
1	D	116[A]	LEU	3.0
1	C	18	SER	2.9
1	A	147	ILE	2.9
1	A	132	ASP	2.9
1	A	146	GLU	2.9
1	C	44	VAL	2.8
1	A	10	ILE	2.7
1	B	8	TYR	2.7
1	A	57	GLY	2.7
1	C	141	HIS	2.7
1	A	115[A]	LEU	2.7
1	A	7	SER	2.7
1	A	17	GLU	2.6
1	D	12	GLU	2.6
1	D	57	GLY	2.6
1	A	68	LEU	2.5
1	A	69	ALA	2.5
1	C	116[A]	LEU	2.3
1	A	76	TYR	2.3
1	A	12	GLU	2.3
1	C	56	ASP	2.3
1	C	146	GLU	2.3
1	C	76	TYR	2.3
1	D	140	TYR	2.3
1	B	123	TRP	2.3
1	B	76	TYR	2.3
1	A	177	TRP	2.2
1	D	69	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	159[A]	TRP	2.2
1	C	57	GLY	2.2
1	D	115[A]	LEU	2.2
1	A	123	TRP	2.2
1	D	68	LEU	2.2
1	B	68	LEU	2.1
1	D	142	PRO	2.1
1	B	132	ASP	2.1
1	C	119	LEU	2.1
1	B	173	ARG	2.1
1	C	177	TRP	2.0
1	C	173	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	THR	D	504	7/8	0.74	0.26	29,37,40,42	7
7	GLY	D	506	4/5	0.76	0.26	30,32,38,42	4
5	ALA	D	503	5/6	0.78	0.26	31,35,38,40	5
4	SER	D	502	6/7	0.81	0.27	34,35,39,42	6
4	SER	D	505	6/7	0.85	0.17	32,36,39,40	6
3	GLU	D	501	9/10	0.96	0.16	23,28,42,47	9
2	NI	D	300	1/1	0.97	0.11	26,26,26,26	0
2	NI	B	300	1/1	0.99	0.13	22,22,22,22	0
2	NI	C	300	1/1	0.99	0.16	19,19,19,19	0
2	NI	A	300	1/1	0.99	0.10	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.