wwPDB X-ray Structure Validation Summary Report (i)

Jan 13, 2024 – 12:32 pm GMT

PDB ID	:	6TK1
Title	:	Femtosecond to millisecond structural changes in a light-driven sodium pump: 20ms structure of KR2 with extrapolated, light and dark datasets
Authors	:	Skopintsev, P.; Ehrenberg, D.; Weinert, T.; James, D.; Kar, R.; Johnson, P.; Ozerov, D.; Furrer, A.; Martiel, I.; Dworkowski, F.; Nass, K.; Knopp, G.; Cirelli, C.; Gashi, D.; Mous, S.; Wranik, M.; Gruhl, T.; Kekilli, D.; Bruenle, S.; Deupi, X.; Schertler, G.F.X.; Benoit, R.; Panneels, V.; Nogly, P.; Schapiro, I.; Milne, C.; Heberle, J.; Standfuss, J.
Deposited on	:	2019-11-28
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	4%, 70%, 20%, • 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LFA	A	314	-	-	-	X
3	LFA	A	316	-	-	-	X
3	LFA	A	318	-	-	-	X
3	LFA	A	321	-	-	-	X
3	LFA	A	325	-	-	-	X
4	NA	A	328	-	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium pumping rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	2099	1405	318	367	9	3	3	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	276	GLU	-	expression tag	UNP N0DKS8
A	277	ASN	-	expression tag	UNP N0DKS8
A	278	LEU	-	expression tag	UNP N0DKS8
A	279	TYR	-	expression tag	UNP N0DKS8
A	280	PHE	-	expression tag	UNP N0DKS8
A	281	GLN	-	expression tag	UNP N0DKS8
A	282	SER	-	expression tag	UNP N0DKS8
A	283	GLY	-	expression tag	UNP N0DKS8
A	284	SER	-	expression tag	UNP N0DKS8
A	285	HIS	-	expression tag	UNP N0DKS8
A	286	HIS	-	expression tag	UNP N0DKS8
A	287	HIS	-	expression tag	UNP N0DKS8
A	288	HIS	-	expression tag	UNP N0DKS8
A	289	HIS	-	expression tag	UNP N0DKS8
A	290	HIS	-	expression tag	UNP N0DKS8

- Molecule 2 is RETINAL (three-letter code: RET) (formula: $C_{20}H_{28}O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 20 20	0	0

- Molecule 3 is EICOSANE (three-letter code: LFA) (formula: $C_{20}H_{42}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 16 16	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 6 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C		
3	A	1	6	6	0	0
			Total	C		
3	A	1	5	5	0	0
			Total	C		
3	A	1	13	13	0	0
			Total	C		
3	A	1	10	10	0	0
			Total	C		
3	A	1	8	8	0	0
			Total	C		
3	A	1	8	8	0	0
			Total	C		
3	A	1	12	12	0	0
			Total	C		
3	A	1	16	16	0	0
			Total	C		
3	A	1	5	5	0	0
			Total	C		
3	A	1	10	10	0	0
			Total	C		
3	A	1	5	5	0	0
			Total	C		
3	A	1	11	11	0	0
			Total	C		
3	A	1	9	9	0	0
			Total	C		
3	A	1	9	9	0	0
			Total	C		
3	A	1	6	6	0	0
			Total	C		
3	A	1	5	5	0	0
			Total	C		
3	A	1	4	4	0	0
			Total	C		
3	A	1	6	6	0	0
			Total	C		
3	A	1	10	10	0	0
			Total	C		
3	A	1	3	3	0	0
			Total	C		
3	A	1	9	9	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C		
3	A	1	10	10	0	0
			Total	C		
3	A	1	12	12	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	47	47	47	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sodium pumping rhodopsin

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α, β, γ	41.54Å 84.48Å 235.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.29 – 2.50 12.29 – 2.50	Depositor EDS
% Data completeness (in resolution range)	89.9 (12.29-2.50) 89.9 (12.29-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.85 (at 2.50Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.278 , 0.353 0.278 , 0.353	Depositor DCC
R_{free} test set	857 reflections (6.46%)	wwPDB-VP
Wilson B-factor (Å^2)	47.7	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 99.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	2389	wwPDB-VP
Average B, all atoms (Å^2)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: RET, NA, LFA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.42	0/2167	0.54	0/2951

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2099	0	2071	37	0
2	A	20	0	27	2	0
3	A	222	0	403	7	0
4	A	1	0	0	0	0
5	A	47	0	0	6	0
All	All	2389	0	2501	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:SER:OG	5:A:420:HOH:O	2.00	0.80
1:A:117:VAL:HG23	1:A:118:PRO:HD3	1.67	0.76
1:A:109:ARG:NH2	5:A:461:HOH:O	2.21	0.73
1:A:55:LYS:HA	1:A:58:GLN:HG3	1.72	0.70
1:A:191:GLU:OE2	5:A:414:HOH:O	2.11	0.67

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	266/290 (92%)	252 (95%)	12 (4%)	2 (1%)	19 35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	SER
1	A	233	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	221/250 (88%)	210 (95%)	11 (5%)	24 46

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	TRP
1	A	233	PHE
1	A	271	LYS
1	A	245	LEU
1	A	161	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN
1	A	157	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 1 is monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	LFA	A	322	-	5,5,19	0.31	0	4,4,18	0.57	0
3	LFA	A	312	-	15,15,19	0.28	0	14,14,18	0.90	0
3	LFA	A	325	-	8,8,19	0.30	0	7,7,18	0.77	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	LFA	A	315	-	4,4,19	0.32	0	3,3,18	0.54	0
3	LFA	A	319	-	5,5,19	0.32	0	4,4,18	0.56	0
3	LFA	A	316	-	10,10,19	0.30	0	9,9,18	0.82	0
3	LFA	A	309	-	7,7,19	0.31	0	6,6,18	0.73	0
3	LFA	A	308	-	9,9,19	0.30	0	8,8,18	0.77	0
3	LFA	A	303	-	7,7,19	0.31	0	6,6,18	0.72	0
3	LFA	A	305	-	5,5,19	0.32	0	4,4,18	0.56	0
2	RET	A	301	1	20,20,21	0.69	0	27,27,28	0.69	1 (3%)
3	LFA	A	314	-	9,9,19	0.30	0	8,8,18	0.80	0
3	LFA	A	311	-	11,11,19	0.30	0	10,10,18	0.84	0
3	LFA	A	327	-	11,11,19	0.30	0	10,10,18	0.82	0
3	LFA	A	324	-	2,2,19	0.34	0	0,1,18	-	-
3	LFA	A	304	-	5,5,19	0.31	0	4,4,18	0.57	0
3	LFA	A	321	-	3,3,19	0.41	0	2,2,18	0.73	0
3	LFA	A	313	-	4,4,19	0.32	0	3,3,18	0.55	0
3	LFA	A	310	-	7,7,19	0.30	0	6,6,18	0.73	0
3	LFA	A	317	-	8,8,19	0.30	0	7,7,18	0.76	0
3	LFA	A	302	-	15,15,19	0.30	0	14,14,18	0.86	0
3	LFA	A	326	-	9,9,19	0.31	0	8,8,18	0.77	0
3	LFA	A	307	-	12,12,19	0.30	0	11,11,18	0.85	0
3	LFA	A	323	-	9,9,19	0.31	0	8,8,18	0.77	0
3	LFA	A	318	-	8,8,19	0.31	0	7,7,18	0.77	0
3	LFA	A	320	-	4,4,19	0.32	0	3,3,18	0.56	0
3	LFA	A	306	-	4,4,19	0.32	0	3,3,18	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LFA	A	322	-	-	0/3/3/17	-
3	LFA	A	312	-	-	4/13/13/17	-
3	LFA	A	325	-	-	0/6/6/17	-
3	LFA	A	315	-	-	0/2/2/17	-
3	LFA	A	319	-	-	2/3/3/17	-
3	LFA	A	316	-	-	0/8/8/17	-
3	LFA	A	309	-	-	0/5/5/17	-
3	LFA	A	308	-	-	3/7/7/17	-
3	LFA	A	303	-	-	2/5/5/17	-
3	LFA	A	305	-	-	1/3/3/17	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	301	1	-	2/13/30/31	0/1/1/1
3	LFA	A	314	-	-	1/7/7/17	-
3	LFA	A	311	-	-	5/9/9/17	-
3	LFA	A	327	-	-	2/9/9/17	-
3	LFA	A	304	-	-	0/3/3/17	-
3	LFA	A	321	-	-	1/1/1/17	-
3	LFA	A	313	-	-	0/2/2/17	-
3	LFA	A	310	-	-	1/5/5/17	-
3	LFA	A	317	-	-	4/6/6/17	-
3	LFA	A	302	-	-	4/13/13/17	-
3	LFA	A	326	-	-	3/7/7/17	-
3	LFA	A	307	-	-	3/10/10/17	-
3	LFA	A	323	-	-	1/7/7/17	-
3	LFA	A	318	-	-	2/6/6/17	-
3	LFA	A	320	-	-	0/2/2/17	-
3	LFA	A	306	-	-	1/2/2/17	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	301	RET	C8-C7-C6	2.25	133.51	127.20

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	321	LFA	C3-C4-C5-C6
3	A	302	LFA	C9-C10-C11-C12
3	A	310	LFA	C3-C4-C5-C6
3	A	317	LFA	C6-C7-C8-C9
3	A	311	LFA	C6-C7-C8-C9

There are no ring outliers.

9 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	325	LFA	1	0
3	A	316	LFA	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	RET	2	0
3	A	313	LFA	1	0
3	A	302	LFA	1	0
3	A	326	LFA	1	0
3	A	307	LFA	1	0
3	A	318	LFA	2	0
3	A	306	LFA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LFA A 312



Ligand LFA A 325



Ligand LFA A 319



Ligand LFA A 316



Ligand LFA A 308



Ligand LFA A 303



Ligand LFA A 305



Ligand RET A 301



Ligand LFA A 314



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 311



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 327



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 321



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 310



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 317



Bond lengths



Bond angles



Torsions

Rings

Ligand LFA A 302



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 326



Bond lengths



Bond angles



Torsions



Rings

Ligand LFA A 307



Ligand LFA A 323



Ligand LFA A 318



Ligand LFA A 306



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	265/290 (91%)	0.15	13 (4%) 29 31	37, 51, 75, 102	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	GLY	3.1
1	A	229	GLY	3.0
1	A	270	SER	2.8
1	A	183	TRP	2.7
1	A	93	GLY	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	LFA	A	322	6/20	0.38	0.39	74,79,93,95	0
3	LFA	A	306	5/20	0.41	0.30	88,98,99,103	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	LFA	A	311	12/20	0.42	0.25	64,84,93,94	0
3	LFA	A	323	10/20	0.42	0.30	78,87,93,95	0
3	LFA	A	314	10/20	0.47	0.40	72,87,103,109	0
3	LFA	A	318	9/20	0.47	0.46	72,78,88,90	0
3	LFA	A	320	5/20	0.52	0.31	62,69,77,79	0
4	NA	A	328	1/1	0.52	0.41	64,64,64,64	0
3	LFA	A	316	11/20	0.53	0.41	64,88,90,91	0
3	LFA	A	309	8/20	0.53	0.40	87,105,110,111	0
3	LFA	A	304	6/20	0.60	0.24	63,72,74,74	0
3	LFA	A	325	9/20	0.61	0.53	75,83,97,97	0
3	LFA	A	324	3/20	0.62	0.33	72,72,92,93	0
3	LFA	A	302	16/20	0.63	0.23	64,77,85,89	0
3	LFA	A	312	16/20	0.66	0.21	75,82,92,98	0
3	LFA	A	313	5/20	0.66	0.31	60,64,71,77	0
3	LFA	A	321	4/20	0.66	0.55	71,73,80,81	0
3	LFA	A	317	9/20	0.66	0.25	72,76,87,90	0
3	LFA	A	327	12/20	0.67	0.24	50,67,70,71	0
3	LFA	A	315	5/20	0.69	0.30	63,63,68,73	0
3	LFA	A	305	6/20	0.71	0.26	66,71,87,89	0
3	LFA	A	307	13/20	0.73	0.30	43,57,68,72	0
3	LFA	A	326	10/20	0.76	0.28	48,61,76,77	0
3	LFA	A	319	6/20	0.77	0.27	57,61,64,70	0
3	LFA	A	310	8/20	0.79	0.28	61,63,68,74	0
3	LFA	A	308	10/20	0.87	0.15	63,66,70,70	0
3	LFA	A	303	8/20	0.89	0.17	47,50,50,56	8
2	RET	A	301	20/21	0.91	0.16	38,46,53,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LFA A 306:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 311:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 323:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 314:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 318:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NA A 328:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around LFA A 316:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 325:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 312:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 321:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 317:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 327:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 305:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 307:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 326:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 319:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LFA A 310:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 308:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LFA A 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around RET A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.