



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2023 – 03:40 PM EDT

PDB ID : 5TLK
Title : COMPLEX BETWEEN HUMAN CD27 AND FAB FRAGMENTS OF ANTI-BODIES M2177 AND H2191
Authors : Teplyakov, A.; Obmolova, G.; Malia, T.; Gilliland, G.L.
Deposited on : 2016-10-11
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	 8% 90% 8%
1	E	218	 8% 88% 10%
2	B	229	 3% 85% 8% 7%
2	F	229	 24% 86% 6% 7%
3	C	218	 % 90% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	218	
4	D	226	
4	H	226	
5	X	109	
5	Y	109	
6	I	2	
6	J	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NAG	I	2	-	-	-	X
6	NAG	J	2	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 14287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M2177 LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	Total 1647	C 1029	N 274	O 339	S 5	0	0	0
1	E	215	Total 1647	C 1029	N 274	O 339	S 5	0	0	0

- Molecule 2 is a protein called M2177 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	213	Total 1598	C 1017	N 261	O 314	S 6	0	0	0
2	F	212	Total 1592	C 1014	N 260	O 312	S 6	0	0	0

- Molecule 3 is a protein called H2191 LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	215	Total 1672	C 1048	N 286	O 332	S 6	0	0	0
3	G	215	Total 1672	C 1048	N 286	O 332	S 6	0	0	0

- Molecule 4 is a protein called H2191 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	212	Total 1584	C 1001	N 268	O 309	S 6	0	0	0
4	H	211	Total 1578	C 998	N 267	O 307	S 6	0	0	0

- Molecule 5 is a protein called CD27 antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	X	75	Total	C	N	O	S	0	0	0
			583	351	113	106	13			
5	Y	75	Total	C	N	O	S	0	0	0
			583	351	113	106	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	102	GLY	-	expression tag	UNP P26842
X	103	GLY	-	expression tag	UNP P26842
X	104	HIS	-	expression tag	UNP P26842
X	105	HIS	-	expression tag	UNP P26842
X	106	HIS	-	expression tag	UNP P26842
X	107	HIS	-	expression tag	UNP P26842
X	108	HIS	-	expression tag	UNP P26842
X	109	HIS	-	expression tag	UNP P26842
Y	102	GLY	-	expression tag	UNP P26842
Y	103	GLY	-	expression tag	UNP P26842
Y	104	HIS	-	expression tag	UNP P26842
Y	105	HIS	-	expression tag	UNP P26842
Y	106	HIS	-	expression tag	UNP P26842
Y	107	HIS	-	expression tag	UNP P26842
Y	108	HIS	-	expression tag	UNP P26842
Y	109	HIS	-	expression tag	UNP P26842

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
6	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	14	Total	O	0	0
			14	14		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	17	Total 17	O 17	0	0
7	C	12	Total 12	O 12	0	0
7	D	15	Total 15	O 15	0	0
7	E	1	Total 1	O 1	0	0
7	G	3	Total 3	O 3	0	0
7	H	8	Total 8	O 8	0	0
7	X	4	Total 4	O 4	0	0
7	Y	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: M2177 LIGHT CHAIN

Chain A: 




- Molecule 1: M2177 LIGHT CHAIN

Chain E: 




- Molecule 2: M2177 HEAVY CHAIN

Chain B: 



- Molecule 2: M2177 HEAVY CHAIN


Chain F: 




● Molecule 3: H2191 LIGHT CHAIN





● Molecule 3: H2191 LIGHT CHAIN





● Molecule 4: H2191 HEAVY CHAIN




● Molecule 4: H2191 HEAVY CHAIN



● Molecule 5: CD27 antigen



● Molecule 5: CD27 antigen



GLU
CYS
THR
GLU
CYS
ASP
GLY
GLY
HIS
HIS
HIS
HIS
HIS
HIS

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1
MAG2

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50%  50%

MAG1
MAG2

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	139.71Å 52.96Å 140.60Å 90.00° 109.52° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 29.21 – 2.69	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.70) 98.6 (29.21-2.69)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.213 , 0.263 0.214 , 0.264	Depositor DCC
R_{free} test set	1109 reflections (2.06%)	wwPDB-VP
Wilson B-factor (Å ²)	56.4	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14287	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.52 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2904e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1683	0.61	1/2288 (0.0%)
1	E	0.40	0/1683	0.59	0/2288
2	B	0.40	0/1633	0.59	1/2225 (0.0%)
2	F	0.44	0/1627	0.60	0/2217
3	C	0.39	0/1711	0.61	0/2320
3	G	0.40	0/1711	0.61	0/2320
4	D	0.39	0/1616	0.62	0/2201
4	H	0.40	0/1610	0.60	0/2193
5	X	0.39	0/598	0.59	0/807
5	Y	0.38	0/598	0.59	0/807
All	All	0.40	0/14470	0.60	2/19666 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	185	LEU	CA-CB-CG	5.08	126.98	115.30
1	A	158	LEU	CA-CB-CG	5.02	126.84	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1647	0	1583	10	0
1	E	1647	0	1583	10	0
2	B	1598	0	1554	7	0
2	F	1592	0	1549	7	0
3	C	1672	0	1629	8	0
3	G	1672	0	1629	10	0
4	D	1584	0	1544	5	0
4	H	1578	0	1539	8	0
5	X	583	0	529	10	0
5	Y	583	0	529	7	0
6	I	28	0	25	0	0
6	J	28	0	25	0	0
7	A	14	0	0	0	0
7	B	17	0	0	0	0
7	C	12	0	0	0	0
7	D	15	0	0	0	0
7	E	1	0	0	0	0
7	G	3	0	0	0	0
7	H	8	0	0	0	0
7	X	4	0	0	0	0
7	Y	1	0	0	0	0
All	All	14287	0	13718	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:72:ARG:HE	4:D:74:ASN:HD21	1.26	0.80
1:A:44:PRO:HG3	1:A:169:GLU:HG2	1.67	0.76
2:B:50:ARG:HH11	2:B:59:ASN:HD22	1.37	0.73
2:F:50:ARG:HH11	2:F:59:ASN:HD22	1.46	0.63
1:E:153:LYS:HB2	1:E:197:ALA:HB3	1.83	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/218 (98%)	206 (97%)	7 (3%)	0	100	100
1	E	213/218 (98%)	203 (95%)	10 (5%)	0	100	100
2	B	209/229 (91%)	201 (96%)	7 (3%)	1 (0%)	29	54
2	F	208/229 (91%)	193 (93%)	15 (7%)	0	100	100
3	C	213/218 (98%)	208 (98%)	5 (2%)	0	100	100
3	G	213/218 (98%)	202 (95%)	11 (5%)	0	100	100
4	D	208/226 (92%)	205 (99%)	3 (1%)	0	100	100
4	H	207/226 (92%)	200 (97%)	7 (3%)	0	100	100
5	X	71/109 (65%)	67 (94%)	3 (4%)	1 (1%)	11	28
5	Y	71/109 (65%)	66 (93%)	3 (4%)	2 (3%)	5	11
All	All	1826/2000 (91%)	1751 (96%)	71 (4%)	4 (0%)	47	73

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	Y	14	ALA
5	X	14	ALA
5	Y	8	PRO
2	B	103	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/187 (99%)	181 (98%)	4 (2%)	52 79
1	E	185/187 (99%)	176 (95%)	9 (5%)	25 52
2	B	176/190 (93%)	173 (98%)	3 (2%)	60 84
2	F	175/190 (92%)	169 (97%)	6 (3%)	37 66
3	C	191/193 (99%)	183 (96%)	8 (4%)	30 58
3	G	191/193 (99%)	182 (95%)	9 (5%)	26 54
4	D	175/188 (93%)	170 (97%)	5 (3%)	42 71
4	H	174/188 (93%)	168 (97%)	6 (3%)	37 66
5	X	67/95 (70%)	63 (94%)	4 (6%)	19 42
5	Y	67/95 (70%)	65 (97%)	2 (3%)	41 70
All	All	1586/1706 (93%)	1530 (96%)	56 (4%)	36 65

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	194	LYS
5	Y	77	THR
3	G	1	ASP
5	Y	18	LEU
4	H	213	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	93	GLN
2	F	82	GLN
5	X	15	GLN
2	F	59	ASN
3	G	93	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	F	1	2	7,8,9	0.56	0	9,10,12	0.97	0
2	PCA	B	1	2	7,8,9	0.55	0	9,10,12	1.15	0
4	PCA	D	1	4	7,8,9	0.47	0	9,10,12	1.27	1 (11%)
4	PCA	H	1	4	7,8,9	0.50	0	9,10,12	1.30	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	F	1	2	-	0/0/11/13	0/1/1/1
2	PCA	B	1	2	-	0/0/11/13	0/1/1/1
4	PCA	D	1	4	-	0/0/11/13	0/1/1/1
4	PCA	H	1	4	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1	PCA	OE-CD-CG	-2.46	122.46	126.76
4	D	1	PCA	OE-CD-CG	-2.25	122.84	126.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	I	1	6,5	14,14,15	0.56	0	17,19,21	1.11	3 (17%)
6	NAG	I	2	6	14,14,15	0.52	0	17,19,21	1.05	1 (5%)
6	NAG	J	1	6,5	14,14,15	0.65	0	17,19,21	0.92	0
6	NAG	J	2	6	14,14,15	0.54	0	17,19,21	1.26	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	I	1	6,5	-	0/6/23/26	0/1/1/1
6	NAG	I	2	6	-	1/6/23/26	0/1/1/1
6	NAG	J	1	6,5	-	2/6/23/26	0/1/1/1
6	NAG	J	2	6	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	I	2	NAG	C1-O5-C5	3.11	116.41	112.19
6	J	2	NAG	C1-O5-C5	2.84	116.04	112.19
6	J	2	NAG	C4-C3-C2	2.74	115.04	111.02
6	I	1	NAG	C4-C3-C2	2.43	114.57	111.02
6	I	1	NAG	C2-N2-C7	2.08	125.87	122.90

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	J	1	NAG	C4-C5-C6-O6
6	J	1	NAG	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	J	2	NAG	C4-C5-C6-O6
6	I	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/218 (98%)	0.09	1 (0%) 91 92	29, 45, 63, 78	0
1	E	215/218 (98%)	0.62	17 (7%) 12 10	53, 72, 106, 117	0
2	B	212/229 (92%)	0.15	7 (3%) 46 46	25, 47, 77, 92	0
2	F	211/229 (92%)	1.29	54 (25%) 0 0	57, 89, 120, 133	0
3	C	215/218 (98%)	0.05	3 (1%) 75 77	29, 48, 74, 84	0
3	G	215/218 (98%)	0.28	4 (1%) 66 69	40, 65, 87, 102	0
4	D	211/226 (93%)	0.06	3 (1%) 75 77	31, 50, 77, 98	0
4	H	210/226 (92%)	0.24	8 (3%) 40 39	38, 58, 78, 94	0
5	X	75/109 (68%)	0.39	7 (9%) 8 6	33, 51, 74, 83	0
5	Y	75/109 (68%)	0.91	10 (13%) 3 2	50, 75, 90, 95	0
All	All	1854/2000 (92%)	0.37	114 (6%) 21 20	25, 58, 99, 133	0

The worst 5 of 114 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	200	THR	6.9
2	F	196	LEU	6.5
2	F	217	LYS	5.3
4	H	194	GLY	5.3
2	F	192	PRO	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PCA	H	1	8/9	0.82	0.32	72,75,78,80	0
2	PCA	F	1	8/9	0.83	0.43	76,87,91,94	0
2	PCA	B	1	8/9	0.83	0.41	67,78,88,95	0
4	PCA	D	1	8/9	0.92	0.15	55,67,70,71	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NAG	J	2	14/15	0.61	0.46	120,131,134,140	0
6	NAG	I	2	14/15	0.65	0.43	116,131,139,144	0
6	NAG	J	1	14/15	0.74	0.39	103,110,123,124	0
6	NAG	I	1	14/15	0.80	0.27	79,90,100,114	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.