



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 17, 2023 – 06:59 PM EDT

PDB ID : 1TU5
Title : Crystal structure of bovine plasma copper-containing amine oxidase
Authors : Lunelli, M.; Di Paolo, M.L.; Biadene, M.; Calderone, V.; Scarpa, M.; Battistutta, R.; Rigo, A.; Zanotti, G.
Deposited on : 2004-06-24
Resolution : 2.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 130704 | 5509 (2.40-2.36) |
| Clashscore | 141614 | 6082 (2.40-2.36) |
| Ramachandran outliers | 138981 | 5973 (2.40-2.36) |
| Sidechain outliers | 138945 | 5975 (2.40-2.36) |
| RSRZ outliers | 127900 | 5397 (2.40-2.36) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | A | 746 | <div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 59%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: grey;"></div> </div> |
| 1 | B | 746 | <div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 61%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: grey;"></div> </div> |
| 2 | C | 3 | <div style="display: flex; align-items: center;"> <div style="width: 33%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: orange;"></div> </div> |
| 2 | D | 3 | <div style="display: flex; align-items: center;"> <div style="width: 67%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: orange;"></div> </div> |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 2 | NAG | C | 3 | - | - | - | X |
| 2 | NAG | D | 1 | X | - | - | - |
| 2 | NAG | D | 3 | - | - | - | X |
| 3 | NAG | A | 803 | - | - | - | X |
| 3 | NAG | A | 804 | X | - | - | X |
| 3 | NAG | B | 804 | - | - | - | X |

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper amine oxidase, liver isozyme.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 633 | 5018 | 3217 | 862 | 920 | 19 | 0 | 0 | 0 |
| 1 | B | 633 | 5018 | 3217 | 862 | 920 | 19 | 0 | 0 | 0 |

There are 2 discrepancies between the modelled and reference sequences:


| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| A | 470 | TPQ | TYR | modified residue | UNP Q29437 |
| B | 470 | TPQ | TYR | modified residue | UNP Q29437 |

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|---------|-------|
| | | | Total | C | N | O | | | |
| 2 | C | 3 | 42 | 24 | 3 | 15 | 0 | 0 | 0 |
| 2 | D | 3 | 42 | 24 | 3 | 15 | 0 | 0 | 0 |

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---|---------|---------|
| | | | Total | C | N | O | | |
| 3 | A | 1 | 14 | 8 | 1 | 5 | 0 | 0 |
| 3 | A | 1 | 14 | 8 | 1 | 5 | 0 | 0 |
| 3 | B | 1 | 14 | 8 | 1 | 5 | 0 | 0 |

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| | | | Total | Cu | | |
| 4 | A | 1 | 1 | 1 | 0 | 0 |
| 4 | B | 1 | 1 | 1 | 0 | 0 |

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| | | | Total | Ca | | |
| 5 | A | 2 | 2 | 2 | 0 | 0 |
| 5 | B | 2 | 2 | 2 | 0 | 0 |

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 6 | A | 1 | Total Cl 1 1 | 0 | 0 |
| 6 | B | 2 | Total Cl 2 2 | 0 | 0 |

- Molecule 7 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 7 | A | 209 | Total O 209 209 | 0 | 0 |
| 7 | B | 201 | Total O 201 201 | 0 | 0 |

4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 77.68Å 131.19Å 134.00Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 25.00 – 2.37 27.85 – 2.37 | Depositor EDS |
| % Data completeness (in resolution range) | 93.7 (25.00-2.37) 93.8 (27.85-2.37) | Depositor EDS |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | 0.11 | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.72 (at 2.36Å) | Xtrriage |
| Refinement program | CNS 1.1 | Depositor |
| R, R_{free} | 0.205 , 0.237 0.203 , 0.231 | Depositor DCC |
| R_{free} test set | 3719 reflections (7.07%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 27.9 | Xtrriage |
| Anisotropy | 0.217 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.38 , 49.7 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$ | Xtrriage |
| Estimated twinning fraction | 0.018 for -h,l,k | Xtrriage |
| F_o, F_c correlation | 0.93 | EDS |
| Total number of atoms | 10581 | wwPDB-VP |
| Average B, all atoms (Å ²) | 29.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPQ, CA, NAG, CU, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.57 | 2/5156 (0.0%) | 0.83 | 7/7025 (0.1%) |
| 1 | B | 0.55 | 0/5156 | 0.79 | 6/7025 (0.1%) |
| All | All | 0.56 | 2/10312 (0.0%) | 0.81 | 13/14050 (0.1%) |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|-------|-------------|----------|
| 1 | A | 142 | VAL | CB-CG1 | -6.81 | 1.38 | 1.52 |
| 1 | A | 142 | VAL | CB-CG2 | 5.38 | 1.64 | 1.52 |

The worst 5 of 13 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 1 | A | 448 | SER | N-CA-C | -9.93 | 84.20 | 111.00 |
| 1 | A | 446 | PHE | N-CA-C | 8.70 | 134.49 | 111.00 |
| 1 | A | 447 | LEU | CA-CB-CG | -8.53 | 95.69 | 115.30 |
| 1 | B | 448 | SER | N-CA-C | -8.41 | 88.30 | 111.00 |
| 1 | B | 446 | PHE | N-CA-C | 6.98 | 129.84 | 111.00 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 5018 | 0 | 4801 | 163 | 0 |
| 1 | B | 5018 | 0 | 4802 | 138 | 0 |
| 2 | C | 42 | 0 | 37 | 2 | 0 |
| 2 | D | 42 | 0 | 37 | 5 | 0 |
| 3 | A | 28 | 0 | 26 | 4 | 0 |
| 3 | B | 14 | 0 | 13 | 0 | 0 |
| 4 | A | 1 | 0 | 0 | 0 | 0 |
| 4 | B | 1 | 0 | 0 | 0 | 0 |
| 5 | A | 2 | 0 | 0 | 0 | 0 |
| 5 | B | 2 | 0 | 0 | 0 | 0 |
| 6 | A | 1 | 0 | 0 | 0 | 0 |
| 6 | B | 2 | 0 | 0 | 0 | 0 |
| 7 | A | 209 | 0 | 0 | 5 | 0 |
| 7 | B | 201 | 0 | 0 | 3 | 0 |
| All | All | 10581 | 0 | 9716 | 289 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 289 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|-----------------|--------------------------|-------------------|
| 1:A:591:LYS:HD3 | 1:A:591:LYS:N | 1.63 | 1.09 |
| 1:B:232:ILE:HG12 | 1:B:240:HIS:CD2 | 1.87 | 1.08 |
| 1:B:591:LYS:H | 1:B:591:LYS:CD | 1.62 | 1.07 |
| 1:B:591:LYS:HD2 | 1:B:591:LYS:N | 1.65 | 1.07 |
| 1:A:591:LYS:H | 1:A:591:LYS:CD | 1.63 | 1.02 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|---------|----------|-------------|----|
| 1 | A | 628/746 (84%) | 592 (94%) | 33 (5%) | 3 (0%) | 29 | 39 |
| 1 | B | 628/746 (84%) | 594 (95%) | 31 (5%) | 3 (0%) | 29 | 39 |
| All | All | 1256/1492 (84%) | 1186 (94%) | 64 (5%) | 6 (0%) | 29 | 39 |

5 of 6 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 233 | THR |
| 1 | A | 81 | ASP |
| 1 | A | 235 | GLY |
| 1 | A | 233 | THR |
| 1 | B | 235 | GLY |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|-----------|----------|-------------|----|
| 1 | A | 530/623 (85%) | 491 (93%) | 39 (7%) | 13 | 19 |
| 1 | B | 530/623 (85%) | 494 (93%) | 36 (7%) | 16 | 23 |
| All | All | 1060/1246 (85%) | 985 (93%) | 75 (7%) | 14 | 21 |

5 of 75 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 451 | PHE |
| 1 | B | 673 | LEU |
| 1 | B | 471 | ASP |
| 1 | B | 572 | GLU |
| 1 | A | 473 | VAL |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 456 | GLN |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 692 | ASN |
| 1 | B | 712 | GLN |
| 1 | B | 559 | GLN |
| 1 | B | 184 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 1 | TPQ | B | 470 | 1 | 13,14,15 | 2.77 | 7 (53%) | 15,19,21 | 1.33 | 2 (13%) |
| 1 | TPQ | A | 470 | 1 | 13,14,15 | 2.81 | 7 (53%) | 15,19,21 | 1.22 | 1 (6%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 1 | TPQ | B | 470 | 1 | - | 2/5/22/24 | 0/1/1/1 |
| 1 | TPQ | A | 470 | 1 | - | 2/5/22/24 | 0/1/1/1 |

The worst 5 of 14 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 1 | B | 470 | TPQ | C1-C2 | -7.27 | 1.38 | 1.49 |
| 1 | A | 470 | TPQ | C1-C2 | -7.13 | 1.38 | 1.49 |
| 1 | A | 470 | TPQ | O5-C5 | 3.33 | 1.33 | 1.24 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 1 | B | 470 | TPQ | O5-C5 | 3.22 | 1.33 | 1.24 |
| 1 | A | 470 | TPQ | O2-C2 | 3.20 | 1.33 | 1.24 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 1 | A | 470 | TPQ | CB-CA-C | -3.02 | 105.81 | 111.47 |
| 1 | B | 470 | TPQ | CB-CA-C | -2.86 | 106.10 | 111.47 |
| 1 | B | 470 | TPQ | C6-C1-C2 | 2.16 | 120.30 | 118.64 |

There are no chirality outliers.

All (4) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|------------|
| 1 | A | 470 | TPQ | N-CA-CB-C1 |
| 1 | B | 470 | TPQ | N-CA-CB-C1 |
| 1 | B | 470 | TPQ | C-CA-CB-C1 |
| 1 | A | 470 | TPQ | C-CA-CB-C1 |

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates i

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 2 | NAG | C | 1 | 1,2 | 14,14,15 | 0.57 | 0 | 17,19,21 | 0.98 | 1 (5%) |
| 2 | NAG | C | 2 | 2 | 14,14,15 | 0.69 | 0 | 17,19,21 | 0.61 | 0 |
| 2 | NAG | C | 3 | 2 | 14,14,15 | 1.05 | 1 (7%) | 17,19,21 | 1.07 | 1 (5%) |
| 2 | NAG | D | 1 | 1,2 | 14,14,15 | 0.82 | 1 (7%) | 17,19,21 | 1.17 | 2 (11%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 2 | NAG | D | 2 | 2 | 14,14,15 | 0.54 | 0 | 17,19,21 | 0.91 | 1 (5%) |
| 2 | NAG | D | 3 | 2 | 14,14,15 | 0.73 | 0 | 17,19,21 | 0.85 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 2 | NAG | C | 1 | 1,2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | NAG | C | 2 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | NAG | C | 3 | 2 | - | 2/6/23/26 | 0/1/1/1 |
| 2 | NAG | D | 1 | 1,2 | 1/1/5/7 | 4/6/23/26 | 0/1/1/1 |
| 2 | NAG | D | 2 | 2 | - | 2/6/23/26 | 0/1/1/1 |
| 2 | NAG | D | 3 | 2 | - | 5/6/23/26 | 0/1/1/1 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 2 | C | 3 | NAG | C1-C2 | 3.35 | 1.57 | 1.52 |
| 2 | D | 1 | NAG | O5-C5 | 2.02 | 1.47 | 1.43 |

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 2 | D | 1 | NAG | C1-O5-C5 | 3.00 | 116.26 | 112.19 |
| 2 | C | 3 | NAG | C3-C4-C5 | -2.52 | 105.75 | 110.24 |
| 2 | C | 1 | NAG | C1-C2-N2 | -2.46 | 106.28 | 110.49 |
| 2 | D | 2 | NAG | C2-N2-C7 | -2.36 | 119.55 | 122.90 |
| 2 | D | 1 | NAG | C2-N2-C7 | -2.15 | 119.84 | 122.90 |

All (1) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 2 | D | 1 | NAG | C1 |

5 of 13 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 2 | C | 3 | NAG | C8-C7-N2-C2 |

Continued on next page...

Continued from previous page...


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 2 | C | 3 | NAG | O7-C7-N2-C2 |
| 2 | D | 2 | NAG | O5-C5-C6-O6 |
| 2 | D | 1 | NAG | C4-C5-C6-O6 |
| 2 | D | 1 | NAG | O5-C5-C6-O6 |


There are no ring outliers.

4 monomers are involved in 7 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 2 | C | 1 | NAG | 1 | 0 |
| 2 | C | 3 | NAG | 1 | 0 |
| 2 | D | 3 | NAG | 5 | 0 |
| 2 | D | 2 | NAG | 4 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 3 | NAG | B | 804 | 1 | 14,14,15 | 0.74 | 0 | 17,19,21 | 0.81 | 1 (5%) |
| 3 | NAG | A | 804 | 1 | 14,14,15 | 0.58 | 0 | 17,19,21 | 0.69 | 0 |
| 3 | NAG | A | 803 | 1 | 14,14,15 | 0.72 | 0 | 17,19,21 | 1.04 | 1 (5%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 3 | NAG | B | 804 | 1 | - | 2/6/23/26 | 0/1/1/1 |
| 3 | NAG | A | 804 | 1 | 1/1/5/7 | 4/6/23/26 | 0/1/1/1 |
| 3 | NAG | A | 803 | 1 | - | 3/6/23/26 | 0/1/1/1 |

There are no bond length outliers.

All (2) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 3 | A | 803 | NAG | O5-C1-C2 | -2.12 | 107.94 | 111.29 |
| 3 | B | 804 | NAG | C2-N2-C7 | -2.05 | 119.99 | 122.90 |

All (1) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 3 | A | 804 | NAG | C1 |

5 of 9 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 3 | A | 803 | NAG | C3-C2-N2-C7 |
| 3 | A | 804 | NAG | C8-C7-N2-C2 |
| 3 | A | 804 | NAG | O7-C7-N2-C2 |
| 3 | B | 804 | NAG | C8-C7-N2-C2 |
| 3 | B | 804 | NAG | O7-C7-N2-C2 |

There are no ring outliers.

2 monomers are involved in 4 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 3 | A | 804 | NAG | 2 | 0 |
| 3 | A | 803 | NAG | 2 | 0 |

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------------|-----------------------|-------|
| 1 | A | 632/746 (84%) | 0.30 | 37 (5%) 22 24 | 11, 25, 50, 114 | 0 |
| 1 | B | 632/746 (84%) | 0.37 | 36 (5%) 23 26 | 9, 25, 48, 115 | 0 |
| All | All | 1264/1492 (84%) | 0.33 | 73 (5%) 23 25 | 9, 25, 49, 115 | 0 |

The worst 5 of 73 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | B | 446 | PHE | 16.7 |
| 1 | A | 447 | LEU | 15.1 |
| 1 | B | 451 | PHE | 14.7 |
| 1 | A | 444 | SER | 14.6 |
| 1 | A | 450 | TYR | 14.6 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 1 | TPQ | B | 470 | 14/15 | 0.82 | 0.21 | 40,53,55,56 | 0 |
| 1 | TPQ | A | 470 | 14/15 | 0.85 | 0.18 | 40,50,52,52 | 0 |


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|-----------------------------|-------|
| 2 | NAG | C | 3 | 14/15 | 0.68 | 0.47 | 63,70,73,74 | 0 |
| 2 | NAG | D | 3 | 14/15 | 0.69 | 0.41 | 61,64,66,67 | 0 |
| 2 | NAG | C | 2 | 14/15 | 0.89 | 0.15 | 28,44,49,55 | 0 |
| 2 | NAG | D | 2 | 14/15 | 0.91 | 0.22 | 35,42,49,55 | 0 |
| 2 | NAG | C | 1 | 14/15 | 0.93 | 0.13 | 29,30,32,36 | 0 |
| 2 | NAG | D | 1 | 14/15 | 0.94 | 0.12 | 29,31,35,35 | 0 |

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|------|-------|------|------|----------------------------|-------|
| 3 | NAG | B | 804 | 14/15 | 0.70 | 0.50 | 57,60,62,63 | 0 |
| 3 | NAG | A | 803 | 14/15 | 0.77 | 0.42 | 48,49,51,52 | 0 |
| 3 | NAG | A | 804 | 14/15 | 0.80 | 0.53 | 64,67,67,68 | 0 |
| 6 | CL | A | 904 | 1/1 | 0.94 | 0.08 | 32,32,32,32 | 0 |
| 5 | CA | A | 903 | 1/1 | 0.98 | 0.03 | 19,19,19,19 | 0 |
| 6 | CL | B | 1904 | 1/1 | 0.98 | 0.09 | 32,32,32,32 | 0 |
| 6 | CL | B | 1905 | 1/1 | 0.98 | 0.10 | 25,25,25,25 | 0 |
| 5 | CA | B | 1903 | 1/1 | 0.99 | 0.05 | 23,23,23,23 | 0 |
| 5 | CA | A | 902 | 1/1 | 0.99 | 0.10 | 18,18,18,18 | 0 |
| 4 | CU | B | 901 | 1/1 | 0.99 | 0.07 | 23,23,23,23 | 0 |
| 5 | CA | B | 1902 | 1/1 | 0.99 | 0.06 | 23,23,23,23 | 0 |
| 4 | CU | A | 901 | 1/1 | 1.00 | 0.08 | 21,21,21,21 | 0 |

6.5 Other polymers [i](#)

There are no such residues in this entry.