wwPDB NMR Structure Validation Summary Report (i)

Oct 31, 2021 – 06:50 PM EDT

PDB ID	:	1U4A
Title	:	Solution structure of human SUMO-3 C47S
Authors	:	Ding, H.; Xu, Y.; Dai, H.; Tang, Y.; Wu, J.; Shi, Y.
Deposited on	:	2004-07-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.23.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	87	55% 26% • 7% 9%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:15-A:87 (73)	0.42	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 8, 10, 11, 12, 14, 15, 16, 17, 18, 19
2	2, 5, 6, 7
3	9, 20
Single-model clusters	13

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1256 atoms, of which 623 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Ubiquitin-like protein SMT3A.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	79	1256	390	623	115	125	3	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	SER	CYS	engineered mutation	UNP P55854
A	93	LEU	-	expression tag	UNP P55854
A	94	GLU	-	expression tag	UNP P55854
A	95	HIS	-	expression tag	UNP P55854
A	96	HIS	-	expression tag	UNP P55854
A	97	HIS	-	expression tag	UNP P55854
A	98	HIS	-	expression tag	UNP P55854
A	99	HIS	-	expression tag	UNP P55854
A	100	HIS	-	expression tag	UNP P55854

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Ubiquitin-like protein SMT3A

Chain A: 55% 26% • 7% 9%

N14 I17 N18 L19 K20 K34 R35 H36 T37 P38 L39 S40 K41 L42 M43 R49 Q50 G51 L52 S53 Q56 I57 R58 D62 I66 D70 T71 E78 D79 T82 I83 D84 V85 Q88 Q89 T90 G91 G92 LEU GLU HIS HIS HIS HIS HIS

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 18. Colouring as in section 4.1 above.

- Molecule 1: Ubiquitin-like protein SMT3A

Chain A: 54% 24% 6% 7% 9%

N14 D15 H16 L19 K20 Q30 K34 R35 L39 S40 K41 L42 M43 G51 L52 S53 Q56 I57 R58 F59 D62 I66 N67 E68 T69 D70 T71 E78 D79 E80 D81 T82 I83 Q87 Q88 Q89 T90 G91 G92 LEU GLU HIS HIS HIS HIS HIS HIS

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
MOLMOL	structure solution	2k.2
MOLMOL	refinement	2k.2

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	592	586	584	9±3
All	All	11840	11720	11680	181

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 65 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:19:LEU:HD23	1:A:83:ILE:HD11	0.82	1.51	2	3
1:A:19:LEU:HD12	1:A:81:ASP:HB3	0.70	1.63	7	1
1:A:20:LYS:HB2	1:A:82:THR:HG22	0.69	1.62	2	4
1:A:39:LEU:HD12	1:A:70:ASP:HB2	0.69	1.63	17	8
1:A:21:VAL:HG22	1:A:83:ILE:HD11	0.65	1.69	14	6

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	73/87 (84%)	61±2 (84±2%)	9±2 (13±2%)	3±1 (4±2%)	6 34
All	All	1460/1740 (84%)	1223 (84%)	185 (13%)	52 (4%)	6 34

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	71	THR	20
1	A	51	GLY	13
1	A	79	ASP	8
1	A	65	PRO	3
1	A	36	HIS	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	66/78 (85%)	46±3 (70±4%)	20±3 (30±4%)	1 16
All	All	1320/1560 (85%)	924 (70%)	396 (30%)	1 16

5 of 52 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	71	THR	19
1	A	34	LYS	18
1	A	56	GLN	16
1	A	41	LYS	14
1	A	35	ARG	13

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided