



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 04:15 PM EDT

PDB ID : 5UBP
Title : TREX2 M-region
Authors : Stewart, M.; Gordon, J.
Deposited on : 2016-12-21
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>6%</div> <div>94%</div> </div>
2	B	455	<div> <div>4%</div> <div>96%</div> </div>
3	C	89	<div> <div>27%</div> <div>52%</div> <div>46%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16570 atoms, of which 8013 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine permease transcriptional regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	469	Total	C	H	N	O	S	0	3	0
			7756	2459	3892	671	714	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	SER	-	expression tag	UNP B3LGC5
A	70	GLY	-	expression tag	UNP B3LGC5

- Molecule 2 is a protein called Nuclear mRNA export protein THP1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	455	Total	C	H	N	O	S	0	2	0
			7494	2402	3766	646	662	18			

- Molecule 3 is a protein called 26S proteasome complex subunit SEM1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	48	Total	C	H	N	O		0	0	0
			774	258	355	64	97				


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	326	Total	O	0	0
			326	326		
4	B	200	Total	O	0	0
			200	200		
4	C	20	Total	O	0	0
			20	20		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Leucine permease transcriptional regulator



- Molecule 2: Nuclear mRNA export protein THP1



- Molecule 3: 26S proteasome complex subunit SEM1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.37Å 86.63Å 168.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.95 – 2.30 19.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.95-2.30) 100.0 (19.95-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.30Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.178 , 0.217 0.180 , 0.219	Depositor DCC
R_{free} test set	2607 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16570	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/3956	0.42	0/5338
2	B	0.26	0/3821	0.40	0/5191
3	C	0.24	0/426	0.36	0/575
All	All	0.26	0/8203	0.41	0/11104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3864	3892	3883	10	0
2	B	3728	3766	3768	10	0
3	C	419	355	355	2	0
4	A	326	0	0	3	0
4	B	200	0	0	0	0
4	C	20	0	0	1	0
All	All	8557	8013	8006	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128[B]:GLN:OE1	4:A:601:HOH:O	1.94	0.86
1:A:310:ARG:NH2	4:A:608:HOH:O	2.30	0.64
1:A:420:GLU:OE2	1:A:420:GLU:N	2.44	0.51
1:A:81:LYS:NZ	1:A:193:ALA:O	2.45	0.50
2:B:359:GLY:O	2:B:428:LYS:NZ	2.32	0.49
2:B:338:LEU:N	2:B:339:PRO:CD	2.76	0.49
2:B:340:MET:HE3	2:B:407:LEU:HD23	1.95	0.48
1:A:274:THR:OG1	4:A:602:HOH:O	2.20	0.48
1:A:85:TYR:HA	1:A:117:GLU:HA	1.96	0.47
2:B:110:ILE:HB	2:B:111:PRO:HD3	1.97	0.47
1:A:404:ARG:HD2	2:B:336:GLU:HB2	1.98	0.46
3:C:41:ASP:O	4:C:101:HOH:O	2.21	0.45
2:B:353:SER:HA	2:B:357:GLU:HB2	1.99	0.45
1:A:318:ILE:O	1:A:322:MET:HG2	2.18	0.43
2:B:258:GLN:O	2:B:261:THR:OG1	2.27	0.43
2:B:240:PHE:HB2	2:B:274:THR:HG22	2.01	0.42
1:A:329:PHE:CE1	1:A:331:LEU:HA	2.55	0.41
2:B:255[B]:LEU:HD11	2:B:261:THR:CG2	2.50	0.41
3:C:40:ILE:HD12	3:C:40:ILE:N	2.35	0.41
1:A:331:LEU:HD12	1:A:332:GLN:N	2.36	0.40
2:B:453:VAL:HG13	2:B:453:VAL:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/482 (98%)	463 (98%)	7 (2%)	0	100	100
2	B	455/455 (100%)	446 (98%)	9 (2%)	0	100	100
3	C	44/89 (49%)	43 (98%)	1 (2%)	0	100	100
All	All	969/1026 (94%)	952 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/441 (98%)	430 (99%)	3 (1%)	84	92
2	B	421/419 (100%)	420 (100%)	1 (0%)	93	97
3	C	47/81 (58%)	47 (100%)	0	100	100
All	All	901/941 (96%)	897 (100%)	4 (0%)	91	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	374	ILE
1	A	404	ARG
1	A	445	PHE
2	B	402	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	469/482 (97%)	0.21	31 (6%)	18 23	24, 40, 91, 159	0
2	B	455/455 (100%)	0.08	20 (4%)	34 41	29, 46, 84, 140	0
3	C	48/89 (53%)	2.28	24 (50%)	0 0	36, 94, 152, 185	0
All	All	972/1026 (94%)	0.25	75 (7%)	13 17	24, 45, 100, 185	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	ALA	13.5
1	A	79	PRO	9.8
1	A	244	ALA	9.0
1	A	187	ALA	8.6
1	A	245	ALA	7.9
3	C	69	VAL	7.8
3	C	29	LEU	7.8
1	A	186	SER	7.7
1	A	80	ASP	7.6
3	C	67	VAL	7.1
3	C	28	SER	6.8
1	A	191	TYR	6.7
1	A	189	ASP	6.5
3	C	68	GLU	6.4
2	B	384	GLY	6.4
1	A	185	ASP	6.3
2	B	454	LEU	6.3
3	C	89	GLN	6.0
1	A	125	GLN	5.5
1	A	112	SER	5.5
1	A	82	LYS	5.4
2	B	383	PRO	5.2
1	A	247	ALA	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	57	THR	5.0
2	B	381	GLU	4.8
2	B	452	HIS	4.7
1	A	81	LYS	4.5
3	C	63	ASN	4.4
3	C	39	PRO	4.3
1	A	184	ALA	4.3
3	C	70	ASP	4.2
3	C	66	ASP	4.1
1	A	188	LYS	4.1
2	B	428	LYS	4.0
2	B	451	SER	3.8
2	B	385	ALA	3.7
1	A	227	ASN	3.7
1	A	190	LEU	3.6
2	B	256	THR	3.6
2	B	386	GLN	3.6
2	B	431	MET	3.6
3	C	62	GLU	3.5
3	C	58	ASN	3.5
2	B	253	LEU	3.4
1	A	328	GLU	3.3
3	C	27	LYS	3.2
3	C	41	ASP	3.1
1	A	192	ASP	3.0
2	B	450	HIS	3.0
3	C	71	ASP	2.9
2	B	359	GLY	2.8
2	B	453	VAL	2.8
1	A	329	PHE	2.6
1	A	90	ALA	2.6
1	A	114	LYS	2.6
1	A	101	SER	2.5
1	A	243	PRO	2.5
1	A	468	LYS	2.4
3	C	83	ARG	2.4
3	C	88	ASN	2.4
2	B	1	MET	2.3
3	C	86	ARG	2.3
2	B	380	PHE	2.3
3	C	64	TRP	2.3
1	A	226	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	ASN	2.2
3	C	87	GLU	2.2
2	B	255[A]	LEU	2.2
2	B	254	PRO	2.1
1	A	193	ALA	2.1
3	C	59	ILE	2.0
3	C	40	ILE	2.0
3	C	30	GLU	2.0
1	A	132	ARG	2.0
2	B	430	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.