Full wwPDB X-ray Structure Validation Report (i)

Jun 26, 2024 – 05:04 AM EDT

PDB ID	:	6UB2
Title	:	Crystal structure of a GH128 (subgroup IV) endo-beta-1,3-glucanase from Lentinula edodes (LeGH128_IV)
Authors	:	Santos, C.R.; Lima, E.A.; Mandelli, F.; Vieira, P.S.; Murakami, M.T.
Deposited on	:	2019-09-11
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	86% 6% 8%
1	B	270	88% • 8%
1	C	270	89% • 8%
1	D	270	87% • 9%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-beta-1,3-glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	248	1874	1205	293	369	7	0	3	0
1	B	249	1877	1208	294	368	7	0	1	0
1	C	249	1873	1205	294	367	7	0	0	0
1	D	247	1869	1203	292	367	7	0	2	0

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP G9M5R4
A	-1	GLY	-	expression tag	UNP G9M5R4
A	0	SER	-	expression tag	UNP G9M5R4
A	1	SER	-	expression tag	UNP G9M5R4
A	2	HIS	-	expression tag	UNP G9M5R4
A	3	HIS	-	expression tag	UNP G9M5R4
A	4	HIS	-	expression tag	UNP G9M5R4
A	5	HIS	-	expression tag	UNP G9M5R4
A	6	HIS	-	expression tag	UNP G9M5R4
A	7	HIS	-	expression tag	UNP G9M5R4
A	8	SER	-	expression tag	UNP G9M5R4
A	9	SER	-	expression tag	UNP G9M5R4
A	10	GLY	-	expression tag	UNP G9M5R4
A	11	LEU	-	expression tag	UNP G9M5R4
A	12	VAL	-	expression tag	UNP G9M5R4
A	13	PRO	-	expression tag	UNP G9M5R4
A	14	ARG	-	expression tag	UNP G9M5R4
A	15	GLY	-	expression tag	UNP G9M5R4
A	16	SER	-	expression tag	UNP G9M5R4
A	17	HIS	-	expression tag	UNP G9M5R4
A	18	MET	-	expression tag	UNP G9M5R4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	ALA	-	expression tag	UNP G9M5R4
A	20	SER	-	expression tag	UNP G9M5R4
B	-2	MET	-	initiating methionine	UNP G9M5R4
B	-1	GLY	-	expression tag	UNP G9M5R4
B	0	SER	-	expression tag	UNP G9M5R4
B	1	SER	-	expression tag	UNP G9M5R4
B	2	HIS	-	expression tag	UNP G9M5R4
B	3	HIS	-	expression tag	UNP G9M5R4
B	4	HIS	-	expression tag	UNP G9M5R4
B	5	HIS	-	expression tag	UNP G9M5R4
B	6	HIS	-	expression tag	UNP G9M5R4
B	7	HIS	-	expression tag	UNP G9M5R4
B	8	SER	-	expression tag	UNP G9M5R4
B	9	SER	-	expression tag	UNP G9M5R4
B	10	GLY	-	expression tag	UNP G9M5R4
B	11	LEU	-	expression tag	UNP G9M5R4
B	12	VAL	-	expression tag	UNP G9M5R4
B	13	PRO	-	expression tag	UNP G9M5R4
B	14	ARG	-	expression tag	UNP G9M5R4
B	15	GLY	-	expression tag	UNP G9M5R4
B	16	SER	-	expression tag	UNP G9M5R4
B	17	HIS	-	expression tag	UNP G9M5R4
B	18	MET	-	expression tag	UNP G9M5R4
B	19	ALA	-	expression tag	UNP G9M5R4
B	20	SER	-	expression tag	UNP G9M5R4
C	-2	MET	-	initiating methionine	UNP G9M5R4
C	-1	GLY	-	expression tag	UNP G9M5R4
C	0	SER	-	expression tag	UNP G9M5R4
C	1	SER	-	expression tag	UNP G9M5R4
C	2	HIS	-	expression tag	UNP G9M5R4
C	3	HIS	-	expression tag	UNP G9M5R4
C	4	HIS	-	expression tag	UNP G9M5R4
C	5	HIS	-	expression tag	UNP G9M5R4
C	6	HIS	-	expression tag	UNP G9M5R4
C	7	HIS	-	expression tag	UNP G9M5R4
C	8	SER	-	expression tag	UNP G9M5R4
C	9	SER	-	expression tag	UNP G9M5R4
C	10	GLY	-	expression tag	UNP G9M5R4
C	11	LEU	-	expression tag	UNP G9M5R4
C	12	VAL	-	expression tag	UNP G9M5R4
C	13	PRO	-	expression tag	UNP G9M5R4
C	14	ARG	-	expression tag	UNP G9M5R4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	15	GLY	-	expression tag	UNP G9M5R4
C	16	SER	-	expression tag	UNP G9M5R4
C	17	HIS	-	expression tag	UNP G9M5R4
C	18	MET	-	expression tag	UNP G9M5R4
C	19	ALA	-	expression tag	UNP G9M5R4
C	20	SER	-	expression tag	UNP G9M5R4
D	-2	MET	-	initiating methionine	UNP G9M5R4
D	-1	GLY	-	expression tag	UNP G9M5R4
D	0	SER	-	expression tag	UNP G9M5R4
D	1	SER	-	expression tag	UNP G9M5R4
D	2	HIS	-	expression tag	UNP G9M5R4
D	3	HIS	-	expression tag	UNP G9M5R4
D	4	HIS	-	expression tag	UNP G9M5R4
D	5	HIS	-	expression tag	UNP G9M5R4
D	6	HIS	-	expression tag	UNP G9M5R4
D	7	HIS	-	expression tag	UNP G9M5R4
D	8	SER	-	expression tag	UNP G9M5R4
D	9	SER	-	expression tag	UNP G9M5R4
D	10	GLY	-	expression tag	UNP G9M5R4
D	11	LEU	-	expression tag	UNP G9M5R4
D	12	VAL	-	expression tag	UNP G9M5R4
D	13	PRO	-	expression tag	UNP G9M5R4
D	14	ARG	-	expression tag	UNP G9M5R4
D	15	GLY	-	expression tag	UNP G9M5R4
D	16	SER	-	expression tag	UNP G9M5R4
D	17	HIS	-	expression tag	UNP G9M5R4
D	18	MET	-	expression tag	UNP G9M5R4
D	19	ALA	-	expression tag	UNP G9M5R4
D	20	SER	-	expression tag	UNP G9M5R4

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	12	6	1	4	1	0	0
2	B	1	12	6	1	4	1	0	0
2	C	1	12	6	1	4	1	0	0
2	D	1	12	6	1	4	1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0
3	D	1	1	1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	318	Total O 318 318	0	0
5	B	308	Total O 308 308	0	0
5	C	284	Total O 284 284	0	0
5	D	361	Total O 361 361	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-beta-1,3-glucanase

Chain A:



- Molecule 1: Endo-beta-1,3-glucanase

Chain B:



- Molecule 1: Endo-beta-1,3-glucanase

Chain C:



- Molecule 1: Endo-beta-1,3-glucanase

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	72.82Å 75.16Å 99.45Å 90.00° 103.60° 90.00°	Depositor
Resolution (Å)	49.09 – 1.80 49.09 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.2 (49.09-1.80) 98.2 (49.09-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.35 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.172 , 0.212 0.182 , 0.218	Depositor DCC
R_{free} test set	4748 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	21.2	Xtriage
Anisotropy	0.640	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 45.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	8831	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.57% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.89	2/1945 (0.1%)	0.90	0/2668
1	B	0.85	1/1939 (0.1%)	0.88	0/2660
1	C	0.85	0/1932	0.91	0/2650
1	D	0.89	0/1935	0.92	1/2655 (0.0%)
All	All	0.87	3/7751 (0.0%)	0.91	1/10633 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	35	ASP	CG-OD2	5.39	1.37	1.25
1	A	223	LEU	C-O	5.33	1.33	1.23
1	A	174	GLU	CD-OE2	5.03	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	23	ARG	NE-CZ-NH2	-5.39	117.60	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1749	8	0
1	B	1877	0	1758	6	0
1	C	1873	0	1751	4	0
1	D	1869	0	1746	8	0
2	A	12	0	13	1	0
2	B	12	0	13	0	0
2	C	12	0	13	0	0
2	D	12	0	13	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	5	0	0	0	0
4	D	10	0	0	0	0
5	A	318	0	0	1	0
5	B	308	0	0	0	0
5	C	284	0	0	0	0
5	D	361	0	0	2	0
All	All	8831	0	7056	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:205:VAL:O	1:D:209:GLU:HG3	2.05	0.57
1:D:167:TRP:CZ2	1:D:169:GLY:HA3	2.42	0.55
1:D:167:TRP:O	1:D:196:PHE:HA	2.07	0.54
1:A:167:TRP:O	1:A:196:PHE:HA	2.09	0.52
1:B:167:TRP:O	1:B:196:PHE:HA	2.11	0.50
1:C:167:TRP:O	1:C:196:PHE:HA	2.12	0.50
1:D:85:GLN:OE1	5:D:601:HOH:O	2.20	0.50
1:B:32:SER:N	1:B:33:PRO:CD	2.75	0.49
1:B:236:GLN:HA	1:B:243:VAL:HG21	1.96	0.48
1:D:32:SER:N	1:D:33:PRO:CD	2.78	0.47
1:A:229:ALA:HB2	1:A:252:LEU:HD12	1.99	0.45
1:C:29:TRP:O	1:C:58:PRO:HG2	2.17	0.45
1:B:32:SER:N	1:B:33:PRO:HD3	2.31	0.45
1:B:19:ALA:HA	1:B:45:GLU:OE2	2.17	0.44
1:A:92:GLN:HG3	5:A:804:HOH:O	2.18	0.44
1:D:32:SER:N	1:D:33:PRO:HD3	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:GLN:O	1:B:145:SER:HB3	2.18	0.43
1:C:145:SER:O	1:C:149:GLU:HG3	2.19	0.43
1:A:103:GLU:N	1:A:104:PRO:CD	2.82	0.43
1:D:103:GLU:N	1:D:104:PRO:CD	2.82	0.42
1:A:252:LEU:CD2	1:A:262:VAL:HG12	2.50	0.42
1:D:74:GLY:HA3	5:D:743:HOH:O	2.19	0.42
1:C:119:ILE:O	1:C:123:ASN:HB2	2.20	0.42
1:A:183:HIS:HB2	1:A:220:TYR:CE1	2.56	0.41
1:A:230:THR:O	1:A:251:CYS:HB3	2.21	0.40
1:A:52:TRP:CD2	2:A:501:MES:H31	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/270 (92%)	245 (98%)	4 (2%)	0	100	100
1	B	248/270 (92%)	244 (98%)	4 (2%)	0	100	100
1	C	247/270 (92%)	244 (99%)	3 (1%)	0	100	100
1	D	247/270 (92%)	240 (97%)	7 (3%)	0	100	100
All	All	991/1080 (92%)	973 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/212 (93%)	197 (100%)	0	100	100
1	B	195/212 (92%)	195 (100%)	0	100	100
1	C	194/212 (92%)	193 (100%)	1 (0%)	88	87
1	D	195/212 (92%)	195 (100%)	0	100	100
All	All	781/848 (92%)	780 (100%)	1 (0%)	93	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	188	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	264	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	SO4	D	502	-	4,4,4	0.54	0	6,6,6	0.30	0
2	MES	B	501	-	12,12,12	0.86	0	14,16,16	1.33	2 (14%)
2	MES	D	501	-	12,12,12	0.78	0	14,16,16	0.49	0
2	MES	A	501	-	12,12,12	0.85	0	14,16,16	0.86	1 (7%)
2	MES	C	501	-	12,12,12	0.81	0	14,16,16	0.76	0
4	SO4	D	503	-	4,4,4	0.29	0	6,6,6	0.45	0
4	SO4	B	502	-	4,4,4	0.29	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	501	-	-	5/6/14/14	0/1/1/1
2	MES	B	501	-	-	3/6/14/14	0/1/1/1
2	MES	C	501	-	-	2/6/14/14	0/1/1/1
2	MES	D	501	-	-	0/6/14/14	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	501	MES	O1S-S-C8	-3.26	103.00	106.92
2	B	501	MES	O3S-S-C8	3.07	110.73	105.77
2	A	501	MES	O1S-S-C8	-2.53	103.87	106.92

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	MES	C8-C7-N4-C3
2	A	501	MES	C8-C7-N4-C5
2	A	501	MES	C7-C8-S-O1S
2	A	501	MES	C7-C8-S-O3S
2	B	501	MES	C7-C8-S-O1S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	501	MES	C7-C8-S-O3S
2	C	501	MES	C8-C7-N4-C3
2	C	501	MES	C8-C7-N4-C5
2	A	501	MES	C7-C8-S-O2S
2	B	501	MES	C7-C8-S-O2S

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	MES	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	248/270 (91%)	-0.63	1 (0%) 92 90	17, 22, 30, 50	0
1	B	249/270 (92%)	-0.58	0 100 100	18, 24, 36, 46	0
1	C	249/270 (92%)	-0.62	0 100 100	17, 24, 35, 45	0
1	D	247/270 (91%)	-0.70	0 100 100	15, 20, 26, 31	0
All	All	993/1080 (91%)	-0.63	1 (0%) 95 93	15, 22, 33, 50	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	20	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MES	C	501	12/12	0.82	0.15	57,62,65,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MES	A	501	12/12	0.86	0.16	57,64,70,73	0
4	SO4	D	503	5/5	0.87	0.21	36,46,56,57	0
4	SO4	B	502	5/5	0.91	0.17	64,66,72,73	0
2	MES	B	501	12/12	0.91	0.12	43,45,58,58	0
4	SO4	D	502	5/5	0.92	0.13	42,45,49,53	0
2	MES	D	501	12/12	0.95	0.09	28,32,43,43	0
3	CL	C	502	1/1	0.98	0.05	32,32,32,32	0
3	CL	B	503	1/1	0.98	0.06	34,34,34,34	0
3	CL	D	504	1/1	0.99	0.03	31,31,31,31	0
3	CL	A	502	1/1	1.00	0.03	24,24,24,24	0

6.5 Other polymers (i)

There are no such residues in this entry.