



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:35 PM UTC

PDB ID : 9UHI / pdb_00009uhi
Title : FGFR1 kinase domain with a covalent inhibitor 9o
Authors : Chen, X.J.; Chen, Y.H.
Deposited on : 2025-04-14
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	 2% 92% 7%
1	B	310	 9% 86% 7% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fibroblast growth factor receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	309	Total	C	N	O	S	0	0	0
			2421	1537	413	453	18			
1	B	288	Total	C	N	O	S	0	0	0
			2239	1425	381	415	18			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	456	GLY	-	expression tag	UNP P11362
A	457	PRO	-	expression tag	UNP P11362
A	584	SER	CYS	conflict	UNP P11362
B	456	GLY	-	expression tag	UNP P11362
B	457	PRO	-	expression tag	UNP P11362
B	584	SER	CYS	conflict	UNP P11362

- Molecule 2 is {N}-[1-methyl-3-[3-[1-(2-morpholin-4-ylethyl)pyrazol-4-yl]quinoxalin-5-yl]indol-6-yl]propanamide (CCD ID: A1EPF) (formula: C₂₉H₃₁N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	38	29	7	2	0	0
2	B	1	38	29	7	2	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0
3	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	334	Total	O	0	0
			334	334		
4	B	222	Total	O	0	0
			222	222		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fibroblast growth factor receptor 1

Chain A: 



- Molecule 1: Fibroblast growth factor receptor 1

Chain B: 



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	211.43Å 49.28Å 66.55Å 90.00° 107.42° 90.00°	Depositor
Resolution (Å)	34.75 – 1.76 34.75 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.1 (34.75-1.76) 99.1 (34.75-1.76)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.18 (at 1.76Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.161 , 0.197 0.161 , 0.196	Depositor DCC
R_{free} test set	3248 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	19.2	Xtrriage
Anisotropy	0.235	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5317	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, A1EPF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2477	0.72	0/3362
1	B	0.49	0/2286	0.66	0/3098
All	All	0.54	0/4763	0.69	0/6460

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2421	0	2371	20	0
1	B	2239	0	2209	17	0
2	A	38	0	0	0	0
2	B	38	0	0	0	0
3	A	20	0	0	0	0
3	B	5	0	0	0	0
4	A	334	0	0	2	3
4	B	222	0	0	3	3
All	All	5317	0	4580	36	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:486:GLU:OE1	1:A:491:GLN:NE2	2.00	0.95
1:A:711:LYS:O	1:A:715:GLU:HG3	1.86	0.74
1:A:540:LYS:HD3	1:A:546:ASN:OD1	1.90	0.72
1:A:540:LYS:HZ1	1:A:547:LEU:H	1.39	0.70
1:B:673:PHE:HE1	1:B:710:PHE:HD1	1.45	0.65
1:B:485:GLY:O	4:B:901:HOH:O	2.15	0.60
1:A:496:GLU:HB3	1:A:508:VAL:CG1	2.32	0.60
1:B:707:GLU:HA	1:B:710:PHE:HD2	1.68	0.59
1:B:721:LYS:NZ	1:B:727:ASN:OD1	2.35	0.56
1:B:748:LYS:NZ	1:B:752:GLU:OE2	2.37	0.56
1:A:752:GLU:OE2	4:A:901:HOH:O	2.18	0.56
1:A:579:PRO:HG2	1:B:674:ASP:O	2.08	0.53
1:B:673:PHE:CE1	1:B:710:PHE:HD1	2.27	0.52
1:A:664:VAL:HG23	1:A:672:LEU:HD21	1.90	0.51
1:B:535:MET:HE3	4:B:1056:HOH:O	2.10	0.51
1:B:490:GLY:N	4:B:902:HOH:O	2.18	0.47
1:A:475:ARG:HD3	4:A:1003:HOH:O	2.15	0.47
1:B:707:GLU:HA	1:B:710:PHE:CD2	2.47	0.47
1:A:662:LEU:HD13	1:A:706:VAL:HG21	1.98	0.45
1:A:537:MET:HE3	1:A:537:MET:HB3	1.83	0.45
1:A:654:TYR:HE1	1:A:706:VAL:HG12	1.81	0.44
1:B:477:ARG:HD2	1:B:498:ILE:HG23	1.99	0.44
1:A:664:VAL:CG2	1:A:672:LEU:HD21	2.48	0.43
1:A:496:GLU:HB3	1:A:508:VAL:HG13	1.99	0.43
1:B:664:VAL:HA	1:B:667:MET:HE3	2.00	0.43
1:B:601:VAL:HG11	1:B:757:ILE:HG22	2.00	0.43
1:A:611:MET:HE3	1:A:611:MET:HB3	1.86	0.43
1:A:664:VAL:HG23	1:A:672:LEU:CD2	2.48	0.42
1:A:540:LYS:HD3	1:A:540:LYS:HA	1.65	0.42
1:B:705:PRO:HG2	1:B:708:GLU:HG2	2.02	0.42
1:A:540:LYS:HD2	1:A:546:ASN:HB3	2.02	0.42
1:A:684:TRP:CE3	1:A:737:TRP:HA	2.56	0.41
1:A:654:TYR:CE1	1:A:706:VAL:HG12	2.55	0.41
1:B:500:LEU:HD11	1:B:511:VAL:HG11	2.03	0.41
1:B:707:GLU:HG2	1:B:708:GLU:N	2.36	0.40
1:B:484:LEU:HD23	1:B:484:LEU:HA	1.86	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1059:HOH:O	4:B:990:HOH:O[2_546]	2.12	0.08
4:A:929:HOH:O	4:B:1058:HOH:O[2_546]	2.13	0.07
4:A:1166:HOH:O	4:B:998:HOH:O[1_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/310 (99%)	305 (99%)	2 (1%)	0	100	100
1	B	282/310 (91%)	277 (98%)	5 (2%)	0	100	100
All	All	589/620 (95%)	582 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/271 (96%)	258 (99%)	3 (1%)	65	52
1	B	239/271 (88%)	239 (100%)	0	100	100
All	All	500/542 (92%)	497 (99%)	3 (1%)	78	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	488	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	706	VAL
1	A	738	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	650	HIS
1	A	659	ASN
1	A	717	HIS
1	A	738	HIS
1	A	743	GLN
1	B	717	HIS
1	B	738	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	803	-	4,4,4	0.38	0	6,6,6	0.40	0
3	SO4	A	804	-	4,4,4	0.28	0	6,6,6	0.84	0
3	SO4	B	802	-	4,4,4	0.24	0	6,6,6	0.18	0
2	A1EPF	B	801	1	43,43,43	2.33	11 (25%)	58,61,61	4.05	15 (25%)
3	SO4	A	805	-	4,4,4	0.37	0	6,6,6	0.42	0
2	A1EPF	A	801	1	43,43,43	2.18	11 (25%)	58,61,61	3.59	15 (25%)
3	SO4	A	802	-	4,4,4	0.23	0	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EPF	B	801	1	-	3/19/27/27	0/6/6/6
2	A1EPF	A	801	1	-	2/19/27/27	0/6/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	A1EPF	C11-N5	-6.55	1.32	1.47
2	A	801	A1EPF	C11-N5	-6.19	1.33	1.47
2	B	801	A1EPF	C21-C20	5.45	1.48	1.41
2	B	801	A1EPF	C15-N5	-5.29	1.32	1.46
2	A	801	A1EPF	C15-N5	-5.28	1.32	1.46
2	A	801	A1EPF	C21-C20	5.16	1.47	1.41
2	B	801	A1EPF	C12-N5	-5.15	1.33	1.46
2	A	801	A1EPF	C12-N5	-4.97	1.33	1.46
2	B	801	A1EPF	C27-N7	4.59	1.45	1.35
2	B	801	A1EPF	C19-N6	-4.58	1.30	1.37
2	A	801	A1EPF	C27-N7	4.12	1.44	1.35
2	B	801	A1EPF	C8-N3	3.22	1.37	1.32
2	A	801	A1EPF	C19-N6	-3.21	1.32	1.37
2	A	801	A1EPF	C4-N2	-3.19	1.32	1.37
2	B	801	A1EPF	C6-C3	-2.95	1.39	1.43
2	B	801	A1EPF	C9-N4	2.90	1.39	1.34
2	A	801	A1EPF	C8-N3	2.78	1.37	1.32
2	A	801	A1EPF	C6-C3	-2.67	1.39	1.43
2	A	801	A1EPF	C9-N4	2.63	1.39	1.34
2	B	801	A1EPF	C4-N2	-2.62	1.33	1.37
2	A	801	A1EPF	C6-C7	2.35	1.53	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	A1EPF	O2-C27	-2.01	1.19	1.23

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	A1EPF	C20-N6-C19	16.87	120.78	108.54
2	A	801	A1EPF	C20-N6-C19	15.86	120.05	108.54
2	B	801	A1EPF	C21-C7-C19	12.81	112.69	106.02
2	B	801	A1EPF	C7-C19-N6	-11.62	102.51	110.95
2	A	801	A1EPF	C7-C19-N6	-10.45	103.36	110.95
2	A	801	A1EPF	C21-C20-N6	-9.59	102.06	107.91
2	A	801	A1EPF	C21-C7-C19	9.44	110.93	106.02
2	B	801	A1EPF	C20-C21-C7	-9.12	101.61	106.83
2	B	801	A1EPF	C21-C20-N6	-9.00	102.42	107.91
2	A	801	A1EPF	C26-N6-C20	-7.85	118.41	125.69
2	B	801	A1EPF	C26-N6-C19	-7.08	118.24	125.70
2	A	801	A1EPF	C20-C21-C7	-5.71	103.56	106.83
2	B	801	A1EPF	C26-N6-C20	-5.08	120.98	125.69
2	B	801	A1EPF	C13-C12-N5	4.79	117.40	110.12
2	A	801	A1EPF	C26-N6-C19	-4.03	121.46	125.70
2	B	801	A1EPF	C22-C20-N6	3.92	137.81	131.24
2	B	801	A1EPF	C21-C7-C6	-3.65	120.03	126.62
2	A	801	A1EPF	C24-C23-C22	3.59	124.00	119.66
2	B	801	A1EPF	C10-N4-C9	-3.15	122.95	128.83
2	A	801	A1EPF	C24-C25-C21	-2.95	116.00	120.86
2	A	801	A1EPF	C22-C20-N6	2.93	136.15	131.24
2	B	801	A1EPF	C10-N4-N3	2.65	127.67	121.16
2	B	801	A1EPF	C12-N5-C15	2.63	114.51	108.84
2	A	801	A1EPF	C21-C7-C6	-2.60	121.93	126.62
2	B	801	A1EPF	C8-N3-N4	2.40	108.35	104.68
2	A	801	A1EPF	C23-C22-C20	-2.24	115.68	119.48
2	B	801	A1EPF	C25-C21-C7	2.22	137.30	132.34
2	A	801	A1EPF	C17-C18-C6	2.21	123.52	119.54
2	A	801	A1EPF	C16-C17-C18	-2.19	117.63	121.00
2	A	801	A1EPF	C24-C23-N7	-2.03	113.59	120.41

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	A1EPF	N4-C10-C11-N5
2	B	801	A1EPF	C10-C11-N5-C15

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	801	A1EPF	C11-C10-N4-N3
2	A	801	A1EPF	C10-C11-N5-C12
2	A	801	A1EPF	C3-C6-C7-C21

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/310 (99%)	-0.23	6 (1%) 66 73	10, 19, 36, 66	0
1	B	288/310 (92%)	0.31	28 (9%) 13 15	12, 24, 67, 94	0
All	All	597/620 (96%)	0.03	34 (5%) 29 33	10, 21, 50, 94	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	654	TYR	6.8
1	B	653	TYR	5.4
1	B	488	CYS	5.0
1	B	660	GLY	5.0
1	B	579	PRO	4.8
1	B	489	PHE	4.3
1	B	578	PRO	4.0
1	B	673	PHE	3.8
1	B	672	LEU	3.6
1	B	658	THR	3.5
1	B	710	PHE	3.3
1	B	677	TYR	3.3
1	B	580	GLY	3.2
1	B	657	THR	3.0
1	B	661	ARG	2.9
1	B	676	ILE	2.8
1	B	762	SER	2.8
1	B	662	LEU	2.7
1	A	503	ASP	2.7
1	B	674	ASP	2.7
1	A	456	GLY	2.6
1	B	594	GLN	2.6
1	B	644	LEU	2.6
1	A	458	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	659	ASN	2.5
1	A	651	ILE	2.3
1	B	675	ARG	2.3
1	A	706	VAL	2.2
1	A	504	LYS	2.2
1	B	671	ALA	2.1
1	B	655	LYS	2.1
1	B	763	ASN	2.1
1	B	740	VAL	2.0
1	B	486	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	803	5/5	0.79	0.13	45,47,48,83	0
3	SO4	B	802	5/5	0.87	0.10	46,60,68,77	0
3	SO4	A	805	5/5	0.93	0.09	36,37,57,62	0
2	A1EPF	B	801	38/38	0.95	0.09	16,32,75,78	0
2	A1EPF	A	801	38/38	0.96	0.07	13,17,41,42	0
3	SO4	A	802	5/5	0.99	0.05	21,22,30,30	0
3	SO4	A	804	5/5	0.99	0.05	20,23,25,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1EPF B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1EPF A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.