



Full wwPDB EM Validation Report ⓘ

May 31, 2026 – 12:20 AM JST

PDB ID : 9V7Z / pdb_00009v7z
EMDB ID : EMD-64826
Title : Structure of the Argonaute protein from Verrucomicrobia bacterium in complex with guide DNA and target RNA in target-released state
Authors : Wu, S.
Deposited on : 2025-05-28
Resolution : 2.97 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : **NOT EXECUTED**
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	229148	23984
Ramachandran outliers	224038	23583
Sidechain outliers	223484	23102
RNA backbone	8273	3508

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	782	 83% 14% .
2	B	18	 56% 22% 22%
3	C	19	 42% 5% 53%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VbAgo.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	757	Total	C	N	O	S	0	0
			6044	3811	1108	1110	15		

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*GP*AP*GP*GP*TP*AP*GP*TP*AP*GP*GP*TP*TP*GP*TP*AP*T)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	14	Total	C	N	O	P	0	0
			296	140	55	87	14		

- Molecule 3 is a RNA chain called RNA (5'-R(*UP*AP*UP*AP*CP*AP*AP*CP*CP*UP*AP*CP*UP*AP*CP*CP*UP*CP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	9	Total	C	N	O	P	0	0
			184	83	28	64	9		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
4	A	2	Total	Mg	0
			2	2	
4	B	1	Total	Mg	0
			1	1	


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		AltConf
5	A	12	Total	O	0
			12	12	
5	B	2	Total	O	0
			2	2	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: VbAgo



• Molecule 2: DNA (5'-D(P*TP*GP*AP*GP*GP*TP*AP*GP*TP*AP*GP*GP*TP*TP*GP*T P*AP*T)-3')



• Molecule 3: RNA (5'-R(*UP*AP*UP*AP*CP*AP*AP*CP*CP*UP*AP*CP*UP*AP*CP*CP*UP*CP*A)-3')



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	112179	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	54	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOCONTINUUM (6k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/6176	0.32	0/8382
2	B	0.20	0/332	0.35	0/511
3	C	0.12	0/203	0.17	0/312
All	All	0.15	0/6711	0.32	0/9205

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6044	0	6034	61	0
2	B	296	0	160	3	0
3	C	184	0	97	0	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	12	0	0	0	0
5	B	2	0	0	0	0
All	All	6541	0	6291	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:704:ARG:HH12	2:B:5:DG:H21	1.30	0.77
1:A:385:MET:HE2	1:A:520:ARG:HD2	1.66	0.75
1:A:236:THR:HG23	1:A:238:LYS:H	1.55	0.72
1:A:225:LYS:HG3	1:A:226:ILE:HG13	1.78	0.65
1:A:375:GLU:OE2	1:A:379:ARG:NH2	2.30	0.64
1:A:90:ARG:NH1	1:A:91:GLU:O	2.33	0.61
1:A:351:VAL:HG11	1:A:676:MET:HE2	1.84	0.60
1:A:404:LEU:HD12	1:A:439:GLN:HG2	1.83	0.60
1:A:164:LEU:HD11	1:A:179:ALA:HB1	1.85	0.59
1:A:300:ASP:OD1	1:A:303:ARG:NH1	2.36	0.58
1:A:57:VAL:HG22	1:A:70:THR:HG22	1.84	0.58
1:A:390:GLU:OE1	1:A:391:HIS:NE2	2.38	0.56
2:B:4:DG:H2'	2:B:5:DG:C8	2.40	0.56
1:A:26:LEU:HD12	1:A:68:PHE:HB2	1.87	0.55
1:A:220:LEU:HD21	1:A:227:LYS:HD3	1.89	0.55
1:A:157:ARG:HE	1:A:301:GLU:HG3	1.71	0.55
1:A:530:ALA:HA	1:A:759:LEU:HD13	1.88	0.54
1:A:654:LYS:HA	1:A:709:LEU:HD23	1.89	0.54
1:A:60:THR:HB	1:A:69:LEU:HD13	1.90	0.52
1:A:588:ARG:HG3	1:A:627:ALA:HB2	1.92	0.52
1:A:157:ARG:NE	1:A:301:GLU:HG3	2.24	0.52
1:A:607:VAL:HG23	1:A:609:LEU:HG	1.91	0.52
1:A:150:ASN:ND2	1:A:158:VAL:O	2.43	0.51
1:A:177:LEU:HD22	1:A:339:ILE:HD13	1.92	0.51
1:A:80:ARG:HE	1:A:82:LEU:HD21	1.76	0.50
1:A:408:ASP:OD1	1:A:409:ASN:N	2.44	0.50
1:A:421:ARG:HB3	1:A:519:LEU:HD21	1.95	0.49
1:A:674:ASN:ND2	1:A:700:ASP:OD1	2.46	0.48
1:A:363:GLN:HG3	1:A:391:HIS:ND1	2.29	0.48
1:A:561:VAL:HG22	1:A:569:VAL:HG12	1.95	0.48
1:A:30:GLU:OE1	1:A:90:ARG:NH2	2.46	0.47
1:A:208:SER:O	1:A:216:ARG:NH1	2.44	0.47
1:A:373:ILE:HG22	1:A:676:MET:HE1	1.97	0.47
1:A:35:ASP:OD1	1:A:38:THR:OG1	2.26	0.47
1:A:578:ARG:HD3	1:A:578:ARG:H	1.80	0.47
1:A:276:ARG:HH12	1:A:282:ASP:HA	1.81	0.46
1:A:41:ARG:HG2	1:A:44:ARG:NH2	2.32	0.45
1:A:185:TYR:OH	1:A:307:VAL:O	2.31	0.45
1:A:578:ARG:O	1:A:580:ARG:HG2	2.18	0.44
1:A:329:ASP:HA	1:A:336:THR:HG23	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:PRO:HD2	1:A:20:ALA:HB2	1.99	0.44
1:A:49:LEU:HD11	1:A:94:ILE:HD13	1.99	0.44
1:A:203:LEU:HB3	1:A:206:HIS:HB2	1.99	0.43
1:A:405:LEU:HD23	1:A:473:ILE:HG23	2.00	0.43
1:A:623:ASN:OD1	1:A:624:GLU:N	2.52	0.43
1:A:16:LEU:HD12	1:A:17:PRO:HD2	2.00	0.43
1:A:351:VAL:HG11	1:A:676:MET:CE	2.48	0.43
1:A:139:HIS:CG	1:A:140:TRP:H	2.37	0.43
2:B:10:DA:H2''	2:B:11:DG:C8	2.53	0.43
1:A:109:SER:OG	1:A:110:SER:N	2.51	0.42
1:A:576:GLN:HG2	1:A:577:LEU:H	1.84	0.42
1:A:58:VAL:HG12	1:A:115:LEU:HD11	2.02	0.42
1:A:657:LEU:HA	1:A:658:PRO:HD3	1.85	0.42
1:A:86:ASP:CG	1:A:88:PRO:HD2	2.44	0.42
1:A:449:HIS:HB2	1:A:480:ARG:HG2	2.02	0.41
1:A:148:PRO:HD2	1:A:151:LEU:HD13	2.03	0.41
1:A:48:GLN:O	1:A:52:GLU:OE1	2.38	0.41
1:A:211:ILE:O	1:A:216:ARG:NH2	2.53	0.41
1:A:267:PRO:HG2	1:A:270:ASP:OD2	2.21	0.41
1:A:455:LEU:HB3	1:A:490:ARG:NH1	2.36	0.41
1:A:657:LEU:HG	1:A:659:VAL:HG23	2.02	0.41
1:A:488:LYS:HA	1:A:488:LYS:HD3	1.94	0.40
1:A:46:ALA:CB	1:A:59:PRO:HD3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	755/782 (96%)	737 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	647/671 (96%)	647 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	ASN
1	A	269	ASN
1	A	304	ASN
1	A	325	ASN
1	A	363	GLN
1	A	464	ASN
1	A	646	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	8/19 (42%)	1 (12%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	15	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.