



wwPDB X-ray Structure Validation Summary Report

Aug 8, 2020 – 04:51 AM BST

PDB ID : 6VM6
Title : Structure of Acinetobacter baumannii Cap4 SAVED/CARF-domain containing receptor with the cyclic trinucleotide 2'3'3'-cAAA
Authors : Lowey, B.; Whiteley, A.T.; Keszei, A.F.A.; Morehouse, B.R.; Antine, S.P.; Cabrera, V.; Schwede, F.; Mekalanos, J.J.; Shao, S.; Lee, A.S.Y.; Kranzusch, P.J.
Deposited on : 2020-01-27
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)
RNA backbone	3102	1000 (2.54-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	
1	B	462	
1	C	462	
1	D	462	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	462	
1	F	462	
2	G	3	
2	H	3	
2	I	3	
2	J	3	
2	K	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	501	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 23004 atoms, of which 66 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SAVED domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	438	3518	2245	593	669	11	0	0	0
1	B	440	3528	2250	595	672	11	0	0	0
1	C	439	3522	2247	594	670	11	0	0	0
1	D	440	3528	2250	595	672	11	0	0	0
1	E	432	3472	2217	586	658	11	0	0	0
1	F	437	3510	2239	592	668	11	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP C0VHC9
B	1	SER	-	expression tag	UNP C0VHC9
C	1	SER	-	expression tag	UNP C0VHC9
D	1	SER	-	expression tag	UNP C0VHC9
E	1	SER	-	expression tag	UNP C0VHC9
F	1	SER	-	expression tag	UNP C0VHC9

- Molecule 2 is a RNA chain called 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	G	3	66	30	15	18	3	0	0	0
2	H	3	99	30	33	15	18	3	0	0
2	I	3	66	30	15	18	3	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	J	3	66	30	15	18	3	0	0	0
2	K	3	99	30	33	15	18	3	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0
3	B	1	5	4	1	0	0
3	C	1	5	4	1	0	0
3	C	1	5	4	1	0	0
3	D	1	5	4	1	0	0
3	E	1	5	4	1	0	0
3	F	1	5	4	1	0	0
3	F	1	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	237	Total 237	O 237	0	0
4	B	277	Total 277	O 277	0	0
4	C	242	Total 242	O 242	0	0
4	D	272	Total 272	O 272	0	0
4	E	229	Total 229	O 229	0	0
4	F	220	Total 220	O 220	0	0
4	G	7	Total 7	O 7	0	0
4	H	3	Total 3	O 3	0	0
4	J	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SAVED domain-containing protein




- Molecule 1: SAVED domain-containing protein




- Molecule 1: SAVED domain-containing protein











- Molecule 1: SAVED domain-containing protein



- Molecule 1: SAVED domain-containing protein



- Molecule 1: SAVED domain-containing protein



- Molecule 2: 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3')



A1
A2
A3

- Molecule 2: 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3')

Chain H: 100%

A1
A2
A3

- Molecule 2: 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3')

Chain I: 67% 33%

A1
A2
A3

- Molecule 2: 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3')

Chain J: 67% 33%

A1
A2
A3

- Molecule 2: 2'-5'-Linked Cyclic RNA (5'-R(P*AP*AP*A)-3')

Chain K: 33% 67% 33%

A1
A2
A3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.58Å 111.46Å 164.83Å 90.00° 100.21° 90.00°	Depositor
Resolution (Å)	49.21 – 2.10 49.21 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.1 (49.21-2.10) 98.1 (49.21-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.185 , 0.219 0.185 , 0.218	Depositor DCC
R_{free} test set	2000 reflections (0.92%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	23004	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3593	0.44	0/4856
1	B	0.27	0/3603	0.44	0/4869
1	C	0.27	0/3597	0.44	0/4861
1	D	0.27	0/3603	0.45	0/4869
1	E	0.27	0/3545	0.45	0/4788
1	F	0.26	0/3584	0.44	0/4841
2	G	7.87	26/74 (35.1%)	9.89	52/113 (46.0%)
2	H	7.98	26/74 (35.1%)	10.39	56/113 (49.6%)
2	I	8.02	30/74 (40.5%)	9.90	51/113 (45.1%)
2	J	8.01	29/74 (39.2%)	9.88	51/113 (45.1%)
2	K	8.05	26/74 (35.1%)	10.58	54/113 (47.8%)
All	All	1.07	137/21895 (0.6%)	1.47	264/29649 (0.9%)

The worst 5 of 137 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	2	A	N7-C5	29.72	1.57	1.39
2	H	2	A	N7-C5	28.97	1.56	1.39
2	G	2	A	N7-C5	28.29	1.56	1.39
2	J	2	A	N7-C5	28.14	1.56	1.39
2	I	2	A	N7-C5	28.12	1.56	1.39

The worst 5 of 264 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	2	A	C8-N9-C4	55.54	128.02	105.80
2	H	2	A	C8-N9-C4	55.37	127.95	105.80
2	I	2	A	C8-N9-C4	54.26	127.50	105.80
2	J	2	A	C8-N9-C4	54.04	127.42	105.80
2	G	2	A	C8-N9-C4	53.45	127.18	105.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3518	0	3466	27	0
1	B	3528	0	3474	29	0
1	C	3522	0	3469	23	0
1	D	3528	0	3474	23	0
1	E	3472	0	3424	47	0
1	F	3510	0	3454	38	0
2	G	66	0	33	0	0
2	H	66	33	33	1	0
2	I	66	0	33	3	0
2	J	66	0	33	1	0
2	K	66	33	33	13	0
3	A	5	0	0	0	0
3	B	5	0	0	3	0
3	C	10	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	10	0	0	0	0
4	A	237	0	0	10	0
4	B	277	0	0	13	0
4	C	242	0	0	2	0
4	D	272	0	0	12	0
4	E	229	0	0	11	0
4	F	220	0	0	14	0
4	G	7	0	0	0	0
4	H	3	0	0	0	0
4	J	3	0	0	0	0
All	All	22938	66	20926	190	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:160:ASP:OD1	4:E:601:HOH:O	1.83	0.96
1:E:427:PRO:HB3	2:K:2:A:N7	1.82	0.95
1:B:343:THR:OG1	4:B:602:HOH:O	1.90	0.90
1:B:227:GLU:O	4:B:601:HOH:O	1.90	0.89
1:E:72:SER:OG	4:E:602:HOH:O	1.92	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/462 (94%)	423 (98%)	11 (2%)	0	100	100
1	B	436/462 (94%)	427 (98%)	9 (2%)	0	100	100
1	C	435/462 (94%)	425 (98%)	10 (2%)	0	100	100
1	D	436/462 (94%)	429 (98%)	7 (2%)	0	100	100
1	E	426/462 (92%)	417 (98%)	9 (2%)	0	100	100
1	F	431/462 (93%)	421 (98%)	10 (2%)	0	100	100
All	All	2598/2772 (94%)	2542 (98%)	56 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/411 (95%)	388 (99%)	4 (1%)	76	82
1	B	393/411 (96%)	389 (99%)	4 (1%)	76	82
1	C	392/411 (95%)	387 (99%)	5 (1%)	69	75
1	D	393/411 (96%)	389 (99%)	4 (1%)	76	82
1	E	387/411 (94%)	385 (100%)	2 (0%)	88	92
1	F	391/411 (95%)	387 (99%)	4 (1%)	76	82
All	All	2348/2466 (95%)	2325 (99%)	23 (1%)	76	82

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	375	ASN
1	C	439	ASN
1	F	113	LEU
1	C	412	LYS
1	D	57	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	71	HIS
1	E	324	HIS
1	F	384	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	G	3/3 (100%)	1 (33%)	1 (33%)
2	H	3/3 (100%)	1 (33%)	1 (33%)
2	I	3/3 (100%)	1 (33%)	1 (33%)
2	J	2/3 (66%)	1 (50%)	0
2	K	2/3 (66%)	1 (50%)	0
All	All	13/15 (86%)	5 (38%)	3 (23%)

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	G	3	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	3	A
2	I	3	A
2	J	3	A
2	K	3	A

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	G	1	A
2	H	1	A
2	I	1	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	F	501	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	A	501	-	4,4,4	0.12	0	6,6,6	0.15	0
3	SO4	D	501	-	4,4,4	0.14	0	6,6,6	0.17	0
3	SO4	B	501	-	4,4,4	0.16	0	6,6,6	0.13	0
3	SO4	F	502	-	4,4,4	0.14	0	6,6,6	0.14	0
3	SO4	C	501	-	4,4,4	0.15	0	6,6,6	0.15	0
3	SO4	E	501	-	4,4,4	0.18	0	6,6,6	0.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	C	502	-	4,4,4	0.14	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	SO4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/462 (94%)	0.43	30 (6%) 17 21	28, 47, 77, 101	0
1	B	440/462 (95%)	0.33	19 (4%) 35 41	25, 45, 85, 111	0
1	C	439/462 (95%)	0.58	44 (10%) 7 9	28, 49, 99, 130	0
1	D	440/462 (95%)	0.35	17 (3%) 39 45	28, 45, 82, 110	0
1	E	432/462 (93%)	0.65	49 (11%) 5 6	28, 53, 108, 129	0
1	F	437/462 (94%)	0.85	80 (18%) 1 1	31, 54, 114, 148	0
2	G	3/3 (100%)	-0.69	0 100 100	50, 50, 55, 56	0
2	H	3/3 (100%)	0.99	0 100 100	66, 66, 68, 69	0
2	I	3/3 (100%)	0.84	0 100 100	95, 95, 95, 97	0
2	J	3/3 (100%)	-0.75	0 100 100	46, 46, 52, 53	0
2	K	3/3 (100%)	2.45	1 (33%) 0 0	76, 76, 77, 78	0
All	All	2641/2787 (94%)	0.53	240 (9%) 9 12	25, 49, 100, 148	0

The worst 5 of 240 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	133	VAL	8.4
1	C	347	ALA	7.6
1	C	133	VAL	7.4
1	C	345	ILE	6.9
1	F	413	SER	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	C	502	5/5	0.75	0.12	128,128,129,131	0
3	SO4	F	501	5/5	0.81	0.19	134,134,136,136	0
3	SO4	A	501	5/5	0.96	0.14	49,55,60,62	0
3	SO4	C	501	5/5	0.97	0.17	48,51,60,66	0
3	SO4	E	501	5/5	0.98	0.14	41,49,56,63	0
3	SO4	D	501	5/5	0.98	0.16	46,52,58,66	0
3	SO4	F	502	5/5	0.99	0.12	50,52,60,63	0
3	SO4	B	501	5/5	0.99	0.14	47,48,54,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.