



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 10:18 am BST

PDB ID : 4W1Y
Title : Crystal structure of Escherichia coli Tryptophanase in 'semi-holo' form
Authors : Goldgur, Y.
Deposited on : 2014-08-13
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	<p>2% 30% 49% 14% • 7%</p>
2	B	467	<p>2% 24% 54% 14% • 7%</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tryptophanase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	435	3428	2181	575	650	1	21	0	0	0

- Molecule 2 is a protein called Tryptophanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	433	3412	2175	575	641	21	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	B	1	5	4	1	0	0
3	B	1	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	42	Total 42	O 42	0	0
4	B	65	Total 65	O 65	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophanase



- Molecule 2: Tryptophanase



T465	PRO	H330	A268	S266	GLN	S75
A466	LYS	L331	K269	L207	GLY	
K467	THR	A332	K270	A208	HIS	S79
L468	GLY	Y333	D271	N209	SER	Y80
K469	LYS	R334	A272	L210	GLN	Y81
E470	GLN	I335	M273	K211	ILE	A82
V471	LEU	A336	V274	A272	ASN	L83
	PRO	Q337	F275	M213	GLY	A84
		C413	M276	Y214	CYS	E85
		P414	G277	S215	T149	S86
		E415	G278	I216	V153	V87
		E416	G279	I217	Y154	K88
		L417	L280	Y220	Y154	M89
		L418	C281	D221	I90	
		R419	M282	I222	E157	F91
		L420	K283	I222	A158	G92
		T421	D284	P223	F159	Y93
		I422	D285	V224	D160	Q94
		P423	G357	V225	T161	Y95
		R424	H358	M226	G162	T96
		A425	A359	D227	V163	T98
		T426	A360	S228	R164	T99
		T428	F361	A229	Y165	P98
		Q429	V362	R230	D166	H100
		T430	D363	F231	F167	Q101
		H431	A364	A232	K168	G102
		M432	L367	E233	G169	R103
		D433	L368	E233	M170	
		I435	I371	N234	F171	E106
		I436	P372	T296	D172	Q107
		I436	Q375	L297	L173	I108
		E437	F376	C298	E174	Y109
		A438	P377	V299	E177	I110
		F439	A378	V300	R178	P111
		H441	O379	Q301	G179	V112
		V442	A380	GLY	I113	I113
		K443	L381	PHE	I114	
		E444	A382	PRO	I180	
		M445	C383	THR	E181	
		A446	E384	TYR	E182	
		A447	E384	GLY	V183	
		M448	K387	G309	G184	
		I449	V388	L310	M187	
		G451	A389	E311	V188	
		L452	R392	A314	P189	
		F453	A393	M315	Y190	
		F454	V394	E316	I191	
		T455	E395	R255		
		Y456	I396	E256	T194	
		E457	G397	T257	I195	
		P458	S398	Y258	T196	
		K459	PHE	K259	S197	
		V460	LEU	Y260	M198	
		L461	LEU	A261	M198	
		R462	LEU	D262	S199	
		H463	GLY	M263	A200	
		F464	ARG	L264	G201	
			ASP	M266	G202	
				S267	Q203	
					P204	
					THR	
					THR	
					THR	

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	109.97Å 109.97Å 238.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 3.20 29.06 – 3.20	Depositor EDS
% Data completeness (in resolution range)	94.7 (15.00-3.20) 94.9 (29.06-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 3.18Å)	Xtrriage
Refinement program	REFMAC 5.4.0077	Depositor
R, R_{free}	0.214 , 0.290 0.212 , 0.288	Depositor DCC
R_{free} test set	1207 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	86.8	Xtrriage
Anisotropy	0.071	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 81.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6957	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	2/3473 (0.1%)	1.12	11/4692 (0.2%)
2	B	1.09	4/3478 (0.1%)	1.20	16/4694 (0.3%)
All	All	1.05	6/6951 (0.1%)	1.16	27/9386 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	GLU	CG-CD	6.80	1.62	1.51
2	B	281	CYS	CB-SG	-6.40	1.71	1.82
1	A	427	TYR	CD1-CE1	6.06	1.48	1.39
2	B	154	TYR	CD2-CE2	-5.92	1.30	1.39
2	B	74	TYR	CE1-CZ	5.05	1.45	1.38

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	266	MET	CB-CG-SD	-7.79	89.05	112.40
2	B	14	ARG	NE-CZ-NH1	-7.49	116.56	120.30
2	B	418	LEU	CA-CB-CG	7.07	131.56	115.30
2	B	7	LEU	CB-CG-CD2	-6.96	99.17	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	282	MET	CG-SD-CE	6.80	111.08	100.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	453	THR	Peptide
2	B	162	GLY	Peptide
2	B	309	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	0	3371	418	0
2	B	3412	0	3395	515	1
3	B	10	0	0	1	0
4	A	42	0	0	28	0
4	B	65	0	0	40	0
All	All	6957	0	6766	916	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 67.

The worst 5 of 916 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:22:THR:CG2	2:B:27:ARG:HH11	1.02	1.58
2:B:22:THR:CG2	2:B:27:ARG:HG3	1.39	1.52
2:B:450:LYS:HB2	2:B:470:GLU:CD	1.31	1.46
2:B:375:GLN:HE21	2:B:471:VAL:CG1	1.26	1.46
1:A:170:ASN:CB	1:A:209:ASN:ND2	1.79	1.45

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:103:ARG:NH1	2:B:301:GLN:O[8_665]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/467 (92%)	331 (77%)	81 (19%)	18 (4%)	3	20
2	B	425/467 (91%)	327 (77%)	83 (20%)	15 (4%)	3	24
All	All	855/934 (92%)	658 (77%)	164 (19%)	33 (4%)	3	22

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328	LEU
2	B	246	LYS
1	A	157	GLU
1	A	238	ILE
2	B	243	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/385 (92%)	286 (80%)	70 (20%)	1	7
2	B	358/386 (93%)	290 (81%)	68 (19%)	1	8
All	All	714/771 (93%)	576 (81%)	138 (19%)	1	8

5 of 138 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	430	THR
2	B	50	LEU
2	B	371	ILE
1	A	441	HIS
2	B	13	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	327	ASN
1	A	445	ASN
2	B	133	ASN
1	A	209	ASN
2	B	170	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	270	1	23,24,25	2.17	6 (26%)	25,32,34	2.43	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	270	1	-	4/16/17/19	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	270	LLP	O3-C3	-6.59	1.21	1.37
1	A	270	LLP	C4-C4'	3.71	1.53	1.46
1	A	270	LLP	C2-N1	2.99	1.39	1.33
1	A	270	LLP	C6-N1	2.81	1.40	1.34
1	A	270	LLP	C4'-NZ	2.63	1.36	1.27

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	LLP	OP4-C5'-C5	7.15	122.98	109.35
1	A	270	LLP	C4-C3-C2	6.41	124.16	120.19
1	A	270	LLP	CE-NZ-C4'	-3.67	107.62	118.90
1	A	270	LLP	OP3-P-OP1	3.06	122.68	110.68
1	A	270	LLP	C3-C2-N1	-2.44	117.62	120.77

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	270	LLP	C4-C5-C5'-OP4
1	A	270	LLP	C6-C5-C5'-OP4
1	A	270	LLP	C4-C4'-NZ-CE
1	A	270	LLP	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	270	LLP	6	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	500	-	4,4,4	0.14	0	6,6,6	0.60	0
3	SO4	B	501	-	4,4,4	0.17	0	6,6,6	0.20	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	434/467 (92%)	-0.32	10 (2%) 60 47	50, 82, 120, 126	0
2	B	433/467 (92%)	-0.27	9 (2%) 63 49	48, 82, 129, 170	0
All	All	867/934 (92%)	-0.29	19 (2%) 62 48	48, 82, 121, 170	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	471	VAL	5.9
2	B	414	PRO	4.6
2	B	413	CYS	3.8
1	A	415	ALA	3.2
2	B	398	SER	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	270	24/25	0.97	0.18	63,73,78,80	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	501	5/5	0.81	0.26	184,184,184,185	0
3	SO4	B	500	5/5	0.92	0.15	110,110,111,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.