Full wwPDB X-ray Structure Validation Report (i)

Dec 5, 2023 – 06:45 am GMT

PDB ID	:	1W6O
Title	:	X-RAY CRYSTAL STRUCTURE OF C2S HUMAN GALECTIN-1 COMPLEXED WITH LACTOSE
Authors	:	Lopez-Lucendo, M.I.F.; Gabius, H.J.; Romero, A.
Deposited on	:	2004-08-20
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	8%, 74%, 24%, ••
2	B	134	4%, 81%, 19%
3	C	2	100%
3	D	2	100%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 2380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GALECTIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	134	1030	648	178	198	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1002	SER	CYS	engineered mutation	UNP P09382
A	1065	ASP	GLY	engineered mutation	UNP P09382

- Molecule 2 is a protein called GALECTIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	134	1029	648	178	197	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2002	SER	CYS	engineered mutation	UNP P09382
B	2065	ASP	GLY	engineered mutation	UNP P09382

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	C	2	23	12	11	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	D	2	23	12	11	0	0	0

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
4	A	1	4	2	1	1	0	0
			Total	C	O	S		
4	A	1	4	2	1	1	0	0
			Total	C	O	S		
4	B	1	4	2	1	1	0	0
			Total	C	O	S		
4	B	1	4	2	1	1	0	0
			Total	C	O	S		
4	B	1	4	2	1	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	115	Total O 115 115	0	0
6	B	135	Total O 135 135	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GALECTIN-1

Chain A:





- Molecule 2: GALECTIN-1

Chain B:



- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain C:



- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	36.71Å 88.20Å 93.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.18 – 1.90 34.18 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (34.18-1.90) 99.4 (34.18-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	8.83 (at 1.81Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.224 0.198 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å^2)	14.4	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 46.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	2380	wwPDB-VP
Average B, all atoms (Å^2)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, SO4, GAL, CSO, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.33	0/1044	0.69	2/1412 (0.1%)
2	B	0.33	0/1051	0.62	0/1423
All	All	0.33	0/2095	0.65	2/2835 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	1001	ALA	C-N-CA	10.01	146.72	121.70
1	A	1001	ALA	CA-C-N	-6.44	103.03	117.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1001	ALA	Mainchain,Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1030	0	987	30	0
2	B	1029	0	988	23	0
3	C	23	0	21	0	0
3	D	23	0	21	0	0
4	A	8	0	10	2	0
4	B	12	0	15	1	0
5	B	5	0	0	0	0
6	A	115	0	0	2	1
6	B	135	0	0	1	1
All	All	2380	0	2042	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1134:ASP:OXT	2:B:2129:LYS:CE	2.09	1.01
1:A:1134:ASP:OXT	2:B:2129:LYS:HE3	1.63	0.99
1:A:1134:ASP:OXT	2:B:2129:LYS:NZ	2.17	0.77
2:B:2080:GLN:HG2	2:B:2083:SER:OG	1.86	0.76
1:A:1003:GLY:HA3	1:A:1122:ALA:O	1.94	0.66
2:B:2112:LEU:HB3	2:B:2114:LEU:HD13	1.77	0.65
2:B:2042:CYS:HB3	2:B:2114:LEU:HD22	1.81	0.61
2:B:2010:ASN:H	2:B:2118:ASN:ND2	2.00	0.59
1:A:1134:ASP:HB3	2:B:2129:LYS:HZ1	1.69	0.57
2:B:2020:ARG:HB3	2:B:2130:CYS:SG	2.45	0.56
2:B:2004:LEU:HD23	2:B:2005:VAL:N	2.20	0.56
1:A:1042:CYS:HB3	1:A:1114:LEU:HD22	1.87	0.55
1:A:1020:ARG:HB3	1:A:1130:CYS:SG	2.46	0.55
1:A:1133:PHE:O	1:A:1134:ASP:OXT	2.24	0.55
1:A:1122:ALA:HB2	1:A:1128:ILE:HD11	1.89	0.54
1:A:1001:ALA:N	1:A:1002:SER:CA	2.72	0.53
1:A:1127:LYS:HE2	6:A:2111:HOH:O	2.07	0.52
1:A:1009:LEU:HD13	1:A:1118:ASN:O	2.09	0.52
2:B:2004:LEU:HD23	2:B:2004:LEU:C	2.30	0.51
1:A:1001:ALA:N	1:A:1002:SER:HA	2.25	0.51
2:B:2010:ASN:H	2:B:2118:ASN:HD22	1.57	0.51
1:A:1024:ALA:HB3	1:A:1125:ASP:HB3	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1134:ASP:HB3	2:B:2129:LYS:NZ	2.25	0.50
2:B:2112:LEU:CB	2:B:2114:LEU:HD13	2.40	0.50
1:A:1086:GLU:HG2	4:A:2137:BME:H12	1.95	0.49
1:A:1030:PHE:HA	1:A:1124:GLY:HA3	1.95	0.49
1:A:1092:ASP:HB2	1:A:1095:ASN:O	2.13	0.49
1:A:1095:ASN:OD1	1:A:1109:PRO:HA	2.14	0.48
2:B:2100:LEU:HB3	2:B:2101:PRO:CD	2.44	0.47
2:B:2100:LEU:HB3	2:B:2101:PRO:HD2	1.96	0.47
2:B:2033:ASN:O	2:B:2120:MET:HA	2.15	0.46
1:A:1007:SER:O	1:A:1008:ASN:HB2	2.17	0.45
1:A:1030:PHE:CG	1:A:1126:PHE:HB2	2.52	0.45
1:A:1004:LEU:HD13	2:B:2009:LEU:HD12	1.99	0.44
1:A:1100:LEU:HB3	1:A:1101:PRO:HD2	1.99	0.44
2:B:2127:LYS:HE2	6:B:5128:HOH:O	2.16	0.44
1:A:1011:LEU:HB3	1:A:1117:ILE:HB	2.00	0.44
2:B:2086:GLU:O	4:B:3139:BME:H11	2.19	0.42
1:A:1093:GLN:NE2	1:A:1093:GLN:HA	2.34	0.42
1:A:1129:LYS:HE3	6:A:2110:HOH:O	2.19	0.42
1:A:1129:LYS:HE2	2:B:2133:PHE:O	2.21	0.41
1:A:1112:LEU:CB	1:A:1114:LEU:HD13	2.50	0.41
2:B:2024:ALA:HB3	2:B:2027:ALA:HB2	2.01	0.41
1:A:1099:LYS:HD2	4:A:2137:BME:S2	2.61	0.40
2:B:2112:LEU:HB3	2:B:2114:LEU:CD1	2.48	0.40
1:A:1100:LEU:HB3	1:A:1101:PRO:CD	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2097:HOH:O	6:B:5130:HOH:O[3_655]	2.05	0.15

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/134 (98%)	127 (97%)	3 (2%)	1 (1%)	19	9
2	B	132/134 (98%)	128 (97%)	3 (2%)	1 (1%)	19	9
All	All	263/268 (98%)	255 (97%)	6 (2%)	2 (1%)	19	9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1002	SER
2	B	2002	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/109 (100%)	107 (98%)	2 (2%)	59	55
2	B	110/110 (100%)	108 (98%)	2 (2%)	59	55
All	All	219/219 (100%)	215 (98%)	4 (2%)	59	55

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1080	GLN
1	A	1134	ASP
2	B	2032	LEU
2	B	2050	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1008	ASN
1	A	1080	GLN
1	A	1093	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1118	ASN
2	B	2050	ASN
2	B	2072	GLN
2	B	2118	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	CSO	A	1016	1	3,6,7	0.65	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	1016	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	BGC	C	1	3	12,12,12	1.43	1 (8%)	17,17,17	0.67	1 (5%)
3	GAL	C	2	3	11,11,12	1.63	3 (27%)	15,15,17	0.82	0
3	BGC	D	1	3	12,12,12	1.41	1 (8%)	17,17,17	0.67	1 (5%)
3	GAL	D	2	3	11,11,12	1.79	4 (36%)	15,15,17	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	C	1	3	-	0/2/22/22	0/1/1/1
3	GAL	C	2	3	-	0/2/19/22	0/1/1/1
3	BGC	D	1	3	-	1/2/22/22	0/1/1/1
3	GAL	D	2	3	-	0/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	GAL	O5-C5	3.25	1.50	1.43
3	D	2	GAL	C2-C3	2.85	1.56	1.52
3	C	1	BGC	C4-C5	2.82	1.59	1.53
3	C	2	GAL	C2-C3	2.76	1.56	1.52
3	C	2	GAL	O5-C5	2.72	1.49	1.43
3	D	1	BGC	C4-C5	2.57	1.58	1.53
3	D	2	GAL	O5-C1	2.40	1.47	1.43
3	D	2	GAL	C4-C5	2.16	1.57	1.53
3	C	2	GAL	C4-C5	2.08	1.57	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	1	BGC	C4-C3-C2	-2.06	107.23	110.82
3	D	1	BGC	C4-C3-C2	-2.05	107.25	110.82

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Oligosaccharide Chain D



5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	BME	A	2138	1	3,3,3	0.29	0	1,2,2	0.15	0
4	BME	B	3140	2	3,3,3	0.29	0	1,2,2	0.09	0
5	SO4	B	3135	-	4,4,4	0.76	0	6,6,6	0.34	0
4	BME	B	3139	2	3,3,3	0.26	0	1,2,2	0.11	0
4	BME	B	3138	2	3,3,3	0.30	0	1,2,2	0.18	0
4	BME	A	2137	1	3,3,3	0.32	0	1,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	A	2138	1	-	1/1/1/1	-
4	BME	B	3140	2	-	0/1/1/1	-
4	BME	B	3139	2	-	0/1/1/1	-
4	BME	B	3138	2	-	0/1/1/1	-
4	BME	A	2137	1	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2138	BME	O1-C1-C2-S2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	3139	BME	1	0
4	A	2137	BME	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	133/134 (99%)	0.57	11 (8%)	11	13	9, 18, 28, 44	0
2	B	134/134 (100%)	0.02	5 (3%)	41	44	6, 13, 22, 36	0
All	All	267/268 (99%)	0.29	16 (5%)	21	24	6, 14, 28, 44	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1001	ALA	29.3
2	B	2001	ALA	10.7
1	A	1134	ASP	4.5
1	A	1002	SER	4.5
2	B	2134	ASP	3.6
2	B	2002	SER	3.4
2	B	2026	ASP	3.2
1	A	1025	PRO	2.9
1	A	1082	GLY	2.7
1	A	1026	ASP	2.7
1	A	1084	VAL	2.5
2	B	2080	GLN	2.4
1	A	1127	LYS	2.3
1	A	1067	ALA	2.2
1	A	1112	LEU	2.1
1	A	1083	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	CSO	A	1016	7/8	0.86	0.12	19,22,24,30	0

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BGC	D	1	12/12	0.68	0.34	20,23,25,26	12
3	GAL	D	2	11/12	0.72	0.38	6,19,22,22	11
3	BGC	C	1	12/12	0.76	0.20	25,29,32,33	0
3	GAL	C	2	11/12	0.94	0.10	15,20,23,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	BME	B	3140	4/4	0.66	0.28	36,39,41,44	0
4	BME	A	2138	4/4	0.83	0.21	33,35,37,40	0
4	BME	A	2137	4/4	0.85	0.25	30,31,32,34	0
4	BME	B	3139	4/4	0.93	0.16	24,25,27,30	0
5	SO4	B	3135	5/5	0.93	0.13	45,45,45,46	0
4	BME	B	3138	4/4	0.94	0.12	26,27,28,29	0

6.5 Other polymers (i)

There are no such residues in this entry.