



## wwPDB EM Validation Summary Report ⓘ

Aug 17, 2026 – 12:54 PM JST

PDB ID : 9WEZ / pdb\_00009wez  
EMDB ID : EMD-65919  
Title : Cryo-EM structure of human pyruvate kinase R (PKR) in complex with FBP  
Authors : Ma, H.Y.; Su, Z.M.  
Deposited on : 2025-08-20  
Resolution : 2.64 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.50

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-----------------------------|----------------------------------------------------------|
| Clashscore            | 229148                      | 23984                       | -                                                        |
| Ramachandran outliers | 224038                      | 23583                       | -                                                        |
| Sidechain outliers    | 223484                      | 23102                       | -                                                        |
| Q-score               | -                           | 25397                       | 8968 ( 2.14 - 3.14 )                                     |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain                                                       |
|-----|-------|--------|------------------------------------------------------------------------|
| 1   | A     | 574    | <div> <div>5%</div> <div>64%</div> <div>7%</div> <div>29%</div> </div> |
| 1   | B     | 574    | <div> <div>6%</div> <div>68%</div> <div>•</div> <div>29%</div> </div>  |
| 1   | C     | 574    | <div> <div>6%</div> <div>66%</div> <div>5%</div> <div>29%</div> </div> |
| 1   | D     | 574    | <div> <div>5%</div> <div>67%</div> <div>5%</div> <div>29%</div> </div> |

## 2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 12424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase PKLR.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1   | A     | 408      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3085  | 1932 | 564 | 573 | 16 |         |       |
| 1   | B     | 408      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3085  | 1932 | 564 | 573 | 16 |         |       |
| 1   | C     | 408      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3087  | 1934 | 564 | 573 | 16 |         |       |
| 1   | D     | 408      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3087  | 1934 | 564 | 573 | 16 |         |       |

- Molecule 2 is 1,6-di-O-phosphono-alpha-D-fructofuranose (CCD ID: AFP) (formula:  $C_6H_{14}O_{12}P_2$ ) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms |   |    |   | AltConf |
|-----|-------|----------|-------|---|----|---|---------|
| 2   | A     | 1        | Total | C | O  | P | 0       |
|     |       |          | 20    | 6 | 12 | 2 |         |
| 2   | B     | 1        | Total | C | O  | P | 0       |
|     |       |          | 20    | 6 | 12 | 2 |         |

*Continued on next page...*


*Continued from previous page...*

| Mol | Chain | Residues | Atoms |   |    |   | AltConf |
|-----|-------|----------|-------|---|----|---|---------|
| 2   | C     | 1        | Total | C | O  | P | 0       |
|     |       |          | 20    | 6 | 12 | 2 |         |
| 2   | D     | 1        | Total | C | O  | P | 0       |
|     |       |          | 20    | 6 | 12 | 2 |         |


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Pyruvate kinase PKLR



#### • Molecule 1: Pyruvate kinase PKLR





• Molecule 1: Pyruvate kinase PKLR



MET SER ILE GLN GLU MET ASN ILE SER SER LEU GLN LEU ARG SER TRP VAL SER ASP LYS SER SER GLN ARG ASP LEU ALA LYS SER ILE ILE ILE GLY THR PRO GLY PRO GLY ALA TYR LEU ARG ALA VAL SER ARG ALA VAL LEU VAL ALA GLN THR GLN GLU LEU GLY THR ALA PHE PHE GLN GLN

LEU PRO ALA ALA MET ASP THR PHE LEU PRO ASN HIS LEU CYS LEU VAL ASP I778 D79 S80 E81 P82 V83 A84 E161 ILE ARG THR ILE LEU GLN GLY PRO GLU SER LEU VAL THR VAL GLN SER GLY THR VAL VAL GLN LYS GLY SER GLN THR VAL LEU VAL THR ARG ASP PRO PHE ARG THR ARG GLY

ASN ALA ASN THR VAL TRP ASP TYR PHE LEU PRO ASN ILE ARG VAL VAL VAL PRO VAL GLY GLY ARG ILE TYR ILE ASP ASP GLY LEU ILE SER LEU VAL VAL GLN LYS GLY SER GLY VAL LEU VAL THR LYS G251 L254 P255 G256

A257 Q258 V259 D260 L261 G263 L264 S265 E266 D271 L272 R273 D293 K313 H317 E325 V329 S330 D331 M334 D339 I342 E343 I344 P345 A346 E347 K348 M355 R359 T371 Q372 M373 L374 E387 E407 E427 Y433 H434 R435 E439

K465 R498 A502 V506 E519 I524 D527 D528 R547 G561 N566 R569 W570 L571 S574

• Molecule 1: Pyruvate kinase PKLR



MET SER ILE GLN GLU ASN ILE SER SER LEU GLN LEU ARG SER TRP VAL VAL SER LYS SER SER GLN ARG ASP LEU ALA LYS SER ILE ILE ILE GLY THR PRO GLY PRO GLY ALA TYR LEU ARG ALA VAL SER ARG ALA VAL LEU VAL ALA GLN THR GLN GLU LEU GLY THR ALA PHE PHE GLN GLN

LEU PRO ALA ALA MET ASP THR PHE LEU PRO ASN HIS LEU CYS LEU VAL ASP I778 D79 S80 E81 P82 V83 A84 E161 ILE ARG THR ILE LEU GLN GLY PRO GLU SER LEU VAL THR VAL GLN LYS GLY SER GLY THR VAL VAL GLN LYS G251 L254 P255 G256

ASN ALA ASN THR VAL TRP VAL ASP TYR PHE LEU PRO ASN ILE ARG VAL VAL VAL PRO VAL GLY GLY ARG ILE TYR ILE ASP ASP GLY LEU ILE SER LEU VAL VAL GLN LYS GLY SER GLN THR VAL LEU VAL THR ARG LYS G251 L254 P255 G256

A257 Q258 V259 D260 L261 G263 L264 S265 E266 D271 R289 D293 L301 I310 H317 A336 D339 L340 E343 I344 P345 A346 A370 T371 Q372 M373 L374 M377 R382 E387 D400 E407 A430 R435 E439 V454 I457 A502

V506 I524 D527 R547 R569 W570 L571 S574

## 4 Experimental information

| Property                             | Value                                   | Source    |
|--------------------------------------|-----------------------------------------|-----------|
| EM reconstruction method             | SINGLE PARTICLE                         | Depositor |
| Imposed symmetry                     | POINT, C2                               | Depositor |
| Number of particles used             | 837441                                  | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF                       | Depositor |
| CTF correction method                | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope                           | FEI POLARA 300                          | Depositor |
| Voltage (kV)                         | 300                                     | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 60                                      | Depositor |
| Minimum defocus (nm)                 | 800                                     | Depositor |
| Maximum defocus (nm)                 | 2100                                    | Depositor |
| Magnification                        | Not provided                            |           |
| Image detector                       | GATAN K2 QUANTUM (4k x 4k)              | Depositor |
| Maximum map value                    | 1.669                                   | Depositor |
| Minimum map value                    | -0.002                                  | Depositor |
| Average map value                    | 0.003                                   | Depositor |
| Map value standard deviation         | 0.036                                   | Depositor |
| Recommended contour level            | 0.14                                    | Depositor |
| Map size (Å)                         | 217.6, 217.6, 217.6                     | wwPDB     |
| Map dimensions                       | 256, 256, 256                           | wwPDB     |
| Map angles (°)                       | 90.0, 90.0, 90.0                        | wwPDB     |
| Pixel spacing (Å)                    | 0.85, 0.85, 0.85                        | Depositor |

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AFP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |             | Bond angles |             |
|-----|-------|--------------|-------------|-------------|-------------|
|     |       | RMSZ         | $\# Z  > 5$ | RMSZ        | $\# Z  > 5$ |
| 1   | A     | 0.22         | 0/3135      | 0.45        | 0/4240      |
| 1   | B     | 0.22         | 0/3135      | 0.45        | 0/4240      |
| 1   | C     | 0.16         | 0/3137      | 0.39        | 0/4243      |
| 1   | D     | 0.16         | 0/3137      | 0.37        | 0/4243      |
| All | All   | 0.19         | 0/12544     | 0.42        | 0/16966     |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 3085  | 0        | 3136     | 35      | 0            |
| 1   | B     | 3085  | 0        | 3136     | 11      | 0            |
| 1   | C     | 3087  | 0        | 3143     | 18      | 0            |
| 1   | D     | 3087  | 0        | 3143     | 18      | 0            |
| 2   | A     | 20    | 0        | 9        | 0       | 0            |
| 2   | B     | 20    | 0        | 9        | 0       | 0            |
| 2   | C     | 20    | 0        | 9        | 1       | 0            |
| 2   | D     | 20    | 0        | 9        | 0       | 0            |
| All | All   | 12424 | 0        | 12594    | 80      | 0            |



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1          | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 1:A:341:GLY:HA2 | 1:A:344:ILE:CD1  | 1.44                     | 1.47              |
| 1:A:341:GLY:CA  | 1:A:344:ILE:HD12 | 1.79                     | 1.12              |
| 1:A:341:GLY:HA2 | 1:A:344:ILE:HD11 | 1.48                     | 0.96              |
| 1:A:341:GLY:CA  | 1:A:344:ILE:CD1  | 2.39                     | 0.95              |
| 1:A:341:GLY:HA2 | 1:A:344:ILE:HD12 | 0.97                     | 0.94              |

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Favoured   | Allowed | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1   | A     | 404/574 (70%)   | 386 (96%)  | 18 (4%) | 0        | 100         | 100 |
| 1   | B     | 404/574 (70%)   | 389 (96%)  | 15 (4%) | 0        | 100         | 100 |
| 1   | C     | 404/574 (70%)   | 388 (96%)  | 16 (4%) | 0        | 100         | 100 |
| 1   | D     | 404/574 (70%)   | 391 (97%)  | 13 (3%) | 0        | 100         | 100 |
| All | All   | 1616/2296 (70%) | 1554 (96%) | 62 (4%) | 0        | 100         | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Rotameric   | Outliers | Percentiles |     |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 1   | A     | 322/460 (70%)   | 322 (100%)  | 0        | 100         | 100 |
| 1   | B     | 322/460 (70%)   | 322 (100%)  | 0        | 100         | 100 |
| 1   | C     | 323/460 (70%)   | 323 (100%)  | 0        | 100         | 100 |
| 1   | D     | 323/460 (70%)   | 323 (100%)  | 0        | 100         | 100 |
| All | All   | 1290/1840 (70%) | 1290 (100%) | 0        | 100         | 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | D     | 267 | GLN  |
| 1   | D     | 393 | ASN  |
| 1   | D     | 421 | GLN  |
| 1   | B     | 434 | HIS  |
| 1   | C     | 267 | GLN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths |      |             | Bond angles |      |             |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
|     |      |       |     |      | Counts       | RMSZ | $\# Z  > 2$ | Counts      | RMSZ | $\# Z  > 2$ |
| 2   | AFP  | C     | 601 | -    | 18,20,20     | 3.69 | 5 (27%)     | 23,32,32    | 0.77 | 0           |
| 2   | AFP  | D     | 601 | -    | 18,20,20     | 3.68 | 5 (27%)     | 23,32,32    | 0.73 | 0           |
| 2   | AFP  | A     | 601 | -    | 18,20,20     | 3.67 | 5 (27%)     | 23,32,32    | 0.73 | 0           |
| 2   | AFP  | B     | 601 | -    | 18,20,20     | 3.68 | 5 (27%)     | 23,32,32    | 0.76 | 0           |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions   | Rings   |
|-----|------|-------|-----|------|---------|------------|---------|
| 2   | AFP  | C     | 601 | -    | -       | 6/13/32/32 | 0/1/1/1 |
| 2   | AFP  | D     | 601 | -    | -       | 3/13/32/32 | 0/1/1/1 |
| 2   | AFP  | A     | 601 | -    | -       | 3/13/32/32 | 0/1/1/1 |
| 2   | AFP  | B     | 601 | -    | -       | 2/13/32/32 | 0/1/1/1 |

The worst 5 of 20 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z     | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 2   | B     | 601 | AFP  | C4-C5 | -8.79 | 1.30        | 1.53     |
| 2   | A     | 601 | AFP  | C4-C5 | -8.76 | 1.30        | 1.53     |
| 2   | D     | 601 | AFP  | C4-C5 | -8.75 | 1.30        | 1.53     |
| 2   | C     | 601 | AFP  | C4-C5 | -8.65 | 1.30        | 1.53     |
| 2   | C     | 601 | AFP  | O5-C2 | -8.39 | 1.30        | 1.43     |

There are no bond angle outliers.

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


| Mol | Chain | Res | Type | Atoms        |
|-----|-------|-----|------|--------------|
| 2   | A     | 601 | AFP  | O1-C1-C2-C3  |
| 2   | A     | 601 | AFP  | O1-C1-C2-O5  |
| 2   | C     | 601 | AFP  | C6-O6-P6-O4P |
| 2   | D     | 601 | AFP  | O1-C1-C2-C3  |
| 2   | D     | 601 | AFP  | O1-C1-C2-O5  |


There are no ring outliers.


1 monomer is involved in 1 short contact:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 2   | C     | 601 | AFP  | 1       | 0            |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-65919. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)

### 6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


### 6.2.2 Raw map



X Index: 128



Y Index: 128



Z Index: 128

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 125




Y Index: 90




Z Index: 156


### 6.3.2 Raw map



X Index: 125



Y Index: 113




Z Index: 165


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.14. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 56  $\text{nm}^3$ ; this corresponds to an approximate mass of 50 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of  $0.379 \text{ \AA}^{-1}$



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.379  $\text{\AA}^{-1}$

## 8.2 Resolution estimates


| Resolution estimate (Å)   | Estimation criterion (FSC cut-off) |      |          |
|---------------------------|------------------------------------|------|----------|
|                           | 0.143                              | 0.5  | Half-bit |
| Reported by author        | 2.64                               | -    | -        |
| Author-provided FSC curve | 2.64                               | 2.98 | 2.67     |
| Unmasked-calculated*      | 3.13                               | 3.53 | 3.17     |

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.13 differs from the reported value 2.64 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-65919 and PDB model 9WEZ. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.14 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.14).

## 9.4 Atom inclusion ⓘ



At the recommended contour level, 84% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.14) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion     | Q-score            |
|-------|--------------------|--------------------|
| All   | <div></div> 0.7770 | <div></div> 0.5400 |
| A     | <div></div> 0.7830 | <div></div> 0.5380 |
| B     | <div></div> 0.7720 | <div></div> 0.5390 |
| C     | <div></div> 0.7750 | <div></div> 0.5410 |
| D     | <div></div> 0.7780 | <div></div> 0.5420 |

