



Full wwPDB EM Validation Report ⓘ

Aug 17, 2026 – 02:12 PM JST

PDB ID : 9WHX / pdb_00009whx
EMDB ID : EMD-65978
Title : Cryo-EM structure of the IS621 recombinase in complex with bridge RNA, left-half DNA, and right-half DNA in the pre-strand exchange state
Authors : Hiraizumi, M.; Tsujimoto, E.; Shiojiri, N.; Nagahata, N.; Yamashita, K.; Nishimasu, H.
Deposited on : 2025-08-27
Resolution : 2.50 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	7115 (2.00 - 3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	 89% 5% • 5%
1	B	328	 91% • • •
1	C	328	 90% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	328	89% 8%
2	E	180	23% 7% 67%
2	F	180	20% 11% 67%
3	G	38	74% 24%
4	H	38	58% 16% 24%
5	I	44	52% 9% 39%
6	J	44	45% 16% 39%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 14677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IS621 transposase.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	311	Total	C	N	O	S	0	0
			2452	1540	464	436	12		
1	B	318	Total	C	N	O	S	0	0
			2503	1569	473	449	12		
1	C	308	Total	C	N	O	S	0	0
			2427	1526	458	431	12		
1	D	317	Total	C	N	O	S	0	0
			2494	1564	472	446	12		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A0E0Y1P1
A	0	PRO	-	expression tag	UNP A0A0E0Y1P1
B	-1	GLY	-	expression tag	UNP A0A0E0Y1P1
B	0	PRO	-	expression tag	UNP A0A0E0Y1P1
C	-1	GLY	-	expression tag	UNP A0A0E0Y1P1
C	0	PRO	-	expression tag	UNP A0A0E0Y1P1
D	-1	GLY	-	expression tag	UNP A0A0E0Y1P1
D	0	PRO	-	expression tag	UNP A0A0E0Y1P1

- Molecule 2 is a RNA chain called bridge RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	59	Total	C	N	O	P	0	0
			1249	557	211	422	59		
2	F	59	Total	C	N	O	P	0	0
			1252	560	218	415	59		

- Molecule 3 is a DNA chain called LH, top strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	29	Total	C	N	O	P	0	0
			580	278	97	176	29		

- Molecule 4 is a DNA chain called LH, bottom strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	29	Total	C	N	O	P	0	0
			607	286	122	170	29		

- Molecule 5 is a DNA chain called RH, top strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	27	Total	C	N	O	P	0	0
			555	264	105	159	27		

- Molecule 6 is a DNA chain called RH, bottom strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	27	Total	C	N	O	P	0	0
			552	264	96	165	27		

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
7	A	1	Total	Mg	0
			1	1	
7	C	1	Total	Mg	0
			1	1	

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		AltConf
8	A	2	Total	O	0
			2	2	
8	C	2	Total	O	0
			2	2	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IS621 transposase

Chain A: 



- Molecule 1: IS621 transposase

Chain B: 




- Molecule 1: IS621 transposase

Chain C: 



- Molecule 1: IS621 transposase


Chain D: 




- Molecule 2: bridge RNA

Chain E: 






- Molecule 2: bridge RNA




- Molecule 3: LH, top strand




- Molecule 4: LH, bottom strand



- Molecule 5: RH, top strand



- Molecule 6: RH, bottom strand



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	558374	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.256	Depositor
Minimum map value	-0.070	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.013	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	172.64, 172.64, 172.64	wwPDB
Map dimensions	208, 208, 208	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.83, 0.83, 0.83	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2499	1.10	2/3365 (0.1%)
1	B	0.59	0/2552	1.06	4/3438 (0.1%)
1	C	0.59	0/2474	1.08	2/3332 (0.1%)
1	D	0.63	0/2543	1.09	4/3426 (0.1%)
2	E	0.59	0/1392	1.08	3/2166 (0.1%)
2	F	0.58	0/1398	1.17	10/2175 (0.5%)
3	G	0.43	0/645	1.03	1/987 (0.1%)
4	H	0.41	0/684	0.93	1/1056 (0.1%)
5	I	0.43	0/622	0.95	0/955
6	J	0.46	0/617	1.08	1/950 (0.1%)
All	All	0.58	0/15426	1.08	28/21850 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	132	U	O3'-P-O5'	12.37	122.55	104.00
2	F	138	G	O3'-P-O5'	-8.38	91.43	104.00
6	J	21	DG	O3'-P-O5'	-8.33	91.50	104.00
2	F	168	U	O3'-P-O5'	7.91	115.87	104.00
2	E	89	A	O3'-P-O5'	-6.78	93.82	104.00
3	G	29	DA	O3'-P-O5'	-6.51	94.23	104.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	260	ARG	N-CA-CB	-6.31	100.50	110.22
1	C	105	ASP	CA-CB-CG	6.16	118.75	112.60
1	C	11	ASP	CA-CB-CG	6.09	118.69	112.60
4	H	25	DA	C4'-C3'-O3'	5.65	118.48	110.00
2	F	134	G	O3'-P-O5'	5.59	112.38	104.00
2	E	71	U	O3'-P-O5'	-5.57	95.65	104.00
1	D	11	ASP	CA-CB-CG	5.49	118.09	112.60
2	F	149	U	O3'-P-O5'	-5.44	95.84	104.00
2	F	133	G	C2'-C3'-O3'	-5.41	105.59	113.70
2	F	156	U	O3'-P-O5'	-5.38	95.93	104.00
1	B	208	THR	CA-CB-OG1	-5.22	101.77	109.60
2	E	52	C	C4'-C3'-C2'	-5.20	97.40	102.60
1	D	211	VAL	N-CA-CB	5.19	116.62	110.55
1	A	208	THR	CA-CB-OG1	-5.17	101.84	109.60
2	F	132	U	C2'-C3'-O3'	5.17	117.26	109.50
1	B	11	ASP	CA-CB-CG	5.14	117.75	112.60
2	F	137	C	O3'-P-O5'	-5.13	96.30	104.00
1	D	60	GLU	CB-CA-C	5.12	118.18	109.53
1	B	60	GLU	N-CA-CB	5.09	118.48	110.23
1	A	117	ARG	O-C-N	-5.07	119.20	121.53
2	F	126	G	C1'-C2'-O2'	5.02	115.93	108.40
1	D	98	ARG	N-CA-CB	-5.02	103.92	111.25

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	157	A	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2452	0	2494	7	0
1	B	2503	0	2537	6	0
1	C	2427	0	2473	4	0
1	D	2494	0	2531	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	1249	0	632	7	0
2	F	1252	0	633	7	0
3	G	580	0	328	0	0
4	H	607	0	325	11	0
5	I	555	0	304	3	0
6	J	552	0	307	7	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
8	A	2	0	0	0	0
8	C	2	0	0	0	0
All	All	14677	0	12564	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:J:12:DG:H4'	6:J:13:DT:H5'	1.55	0.88
4:H:24:DA:H5'	4:H:24:DA:C8	2.27	0.69
1:B:89:LYS:HD2	4:H:27:DG:H5''	1.78	0.65
6:J:11:DT:H2''	6:J:12:DG:H5'	1.81	0.61
2:F:124:C:H2'	2:F:125:A:O4'	2.02	0.60
1:A:95:GLU:OE1	1:D:116:LYS:NZ	2.33	0.58
5:I:23:DA:H2''	5:I:24:DG:H5'	1.84	0.58
2:E:51:U:H2'	2:E:52:C:O4'	2.04	0.58
1:A:301:LYS:NZ	2:E:44:U:OP1	2.38	0.57
1:B:194:ARG:NH1	1:B:217:GLY:O	2.37	0.57
1:D:98:ARG:NH2	2:E:83:U:OP1	2.40	0.55
2:E:75:C:H2'	2:E:76:U:O4'	2.08	0.53
1:B:250:ARG:HE	4:H:23:DC:H4'	1.74	0.53
5:I:16:DA:H5''	5:I:16:DA:C8	2.44	0.52
1:D:253:LYS:HZ2	6:J:14:DA:H2	1.56	0.52
1:D:238:ARG:NH2	1:D:240:GLU:OE2	2.41	0.52
6:J:7:DC:H2''	6:J:8:DA:C8	2.44	0.52
1:A:254:ALA:O	1:D:98:ARG:NH1	2.44	0.51
1:D:98:ARG:NH2	2:E:83:U:H5''	2.26	0.50
4:H:23:DC:H2''	4:H:24:DA:H5''	1.96	0.47
1:B:226:ARG:HD3	1:B:226:ARG:HA	1.66	0.46
1:B:101:THR:C	1:B:103:THR:H	2.24	0.46
4:H:27:DG:H2'	4:H:28:DC:C6	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:67:GLU:N	1:C:68:PRO:CD	2.80	0.45
1:A:238:ARG:HD3	2:E:85:A:C2	2.51	0.45
2:F:123:A:H2'	2:F:124:C:O4'	2.17	0.45
4:H:24:DA:H2''	4:H:25:DA:OP1	2.18	0.44
1:D:250:ARG:HD2	6:J:13:DT:O2	2.18	0.44
1:A:107:ARG:HE	1:A:107:ARG:HB2	1.36	0.43
4:H:24:DA:N3	6:J:12:DG:N2	2.67	0.43
1:D:256:HIS:NE2	2:F:131:A:OP2	2.45	0.43
5:I:15:DC:H2''	5:I:16:DA:C5	2.54	0.43
1:D:61:ALA:HB2	1:D:82:VAL:HG12	2.01	0.43
4:H:24:DA:H5'	4:H:24:DA:N9	2.34	0.43
1:A:141:LEU:HD12	1:A:177:LEU:HD12	2.00	0.42
1:C:301:LYS:NZ	2:F:117:A:OP1	2.48	0.42
4:H:23:DC:C2'	4:H:24:DA:H5''	2.50	0.42
1:B:226:ARG:NH2	2:E:60:G:O6	2.52	0.42
1:D:194:ARG:NH1	1:D:217:GLY:O	2.52	0.42
1:D:222:PHE:HB2	1:D:307:TYR:CZ	2.55	0.42
1:D:274:THR:CG2	1:D:276:TRP:CD1	3.04	0.41
1:D:301:LYS:NZ	2:F:151:A:OP1	2.48	0.41
1:C:89:LYS:HZ2	1:C:89:LYS:HG3	1.73	0.41
1:D:144:MET:HB3	1:D:144:MET:HE2	1.84	0.41
4:H:21:DT:H5'	4:H:22:DA:OP1	2.21	0.41
6:J:11:DT:H6	6:J:11:DT:H2'	1.76	0.41
4:H:22:DA:H2''	4:H:23:DC:C6	2.56	0.41
1:D:111:GLU:OE2	1:D:114:ARG:NH1	2.53	0.41
1:C:289:LYS:NZ	2:F:120:C:OP1	2.54	0.41
1:D:283:ARG:NH2	2:F:152:U:OP1	2.54	0.40
1:A:265:MET:SD	1:A:268:MET:CE	3.10	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/328 (94%)	306 (100%)	1 (0%)	0	100	100
1	B	316/328 (96%)	312 (99%)	4 (1%)	0	100	100
1	C	304/328 (93%)	300 (99%)	4 (1%)	0	100	100
1	D	315/328 (96%)	313 (99%)	2 (1%)	0	100	100
All	All	1242/1312 (95%)	1231 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/265 (95%)	241 (96%)	10 (4%)	28	54
1	B	257/265 (97%)	245 (95%)	12 (5%)	23	47
1	C	249/265 (94%)	241 (97%)	8 (3%)	34	62
1	D	256/265 (97%)	253 (99%)	3 (1%)	63	83
All	All	1013/1060 (96%)	980 (97%)	33 (3%)	34	61

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	MET
1	A	89	LYS
1	A	95	GLU
1	A	107	ARG
1	A	154	THR
1	A	216	ILE
1	A	237	ARG
1	A	250	ARG
1	A	251	MET
1	A	257	VAL
1	B	60	GLU
1	B	101	THR
1	B	103	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	105	ASP
1	B	135	VAL
1	B	136	VAL
1	B	153	GLU
1	B	177	LEU
1	B	226	ARG
1	B	250	ARG
1	B	258	SER
1	B	284	LEU
1	C	66	MET
1	C	89	LYS
1	C	135	VAL
1	C	168	LEU
1	C	175	LYS
1	C	182	LYS
1	C	185	THR
1	C	250	ARG
1	D	135	VAL
1	D	136	VAL
1	D	311	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	HIS
1	B	115	GLN
1	B	193	HIS
1	C	54	HIS
1	C	115	GLN
1	C	139	GLN
1	C	147	GLN
1	C	224	HIS
1	D	304	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	58/180 (32%)	10 (17%)	3 (5%)
2	F	58/180 (32%)	8 (13%)	0
All	All	116/360 (32%)	18 (15%)	3 (2%)

All (18) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	41	G
2	E	46	C
2	E	59	U
2	E	60	G
2	E	70	C
2	E	76	U
2	E	85	A
2	E	86	U
2	E	87	G
2	E	90	G
2	F	115	U
2	F	126	G
2	F	133	G
2	F	135	A
2	F	138	G
2	F	141	U
2	F	169	G
2	F	171	U

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	E	52	C
2	E	59	U
2	E	86	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	G	1
5	I	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	G	25:DT	O3'	26:DT	P	3.25
1	I	20:DT	O3'	21:DA	P	2.96


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-65978. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 104




Y Index: 104




Z Index: 104


6.2.2 Raw map



X Index: 104



Y Index: 104




Z Index: 104


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 91




Y Index: 91




Z Index: 82


6.3.2 Raw map



X Index: 91



Y Index: 91




Z Index: 82


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_65978_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 170 nm³; this corresponds to an approximate mass of 153 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.400 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.400 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.50	-	-
Author-provided FSC curve	2.53	2.83	2.58
Unmasked-calculated*	2.74	3.00	2.79

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-65978 and PDB model 9WHX. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


9.1 Map-model overlay [i](#)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9770	0.5920
A	0.9770	0.6090
B	0.9730	0.5910
C	0.9710	0.5860
D	0.9760	0.6030
E	0.9920	0.5810
F	0.9970	0.6100
G	0.9850	0.5670
H	0.9740	0.5730
I	0.9960	0.5730
J	0.9870	0.5500

1.0

0.0

<0.0