Full wwPDB X-ray Structure Validation Report (i)

Jun 16, 2024 – 08:12 AM EDT

PDB ID	:	4WLN
Title	:	Crystal structure of apo MDH2
Authors	:	Eo, Y.M.; Han, B.G.; Ahn, H.C.
Deposited on	:	2014-10-07
Resolution	:	2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	% 89% • 8%
1	B	340	4% 88% • • 8%
1	C	340	6% 88% • 8%
1	D	340	5% 87% 5% 8%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malate dehydrogenase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2312	1472	390	437	13	0	0	0
1	B	314	2312	1472	390	437	13	0	0	0
1	C	314	2312	1472	390	437	13	0	0	0
1	D	314	2312	1472	390	437	13	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P40926
A	0	GLY	-	expression tag	UNP P40926
A	1	SER	-	expression tag	UNP P40926
A	2	SER	-	expression tag	UNP P40926
A	3	HIS	-	expression tag	UNP P40926
A	4	HIS	-	expression tag	UNP P40926
A	5	HIS	-	expression tag	UNP P40926
A	6	HIS	-	expression tag	UNP P40926
A	7	HIS	-	expression tag	UNP P40926
A	8	HIS	-	expression tag	UNP P40926
A	9	SER	-	expression tag	UNP P40926
A	10	SER	-	expression tag	UNP P40926
A	11	GLY	-	expression tag	UNP P40926
A	12	LEU	-	expression tag	UNP P40926
A	13	VAL	-	expression tag	UNP P40926
A	14	PRO	-	expression tag	UNP P40926
A	15	ARG	-	expression tag	UNP P40926
A	16	GLY	-	expression tag	UNP P40926
A	17	SER	-	expression tag	UNP P40926
A	18	HIS	-	expression tag	UNP P40926
A	19	MET	-	expression tag	UNP P40926

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MET	-	expression tag	UNP P40926
B	0	GLY	-	expression tag	UNP P40926
B	1	SER	-	expression tag	UNP P40926
B	2	SER	-	expression tag	UNP P40926
B	3	HIS	-	expression tag	UNP P40926
B	4	HIS	-	expression tag	UNP P40926
B	5	HIS	-	expression tag	UNP P40926
B	6	HIS	-	expression tag	UNP P40926
B	7	HIS	-	expression tag	UNP P40926
B	8	HIS	-	expression tag	UNP P40926
B	9	SER	-	expression tag	UNP P40926
B	10	SER	-	expression tag	UNP P40926
B	11	GLY	-	expression tag	UNP P40926
B	12	LEU	-	expression tag	UNP P40926
B	13	VAL	-	expression tag	UNP P40926
B	14	PRO	-	expression tag	UNP P40926
B	15	ARG	-	expression tag	UNP P40926
B	16	GLY	-	expression tag	UNP P40926
B	17	SER	-	expression tag	UNP P40926
B	18	HIS	-	expression tag	UNP P40926
B	19	MET	-	expression tag	UNP P40926
C	-1	MET	-	expression tag	UNP P40926
C	0	GLY	-	expression tag	UNP P40926
C	1	SER	-	expression tag	UNP P40926
C	2	SER	-	expression tag	UNP P40926
C	3	HIS	-	expression tag	UNP P40926
C	4	HIS	-	expression tag	UNP P40926
C	5	HIS	-	expression tag	UNP P40926
C	6	HIS	-	expression tag	UNP P40926
C	7	HIS	-	expression tag	UNP P40926
C	8	HIS	-	expression tag	UNP P40926
C	9	SER	-	expression tag	UNP P40926
C	10	SER	-	expression tag	UNP P40926
C	11	GLY	-	expression tag	UNP P40926
C	12	LEU	-	expression tag	UNP P40926
C	13	VAL	-	expression tag	UNP P40926
C	14	PRO	-	expression tag	UNP P40926
C	15	ARG	-	expression tag	UNP P40926
C	16	GLY	-	expression tag	UNP P40926
C	17	SER	-	expression tag	UNP P40926
C	18	HIS	-	expression tag	UNP P40926
C	19	MET	-	expression tag	UNP P40926

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	MET	-	expression tag	UNP P40926
D	0	GLY	-	expression tag	UNP P40926
D	1	SER	-	expression tag	UNP P40926
D	2	SER	-	expression tag	UNP P40926
D	3	HIS	-	expression tag	UNP P40926
D	4	HIS	-	expression tag	UNP P40926
D	5	HIS	-	expression tag	UNP P40926
D	6	HIS	-	expression tag	UNP P40926
D	7	HIS	-	expression tag	UNP P40926
D	8	HIS	-	expression tag	UNP P40926
D	9	SER	-	expression tag	UNP P40926
D	10	SER	-	expression tag	UNP P40926
D	11	GLY	-	expression tag	UNP P40926
D	12	LEU	-	expression tag	UNP P40926
D	13	VAL	-	expression tag	UNP P40926
D	14	PRO	-	expression tag	UNP P40926
D	15	ARG	-	expression tag	UNP P40926
D	16	GLY	-	expression tag	UNP P40926
D	17	SER	-	expression tag	UNP P40926
D	18	HIS	-	expression tag	UNP P40926
D	19	MET	-	expression tag	UNP P40926

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
			Total	O	P		
2	A	1	5	4	1	0	0
			Total	O	P		
2	B	1	5	4	1	0	0
			Total	O	P		
2	B	1	5	4	1	0	0
			Total	O	P		
2	B	1	5	4	1	0	0
			Total	O	P		
2	C	1	5	4	1	0	0
			Total	O	P		
2	C	1	5	4	1	0	0
			Total	O	P		
2	C	1	5	4	1	0	0
			Total	O	P		
2	D	1	5	4	1	0	0
			Total	O	P		
2	D	1	5	4	1	0	0
			Total	O	P		
2	D	1	5	4	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	42	42	42	0	0
			Total	O		
3	B	32	32	32	0	0
			Total	O		
3	C	18	18	18	0	0
			Total	O		
3	D	30	30	30	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Malate dehydrogenase, mitochondrial

Chain A:

• Molecule 1: Malate dehydrogenase, mitochondrial

Chain B:





• Molecule 1: Malate dehydrogenase, mitochondrial

Chain C:





• Molecule 1: Malate dehydrogenase, mitochondrial

Chain D:




V216
D217
L226
E232
K269
E270
G271
S276
Q281
E284
C285
T286
K297
E313
I320
P321
E322
L337
LYS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	59.92Å 151.27Å 155.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.56 – 2.28 32.56 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.8 (32.56-2.28) 99.9 (32.56-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.26 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.217 , 0.246 0.221 , 0.248	Depositor DCC
R_{free} test set	3322 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	31.4	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.42 , 45.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.008 for -h,l,k	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	9430	wwPDB-VP
Average B, all atoms (Å^2)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.29	0/2351	0.47	0/3192
1	B	0.28	0/2351	0.47	0/3192
1	C	0.28	0/2351	0.46	0/3192
1	D	0.28	0/2351	0.46	0/3192
All	All	0.28	0/9404	0.47	0/12768

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2312	0	2400	4	0
1	B	2312	0	2400	5	0
1	C	2312	0	2400	6	0
1	D	2312	0	2400	7	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
2	C	15	0	0	0	0
2	D	15	0	0	0	0
3	A	42	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	32	0	0	0	0
3	C	18	0	0	0	0
3	D	30	0	0	0	0
All	All	9430	0	9600	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:116:THR:HG23	1:C:106:PRO:O	2.00	0.62
1:D:48:PRO:O	1:D:74:LYS:NZ	2.35	0.58
1:A:106:PRO:O	1:D:116:THR:HG23	2.06	0.56
1:D:28:ALA:HB2	1:D:93:CYS:SG	2.50	0.52
1:A:28:ALA:HB2	1:A:93:CYS:SG	2.53	0.49
1:C:286:THR:HG23	1:C:327:ILE:HG21	1.95	0.47
1:B:115:ASN:O	1:B:119:THR:CG2	2.63	0.46
1:C:136:MET:HE3	1:C:167:PHE:HE2	1.81	0.45
1:C:28:ALA:HB2	1:C:93:CYS:SG	2.56	0.45
1:C:136:MET:CE	1:C:167:PHE:HE2	2.30	0.45
1:B:28:ALA:HB2	1:B:93:CYS:SG	2.56	0.45
1:A:266:MET:HA	1:A:296:LYS:HA	1.99	0.45
1:D:320:ILE:HB	1:D:321:PRO:HD3	1.99	0.45
1:C:86:LEU:N	1:C:87:PRO:HD2	2.32	0.44
1:D:203:LYS:NZ	1:D:322:GLU:OE1	2.50	0.44
1:B:148:ILE:HD13	1:B:148:ILE:HA	1.92	0.43
1:D:115:ASN:O	1:D:119:THR:HG23	2.19	0.42
1:B:170:THR:HG22	1:B:200:HIS:HB2	2.02	0.41
1:A:286:THR:HG23	1:A:327:ILE:HG21	2.02	0.41
1:D:181:VAL:HG22	1:D:226:LEU:HD21	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/340 (92%)	304 (97%)	8 (3%)	0	100	100
1	B	312/340 (92%)	309 (99%)	3 (1%)	0	100	100
1	C	312/340 (92%)	304 (97%)	8 (3%)	0	100	100
1	D	312/340 (92%)	307 (98%)	5 (2%)	0	100	100
All	All	1248/1360 (92%)	1224 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/275 (92%)	247 (98%)	6 (2%)	49	63
1	B	253/275 (92%)	244 (96%)	9 (4%)	35	47
1	C	253/275 (92%)	245 (97%)	8 (3%)	39	52
1	D	253/275 (92%)	246 (97%)	7 (3%)	43	57
All	All	1012/1100 (92%)	982 (97%)	30 (3%)	41	54

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	52	ARG
1	A	61	THR
1	A	119	THR
1	A	286	THR
1	A	314	LYS
1	B	52	ARG
1	B	61	THR
1	B	111	ASP
1	B	116	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	119	THR
1	B	220	GLN
1	B	232	GLU
1	B	281	GLN
1	B	307	LYS
1	C	24	ASN
1	C	61	THR
1	C	97	VAL
1	C	119	THR
1	C	220	GLN
1	C	286	THR
1	C	314	LYS
1	C	328	LYS
1	D	61	THR
1	D	97	VAL
1	D	116	THR
1	D	215	LYS
1	D	232	GLU
1	D	281	GLN
1	D	286	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	164	ASN
1	A	267	ASN
1	B	24	ASN
1	B	164	ASN
1	B	267	ASN
1	B	281	GLN
1	C	24	ASN
1	C	164	ASN
1	D	24	ASN
1	D	164	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	PO4	B	402	-	4,4,4	0.92	0	6,6,6	0.51	0
2	PO4	D	403	-	4,4,4	0.88	0	6,6,6	0.46	0
2	PO4	D	402	-	4,4,4	0.97	0	6,6,6	0.49	0
2	PO4	A	403	-	4,4,4	0.83	0	6,6,6	0.61	0
2	PO4	B	401	-	4,4,4	1.01	0	6,6,6	0.62	0
2	PO4	C	402	-	4,4,4	0.96	0	6,6,6	0.55	0
2	PO4	B	403	-	4,4,4	0.97	0	6,6,6	0.58	0
2	PO4	C	401	-	4,4,4	0.93	0	6,6,6	0.63	0
2	PO4	C	403	-	4,4,4	0.96	0	6,6,6	0.54	0
2	PO4	A	402	-	4,4,4	1.02	0	6,6,6	0.49	0
2	PO4	D	401	-	4,4,4	1.02	0	6,6,6	0.64	0
2	PO4	A	401	-	4,4,4	0.90	0	6,6,6	0.35	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	314/340 (92%)	0.02	5 (1%)	72	77	19, 27, 39, 58	0
1	B	314/340 (92%)	0.27	15 (4%)	30	36	19, 32, 50, 61	0
1	C	314/340 (92%)	0.33	19 (6%)	21	26	22, 34, 51, 68	0
1	D	314/340 (92%)	0.26	16 (5%)	28	33	22, 34, 50, 67	0
All	All	1256/1360 (92%)	0.22	55 (4%)	34	40	19, 32, 49, 68	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	LYS	4.6
1	D	169	VAL	3.8
1	D	284	GLU	3.7
1	C	220	GLN	3.7
1	C	169	VAL	3.5
1	D	271	GLY	3.4
1	D	139	VAL	3.4
1	D	313	GLU	3.3
1	D	281	GLN	3.1
1	C	191	ARG	3.1
1	B	217	ASP	3.1
1	B	284	GLU	3.1
1	D	270	GLU	3.0
1	C	215	LYS	3.0
1	C	335	LYS	2.9
1	C	221	ASP	2.9
1	A	169	VAL	2.8
1	D	170	THR	2.8
1	D	276	SER	2.8
1	C	336	THR	2.8
1	C	98	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	170	THR	2.7
1	C	139	VAL	2.7
1	B	337	LEU	2.6
1	B	191	ARG	2.6
1	B	169	VAL	2.6
1	D	140	ILE	2.6
1	C	24	ASN	2.5
1	D	217	ASP	2.5
1	D	161	TYR	2.4
1	C	170	THR	2.4
1	D	269	LYS	2.4
1	A	311	PHE	2.3
1	B	196	VAL	2.3
1	B	171	THR	2.3
1	C	281	GLN	2.3
1	B	281	GLN	2.3
1	C	171	THR	2.2
1	C	337	LEU	2.2
1	C	225	ALA	2.2
1	D	191	ARG	2.2
1	B	221	ASP	2.2
1	C	137	ILE	2.2
1	B	307	LYS	2.2
1	A	24	ASN	2.1
1	C	214	PRO	2.1
1	B	268	GLY	2.1
1	B	314	LYS	2.1
1	C	172	LEU	2.1
1	A	52	ARG	2.1
1	B	270	GLU	2.1
1	D	98	ILE	2.1
1	D	297	LYS	2.1
1	B	167	PHE	2.0
1	C	217	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	PO4	D	403	5/5	0.95	0.14	31,32,33,34	0
2	PO4	B	402	5/5	0.96	0.13	40,41,43,44	0
2	PO4	B	403	5/5	0.97	0.12	32,32,34,35	0
2	PO4	C	403	5/5	0.97	0.13	42,42,44,45	0
2	PO4	D	401	5/5	0.97	0.14	37,37,40,40	0
2	PO4	A	402	5/5	0.97	0.13	42,44,45,45	0
2	PO4	B	401	5/5	0.98	0.14	32,33,34,35	0
2	PO4	C	402	5/5	0.99	0.08	30,31,31,32	0
2	PO4	A	403	5/5	0.99	0.09	22,22,22,23	0
2	PO4	A	401	5/5	0.99	0.09	26,26,27,28	0
2	PO4	D	402	5/5	0.99	0.07	25,25,26,26	0
2	PO4	C	401	5/5	0.99	0.09	31,31,32,33	0

6.5 Other polymers (i)

There are no such residues in this entry.