wwPDB NMR Structure Validation Summary Report (i)

Feb 23, 2022 – 08:19 AM EST

PDB ID	:	1X4Q
Title	:	Solution structure of PWI domain in U4/U6 small nuclear ribonucleoprotein Prp3(hPrp3)
Authors	:	He, F.; Muto, Y.; Inoue, M.; Kigawa, T.; Shirouzu, M.; Terada, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2005-05-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	92	33% 37% • 26%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:10-A:77 (68)	0.15	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 4, 6, 10, 13, 14, 19, 20
2	2, 5, 7, 8, 11, 12, 16
3	3, 15
4	17, 18
Single-model clusters	9

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1394 atoms, of which 701 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called U4/U6 small nuclear ribonucleoprotein Prp3.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	92	1394	431	701	123	136	3	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP O43395
A	2	SER	-	cloning artifact	UNP O43395
A	3	SER	-	cloning artifact	UNP O43395
A	4	GLY	-	cloning artifact	UNP O43395
A	5	SER	-	cloning artifact	UNP O43395
A	6	SER	-	cloning artifact	UNP O43395
A	7	GLY	-	cloning artifact	UNP O43395
A	87	SER	-	cloning artifact	UNP O43395
A	88	GLY	-	cloning artifact	UNP O43395
A	89	PRO	-	cloning artifact	UNP O43395
A	90	SER	-	cloning artifact	UNP O43395
A	91	SER	-	cloning artifact	UNP O43395
A	92	GLY	-	cloning artifact	UNP O43395

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: U4/U6 small nuclear ribonucleoprotein Prp3

Chain A: 33% 37% • 26%

G1 S2 S3 G4 S5 S6 G7 M8 A9 K12 D16 K19 P20 W21 I22 E23 K24 T25 V26 K27 R28 V29 L30 G31 F32 S33 E34 V37 V38 K47 G48 M49 D50 K51 K52 K53 A54 L58 K59 P60 F61 L62 D63 D64 S65 T66 F69 V70 D71 K72 L73 F74 E75

A76 V77 E78 E79 G80 R81 S82 S83 R84 H85 S86 S87 G88 P89 S90 S91 G92

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

- Molecule 1: U4/U6 small nuclear ribonucleoprotein Prp3

Chain A: 40% 29% • 26%

G1 S2 S3 G4 S5 S6 G7 M8 A9 K19 P20 W21 I22 E23 K24 T25 V26 L30 G31 F32 S33 E34 V37 V38 T39 A40 C44 K47 G48 M49 D50 K51 K52 K53 A54 L58 K59 P60 F61 L62 D63 D64 S65 T66 V70 L73 F74 E75 A76 V77 E78 E79

G80 R81 S82 S83 R84 H85 S86 S87 G88 P89 S90 S91 G92

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.29
CYANA	refinement	2.0.29

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	539	567	567	28±4
All	All	10780	11340	11340	550

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

5 of 83 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:LEU:HD21	1:A:62:LEU:HD22	0.92	1.40	18	20
1:A:66:THR:O	1:A:70:VAL:HG12	0.88	1.68	8	20
1:A:22:ILE:O	1:A:26:VAL:HG23	0.86	1.70	14	14
1:A:21:TRP:CH2	1:A:76:ALA:HB2	0.85	2.05	14	18
1:A:21:TRP:CZ3	1:A:76:ALA:HB2	0.74	2.17	14	16

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	68/92 (74%)	53±2 (78±3%)	12±2 (18±3%)	3±1 (4±1%)	6 34
All	All	1360/1840 (74%)	1064 (78%)	245 (18%)	51 (4%)	5 33

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	63	ASP	20
1	A	33	SER	14
1	A	49	MET	13
1	A	70	VAL	2
1	A	51	LYS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	60/77 (78%)	50±2 (83±3%)	10±2 (17±3%)	4 39
All	All	1200/1540 (78%)	991 (83%)	209 (17%)	4 39

5 of 26 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	63	ASP	19
1	A	47	LYS	18
1	A	59	LYS	17
1	A	33	SER	15
1	A	12	LYS	14

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided