



wwPDB EM Validation Summary Report ⓘ

Mar 20, 2026 – 02:20 AM UTC

PDB ID : 9XCF / pdb_00009xcf
EMDB ID : EMD-66729
Title : Cryo-EM structure of csy3 with crRNA
Authors : Gao, X.; Cui, S.; Zhu, H.; Zhu, K.
Deposited on : 2025-10-25
Resolution : 4.58 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	2398 (4.08 - 5.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	
1	C	337	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	337	
1	E	337	
1	F	337	
1	G	337	
1	H	337	
1	I	337	
1	J	337	
1	K	337	
1	L	337	
1	N	337	
1	O	337	
1	P	337	
1	Q	337	
1	S	337	
1	T	337	
1	U	337	
1	V	337	
1	W	337	
1	Y	337	
1	Z	337	
1	a	337	
1	b	337	
1	f	337	
1	i	337	
1	o	337	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	p	337	9% 70% 28%
1	q	337	11% 74% 20% 6%
2	M	185	19% 81%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 155526 atoms, of which 75360 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR-associated protein Csy3.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	325	Total	C	H	N	O	S	0	0
			5001	1587	2487	439	485	3		
1	C	330	Total	C	H	N	O	S	0	0
			5091	1614	2532	448	493	4		
1	D	331	Total	C	H	N	O	S	0	0
			5097	1618	2531	449	495	4		
1	E	332	Total	C	H	N	O	S	0	0
			5119	1622	2546	450	497	4		
1	F	332	Total	C	H	N	O	S	0	0
			5119	1622	2546	450	497	4		
1	G	331	Total	C	H	N	O	S	0	0
			5105	1618	2539	449	495	4		
1	I	331	Total	C	H	N	O	S	0	0
			5105	1618	2539	449	495	4		
1	J	331	Total	C	H	N	O	S	0	0
			5105	1618	2539	449	495	4		
1	K	330	Total	C	H	N	O	S	0	0
			5091	1614	2532	448	493	4		
1	L	317	Total	C	H	N	O	S	0	0
			4896	1553	2433	433	473	4		
1	B	331	Total	C	H	N	O	S	0	0
			5094	1616	2530	449	495	4		
1	H	328	Total	C	H	N	O	S	0	0
			5050	1602	2511	442	491	4		
1	N	331	Total	C	H	N	O	S	0	0
			5105	1618	2539	449	495	4		
1	O	330	Total	C	H	N	O	S	0	0
			5081	1612	2526	447	492	4		
1	P	317	Total	C	H	N	O	S	0	0
			4896	1553	2433	433	473	4		
1	Q	323	Total	C	H	N	O	S	0	0
			4960	1576	2466	433	482	3		
1	S	330	Total	C	H	N	O	S	0	0
			5091	1614	2532	448	493	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
1	T	331	Total 5097	C 1618	H 2531	N 449	O 495	S 4	0	0
1	U	332	Total 5119	C 1622	H 2546	N 450	O 497	S 4	0	0
1	V	332	Total 5119	C 1622	H 2546	N 450	O 497	S 4	0	0
1	W	331	Total 5105	C 1618	H 2539	N 449	O 495	S 4	0	0
1	Y	331	Total 5105	C 1618	H 2539	N 449	O 495	S 4	0	0
1	Z	331	Total 5105	C 1618	H 2539	N 449	O 495	S 4	0	0
1	a	330	Total 5091	C 1614	H 2532	N 448	O 493	S 4	0	0
1	b	317	Total 4896	C 1553	H 2433	N 433	O 473	S 4	0	0
1	f	317	Total 4896	C 1553	H 2433	N 433	O 473	S 4	0	0
1	i	322	Total 4940	C 1567	H 2457	N 432	O 481	S 3	0	0
1	o	331	Total 5105	C 1618	H 2539	N 449	O 495	S 4	0	0
1	p	330	Total 5091	C 1614	H 2532	N 448	O 493	S 4	0	0
1	q	317	Total 4896	C 1553	H 2433	N 433	O 473	S 4	0	0


- Molecule 2 is a RNA chain called RNA (185-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	M	185	Total	C	N	O	P	0	0
			3955	1771	731	1268	185		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3






• Molecule 1: CRISPR-associated protein Csy3

Chain G: 61% 37%




• Molecule 1: CRISPR-associated protein Csy3


Chain I: 6% 63% 34%




• Molecule 1: CRISPR-associated protein Csy3

Chain J: 6% 58% 39%







• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3







• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3



• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3






• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3







• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3



• Molecule 1: CRISPR-associated protein Csy3






• Molecule 1: CRISPR-associated protein Csy3







• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3







• Molecule 1: CRISPR-associated protein Csy3




• Molecule 1: CRISPR-associated protein Csy3



• Molecule 1: CRISPR-associated protein Csy3



● Molecule 2: RNA (185-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	168817	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.566	Depositor
Minimum map value	-0.003	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.027	Depositor
Recommended contour level	0.005	Depositor
Map size (Å)	480.0, 480.0, 480.0	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0, 1.0, 1.0	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	2/2559 (0.1%)	0.74	0/3469
1	B	0.82	5/2611 (0.2%)	0.77	0/3539
1	C	0.80	2/2606 (0.1%)	0.75	1/3532 (0.0%)
1	D	0.79	1/2613 (0.0%)	0.72	0/3542
1	E	0.82	2/2620 (0.1%)	0.72	1/3552 (0.0%)
1	F	0.87	3/2620 (0.1%)	0.76	1/3552 (0.0%)
1	G	0.84	2/2613 (0.1%)	0.73	0/3542
1	H	0.84	0/2584	0.81	2/3501 (0.1%)
1	I	0.75	0/2613	0.73	0/3542
1	J	0.72	2/2613 (0.1%)	0.72	0/3542
1	K	0.69	1/2606 (0.0%)	0.70	0/3532
1	L	0.66	0/2509	0.69	0/3404
1	N	0.82	1/2613 (0.0%)	0.75	0/3542
1	O	0.74	0/2602	0.72	1/3527 (0.0%)
1	P	0.79	0/2509	0.70	0/3404
1	Q	0.79	3/2539 (0.1%)	0.75	1/3443 (0.0%)
1	S	0.60	0/2606	0.63	0/3532
1	T	0.64	0/2613	0.70	1/3542 (0.0%)
1	U	0.71	0/2620	0.74	1/3552 (0.0%)
1	V	0.78	1/2620 (0.0%)	0.74	0/3552
1	W	0.82	1/2613 (0.0%)	0.79	0/3542
1	Y	0.74	1/2613 (0.0%)	0.69	0/3542
1	Z	0.74	0/2613	0.71	0/3542
1	a	0.78	0/2606	0.75	1/3532 (0.0%)
1	b	0.74	1/2509 (0.0%)	0.76	2/3404 (0.1%)
1	f	0.58	0/2509	0.62	0/3404
1	i	0.72	1/2527 (0.0%)	0.74	2/3427 (0.1%)
1	o	0.67	1/2613 (0.0%)	0.66	1/3542 (0.0%)
1	p	0.56	0/2606	0.64	0/3532
1	q	0.53	0/2509	0.60	0/3404
2	M	0.78	0/4430	0.77	9/6893 (0.1%)
All	All	0.75	30/82037 (0.0%)	0.72	24/112107 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	5
1	C	0	1
1	D	0	5
1	E	0	2
1	F	0	5
1	G	0	4
1	H	0	3
1	I	0	5
1	J	0	2
1	K	0	4
1	L	0	2
1	N	0	2
1	O	0	3
1	P	0	2
1	Q	0	2
1	S	0	2
1	T	0	3
1	U	0	2
1	V	0	5
1	W	0	7
1	Y	0	2
1	Z	0	4
1	a	0	4
1	b	0	3
1	f	0	1
1	i	0	1
1	o	0	4
1	p	0	3
All	All	0	93

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	83	ALA	C-N	-8.47	1.21	1.33
1	C	83	ALA	C-N	-8.16	1.21	1.33
1	F	149	TRP	CA-CB	-7.45	1.40	1.53
1	o	183	SER	C-N	-7.29	1.22	1.33
1	V	149	TRP	CA-CB	-7.05	1.41	1.53

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	31	C	OP1-P-O3'	-9.66	79.03	108.00
2	M	55	C	OP1-P-O3'	-9.49	79.52	108.00
2	M	145	C	OP1-P-O3'	-9.42	79.74	108.00
2	M	145	C	OP2-P-O3'	-9.42	79.74	108.00
2	M	115	C	OP1-P-O3'	-9.16	80.52	108.00

There are no chirality outliers.

5 of 93 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	122	ILE	Peptide
1	A	148	LEU	Peptide
1	A	152	ARG	Peptide
1	A	168	ARG	Peptide
1	A	44	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2514	2487	2485	96	0
1	B	2564	2530	2531	104	0
1	C	2559	2532	2531	97	0
1	D	2566	2531	2538	104	0
1	E	2573	2546	2545	98	0
1	F	2573	2546	2545	93	0
1	G	2566	2539	2538	104	0
1	H	2539	2511	2509	110	0
1	I	2566	2539	2538	86	0
1	J	2566	2539	2538	106	0
1	K	2559	2532	2531	90	0
1	L	2463	2433	2432	81	0
1	N	2566	2539	2538	104	0
1	O	2555	2526	2525	85	0
1	P	2463	2433	2432	85	0
1	Q	2494	2466	2464	96	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2559	2532	2531	88	0
1	T	2566	2531	2538	75	0
1	U	2573	2546	2545	107	0
1	V	2573	2546	2545	102	0
1	W	2566	2539	2538	104	0
1	Y	2566	2539	2538	96	0
1	Z	2566	2539	2538	108	0
1	a	2559	2532	2531	95	0
1	b	2463	2433	2432	96	0
1	f	2463	2433	2432	72	0
1	i	2483	2457	2455	98	0
1	o	2566	2539	2538	81	0
1	p	2559	2532	2531	74	0
1	q	2463	2433	2432	49	0
2	M	3955	0	2000	627	0
All	All	80166	75360	77344	3007	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 3007 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:ARG:NH1	2:M:56:G:O5'	2.06	0.87
2:M:93:A:O2'	2:M:94:C:O4'	1.92	0.87
1:L:328:ARG:NH2	2:M:27:A:O2'	2.07	0.87
1:F:150:ARG:NH1	2:M:68:G:O5'	2.07	0.87
1:b:26:LEU:HD23	1:b:250:ALA:HB2	1.55	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/337 (95%)	279 (87%)	42 (13%)	0	100	100
1	B	329/337 (98%)	276 (84%)	52 (16%)	1 (0%)	36	71
1	C	328/337 (97%)	290 (88%)	38 (12%)	0	100	100
1	D	329/337 (98%)	290 (88%)	39 (12%)	0	100	100
1	E	330/337 (98%)	292 (88%)	37 (11%)	1 (0%)	36	71
1	F	330/337 (98%)	282 (86%)	47 (14%)	1 (0%)	36	71
1	G	329/337 (98%)	284 (86%)	45 (14%)	0	100	100
1	H	324/337 (96%)	264 (82%)	60 (18%)	0	100	100
1	I	329/337 (98%)	288 (88%)	41 (12%)	0	100	100
1	J	329/337 (98%)	284 (86%)	45 (14%)	0	100	100
1	K	328/337 (97%)	286 (87%)	42 (13%)	0	100	100
1	L	315/337 (94%)	266 (84%)	49 (16%)	0	100	100
1	N	329/337 (98%)	289 (88%)	39 (12%)	1 (0%)	36	71
1	O	328/337 (97%)	295 (90%)	33 (10%)	0	100	100
1	P	315/337 (94%)	276 (88%)	39 (12%)	0	100	100
1	Q	319/337 (95%)	286 (90%)	33 (10%)	0	100	100
1	S	328/337 (97%)	290 (88%)	38 (12%)	0	100	100
1	T	329/337 (98%)	289 (88%)	40 (12%)	0	100	100
1	U	330/337 (98%)	285 (86%)	45 (14%)	0	100	100
1	V	330/337 (98%)	283 (86%)	47 (14%)	0	100	100
1	W	329/337 (98%)	283 (86%)	46 (14%)	0	100	100
1	Y	329/337 (98%)	289 (88%)	40 (12%)	0	100	100
1	Z	329/337 (98%)	285 (87%)	44 (13%)	0	100	100
1	a	328/337 (97%)	294 (90%)	34 (10%)	0	100	100
1	b	315/337 (94%)	269 (85%)	46 (15%)	0	100	100
1	f	315/337 (94%)	281 (89%)	34 (11%)	0	100	100
1	i	318/337 (94%)	276 (87%)	42 (13%)	0	100	100
1	o	329/337 (98%)	295 (90%)	34 (10%)	0	100	100
1	p	328/337 (97%)	298 (91%)	30 (9%)	0	100	100
1	q	315/337 (94%)	286 (91%)	29 (9%)	0	100	100
All	All	9764/10110 (97%)	8530 (87%)	1230 (13%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	149	TRP
1	B	81	VAL
1	E	149	TRP
1	N	246	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/266 (97%)	257 (100%)	0	100	100
1	B	262/266 (98%)	261 (100%)	1 (0%)	84	82
1	C	262/266 (98%)	262 (100%)	0	100	100
1	D	263/266 (99%)	263 (100%)	0	100	100
1	E	264/266 (99%)	264 (100%)	0	100	100
1	F	264/266 (99%)	263 (100%)	1 (0%)	84	82
1	G	263/266 (99%)	263 (100%)	0	100	100
1	H	260/266 (98%)	260 (100%)	0	100	100
1	I	263/266 (99%)	263 (100%)	0	100	100
1	J	263/266 (99%)	262 (100%)	1 (0%)	84	82
1	K	262/266 (98%)	262 (100%)	0	100	100
1	L	252/266 (95%)	252 (100%)	0	100	100
1	N	263/266 (99%)	263 (100%)	0	100	100
1	O	261/266 (98%)	261 (100%)	0	100	100
1	P	252/266 (95%)	252 (100%)	0	100	100
1	Q	255/266 (96%)	255 (100%)	0	100	100
1	S	262/266 (98%)	262 (100%)	0	100	100
1	T	263/266 (99%)	262 (100%)	1 (0%)	84	82
1	U	264/266 (99%)	263 (100%)	1 (0%)	84	82
1	V	264/266 (99%)	263 (100%)	1 (0%)	84	82
1	W	263/266 (99%)	263 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Y	263/266 (99%)	263 (100%)	0	100	100
1	Z	263/266 (99%)	263 (100%)	0	100	100
1	a	262/266 (98%)	262 (100%)	0	100	100
1	b	252/266 (95%)	252 (100%)	0	100	100
1	f	252/266 (95%)	252 (100%)	0	100	100
1	i	254/266 (96%)	254 (100%)	0	100	100
1	o	263/266 (99%)	263 (100%)	0	100	100
1	p	262/266 (98%)	262 (100%)	0	100	100
1	q	252/266 (95%)	252 (100%)	0	100	100
All	All	7805/7980 (98%)	7799 (100%)	6 (0%)	87	88

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	T	28	TYR
1	U	85	PRO
1	V	28	TYR
1	J	46	LYS
1	F	118	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 91 such sidechains are listed below:

Mol	Chain	Res	Type
1	S	254	GLN
1	Y	72	GLN
1	U	112	GLN
1	W	37	ASN
1	a	141	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	M	184/185 (99%)	147 (79%)	12 (6%)

5 of 147 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	M	4	G
2	M	5	A
2	M	6	U
2	M	7	U
2	M	8	C

5 of 12 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	M	134	G
2	M	135	A
2	M	177	G
2	M	136	A
2	M	62	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-66729. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240


6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 221




Y Index: 265




Z Index: 306


6.3.2 Raw map



X Index: 265



Y Index: 266




Z Index: 237


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.005. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 35156 nm^3 ; this corresponds to an approximate mass of 31758 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.218 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.218 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.58	-	-
Author-provided FSC curve	4.58	5.72	4.68
Unmasked-calculated*	7.20	9.17	7.44

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.20 differs from the reported value 4.58 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-66729 and PDB model 9XCF. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.005 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.005).

9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.005) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8200	 0.1500
A	 0.8260	 0.1630
B	 0.8280	 0.1540
C	 0.8270	 0.1630
D	 0.8290	 0.1690
E	 0.8460	 0.1700
F	 0.8420	 0.1760
G	 0.8520	 0.1790
H	 0.8410	 0.1460
I	 0.8190	 0.1450
J	 0.8060	 0.1210
K	 0.7970	 0.1210
L	 0.8130	 0.1200
M	 0.9890	 0.3420
N	 0.8380	 0.1690
O	 0.8280	 0.1700
P	 0.8410	 0.1720
Q	 0.8290	 0.1480
S	 0.7830	 0.0880
T	 0.7940	 0.1060
U	 0.8070	 0.1100
V	 0.8170	 0.1330
W	 0.8390	 0.1470
Y	 0.8210	 0.1340
Z	 0.8210	 0.1420
a	 0.8340	 0.1580
b	 0.8230	 0.1380
f	 0.8200	 0.0810
i	 0.8260	 0.1320
o	 0.8150	 0.1360
p	 0.7890	 0.1130
q	 0.7880	 0.0940

