wwPDB EM Validation Summary Report (i)

Feb 21, 2023 – 06:27 PM JST

PDB ID	:	7XDB
EMDB ID	:	EMD-33142
Title	:	Cryo-EM structure of SARS-CoV-2 Omicron Spike protein in complex with BA7208 fab
Authors	:	Liu, Z.; Liu, S.; Gao, Y.Z.
Deposited on	:	2022-03-26
Resolution	:	2.62 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	106	• 66% 34%
1	F	106	14% 71% 29%
1	H	106	100% 69% 31%
2	A	1116	• 74% 19% 7%
2	B	1116	• 76% 18% 7%
2	C	1116	19% 71% 21% 7%
3	D	118	8% 66% 34%
3	G	118	7% 73% 27%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	118	100%, 73%, 27%
4	J	2	50%, 50%
4	K	2	100%
4	L	2	100%
4	M	2	50%, 50%
4	O	2	50%, 50%
4	S	2	100%
4	U	2	100%
4	Z	2	50%, 50%
4	d	2	100%
4	f	2	100%
4	i	2	50%, 50%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 30189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BA7208 fab.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	E	106	818	518	132	165	3	0	0
1	F	106	818	518	132	165	3	0	0
1	H	106	818	518	132	165	3	0	0

- Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	A	1034	8126	5208	1355	1527	36	0	0
2	C	1033	8121	5205	1354	1526	36	0	0
2	B	1040	8169	5236	1361	1536	36	0	0

There are 123 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	VAL	ALA	variant	UNP P0DTC2
A	?	-	HIS	deletion	UNP P0DTC2
A	?	-	VAL	deletion	UNP P0DTC2
A	96	ILE	THR	variant	UNP P0DTC2
A	143	ASP	GLY	variant	UNP P0DTC2
A	?	-	VAL	deletion	UNP P0DTC2
A	?	-	TYR	deletion	UNP P0DTC2
A	209	ILE	LEU	variant	UNP P0DTC2
A	212	GLU	-	insertion	UNP P0DTC2
A	213	PRO	-	insertion	UNP P0DTC2
A	214	GLU	-	insertion	UNP P0DTC2
A	339	ASP	GLY	variant	UNP P0DTC2
A	371	LEU	SER	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	373	PRO	SER	variant	UNP P0DTC2
A	375	PHE	SER	variant	UNP P0DTC2
A	417	ASN	LYS	variant	UNP P0DTC2
A	440	LYS	ASN	variant	UNP P0DTC2
A	446	SER	GLY	variant	UNP P0DTC2
A	477	ASN	SER	variant	UNP P0DTC2
A	478	LYS	THR	variant	UNP P0DTC2
A	484	ALA	GLU	variant	UNP P0DTC2
A	493	ARG	GLN	variant	UNP P0DTC2
A	496	SER	GLY	variant	UNP P0DTC2
A	498	ARG	GLN	variant	UNP P0DTC2
A	501	TYR	ASN	variant	UNP P0DTC2
A	505	HIS	TYR	variant	UNP P0DTC2
A	547	LYS	THR	variant	UNP P0DTC2
A	614	GLY	ASP	variant	UNP P0DTC2
A	655	TYR	HIS	variant	UNP P0DTC2
A	764	LYS	ASN	variant	UNP P0DTC2
A	796	TYR	ASP	variant	UNP P0DTC2
A	817	PRO	PHE	engineered mutation	UNP P0DTC2
A	856	LYS	ASN	variant	UNP P0DTC2
A	892	PRO	ALA	engineered mutation	UNP P0DTC2
A	899	PRO	ALA	engineered mutation	UNP P0DTC2
A	942	PRO	ALA	engineered mutation	UNP P0DTC2
A	954	HIS	GLN	variant	UNP P0DTC2
A	969	LYS	ASN	variant	UNP P0DTC2
A	981	PHE	LEU	variant	UNP P0DTC2
A	986	PRO	LYS	engineered mutation	UNP P0DTC2
A	987	PRO	VAL	engineered mutation	UNP P0DTC2
C	70	VAL	ALA	variant	UNP P0DTC2
C	?	-	HIS	deletion	UNP P0DTC2
C	?	-	VAL	deletion	UNP P0DTC2
C	96	ILE	THR	variant	UNP P0DTC2
C	143	ASP	GLY	variant	UNP P0DTC2
C	?	-	VAL	deletion	UNP P0DTC2
C	?	-	TYR	deletion	UNP P0DTC2
C	209	ILE	LEU	variant	UNP P0DTC2
C	212	GLU	-	insertion	UNP P0DTC2
C	213	PRO	-	insertion	UNP P0DTC2
C	214	GLU	-	insertion	UNP P0DTC2
C	339	ASP	GLY	variant	UNP P0DTC2
C	371	LEU	SER	variant	UNP P0DTC2
C	373	PRO	SER	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	375	PHE	SER	variant	UNP P0DTC2
C	417	ASN	LYS	variant	UNP P0DTC2
C	440	LYS	ASN	variant	UNP P0DTC2
C	446	SER	GLY	variant	UNP P0DTC2
C	477	ASN	SER	variant	UNP P0DTC2
C	478	LYS	THR	variant	UNP P0DTC2
C	484	ALA	GLU	variant	UNP P0DTC2
C	493	ARG	GLN	variant	UNP P0DTC2
C	496	SER	GLY	variant	UNP P0DTC2
C	498	ARG	GLN	variant	UNP P0DTC2
C	501	TYR	ASN	variant	UNP P0DTC2
C	505	HIS	TYR	variant	UNP P0DTC2
C	547	LYS	THR	variant	UNP P0DTC2
C	614	GLY	ASP	variant	UNP P0DTC2
C	655	TYR	HIS	variant	UNP P0DTC2
C	764	LYS	ASN	variant	UNP P0DTC2
C	796	TYR	ASP	variant	UNP P0DTC2
C	817	PRO	PHE	engineered mutation	UNP P0DTC2
C	856	LYS	ASN	variant	UNP P0DTC2
C	892	PRO	ALA	engineered mutation	UNP P0DTC2
C	899	PRO	ALA	engineered mutation	UNP P0DTC2
C	942	PRO	ALA	engineered mutation	UNP P0DTC2
C	954	HIS	GLN	variant	UNP P0DTC2
C	969	LYS	ASN	variant	UNP P0DTC2
C	981	PHE	LEU	variant	UNP P0DTC2
C	986	PRO	LYS	engineered mutation	UNP P0DTC2
C	987	PRO	VAL	engineered mutation	UNP P0DTC2
B	70	VAL	ALA	variant	UNP P0DTC2
B	?	-	HIS	deletion	UNP P0DTC2
B	?	-	VAL	deletion	UNP P0DTC2
B	96	ILE	THR	variant	UNP P0DTC2
B	143	ASP	GLY	variant	UNP P0DTC2
B	?	-	VAL	deletion	UNP P0DTC2
B	?	-	TYR	deletion	UNP P0DTC2
B	209	ILE	LEU	variant	UNP P0DTC2
B	212	GLU	-	insertion	UNP P0DTC2
B	213	PRO	-	insertion	UNP P0DTC2
B	214	GLU	-	insertion	UNP P0DTC2
B	339	ASP	GLY	variant	UNP P0DTC2
B	371	LEU	SER	variant	UNP P0DTC2
B	373	PRO	SER	variant	UNP P0DTC2
B	375	PHE	SER	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	417	ASN	LYS	variant	UNP P0DTC2
B	440	LYS	ASN	variant	UNP P0DTC2
B	446	SER	GLY	variant	UNP P0DTC2
B	477	ASN	SER	variant	UNP P0DTC2
B	478	LYS	THR	variant	UNP P0DTC2
B	484	ALA	GLU	variant	UNP P0DTC2
B	493	ARG	GLN	variant	UNP P0DTC2
B	496	SER	GLY	variant	UNP P0DTC2
B	498	ARG	GLN	variant	UNP P0DTC2
B	501	TYR	ASN	variant	UNP P0DTC2
B	505	HIS	TYR	variant	UNP P0DTC2
B	547	LYS	THR	variant	UNP P0DTC2
B	614	GLY	ASP	variant	UNP P0DTC2
B	655	TYR	HIS	variant	UNP P0DTC2
B	764	LYS	ASN	variant	UNP P0DTC2
B	796	TYR	ASP	variant	UNP P0DTC2
B	817	PRO	PHE	engineered mutation	UNP P0DTC2
B	856	LYS	ASN	variant	UNP P0DTC2
B	892	PRO	ALA	engineered mutation	UNP P0DTC2
B	899	PRO	ALA	engineered mutation	UNP P0DTC2
B	942	PRO	ALA	engineered mutation	UNP P0DTC2
B	954	HIS	GLN	variant	UNP P0DTC2
B	969	LYS	ASN	variant	UNP P0DTC2
B	981	PHE	LEU	variant	UNP P0DTC2
B	986	PRO	LYS	engineered mutation	UNP P0DTC2
B	987	PRO	VAL	engineered mutation	UNP P0DTC2

- Molecule 3 is a protein called BA7208 fab.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	D	118	929	598	151	175	5	0	0
			Total	C	N	O	S		
3	G	118	929	598	151	175	5	0	0
			Total	C	N	O	S		
3	I	118	929	598	151	175	5	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	O	2	28	16	2	10	0	0
4	S	2	28	16	2	10	0	0
4	U	2	28	16	2	10	0	0
4	Z	2	28	16	2	10	0	0
4	d	2	28	16	2	10	0	0
4	f	2	28	16	2	10	0	0
4	i	2	28	16	2	10	0	0
4	J	2	28	16	2	10	0	0
4	K	2	28	16	2	10	0	0
4	L	2	28	16	2	10	0	0
4	M	2	28	16	2	10	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	A	1	14	8	1	5	0
5	A	1	14	8	1	5	0
5	A	1	14	8	1	5	0
5	A	1	14	8	1	5	0
5	C	1	14	8	1	5	0
5	C	1	14	8	1	5	0
5	C	1	14	8	1	5	0
5	C	1	14	8	1	5	0
5	C	1	14	8	1	5	0
5	B	1	14	8	1	5	0
5	B	1	14	8	1	5	0
5	B	1	14	8	1	5	0
5	B	1	14	8	1	5	0
5	B	1	14	8	1	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	B	1	14	8	1	5	0
			Total	C	N	O	
5	B	1	14	8	1	5	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BA7208 fab

Chain E:



- Molecule 1: BA7208 fab

Chain F:



- Molecule 1: BA7208 fab

Chain H:

- Molecule 2: Spike glycoprotein

Chain A:



- Molecule 2: Spike glycoprotein

Chain C:





- Molecule 2: Spike glycoprotein

Chain B:





• Molecule 3: BA7208 fab

Chain D:

• Molecule 3: BA7208 fab

Chain G:



• Molecule 3: BA7208 fab

Chain I:





• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain U: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Z: 50% 50%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain d: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain f: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain i: 50% 50%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 50% 50%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 100%

NAG1
NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 100%

NAG1
NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 50% 50%

NAG1
NAG2

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	439731	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	1.39	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	4.347	Depositor
Minimum map value	-2.248	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.077	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	342.4, 342.4, 342.4	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.856, 0.856, 0.856	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	E	0.26	0/838	0.49	0/1138
1	F	0.26	0/838	0.51	0/1138
1	H	0.27	0/838	0.51	0/1138
2	A	0.27	0/8320	0.48	0/11319
2	B	0.27	0/8366	0.48	0/11386
2	C	0.28	0/8315	0.50	0/11312
3	D	0.25	0/958	0.48	0/1301
3	G	0.27	0/958	0.49	0/1301
3	I	0.27	0/958	0.49	0/1301
All	All	0.27	0/30389	0.49	0/41334

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	617	CYS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	818	0	790	22	0
1	F	818	0	790	23	0
1	H	818	0	790	24	0
2	A	8126	0	7954	156	0
2	B	8169	0	7992	143	0
2	C	8121	0	7952	179	0
3	D	929	0	885	34	0
3	G	929	0	885	22	0
3	I	929	0	885	22	0
4	J	28	0	25	1	0
4	K	28	0	25	0	0
4	L	28	0	25	0	0
4	M	28	0	25	2	0
4	O	28	0	25	1	0
4	S	28	0	25	0	0
4	U	28	0	25	0	0
4	Z	28	0	25	1	0
4	d	28	0	25	0	0
4	f	28	0	25	0	0
4	i	28	0	25	0	0
5	A	56	0	52	1	0
5	B	98	0	91	1	0
5	C	70	0	65	1	0
All	All	30189	0	29406	568	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 568 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:165:THR:HG23	2:C:357:ARG:NH2	1.71	1.04
2:C:328:ARG:NH2	2:C:580:GLN:HB3	1.93	0.83
2:A:329:PHE:O	2:A:580:GLN:NE2	2.11	0.83
2:A:903:ALA:HB1	2:A:913:GLN:HG2	1.63	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:332:ILE:HG22	2:C:362:VAL:HG11	1.63	0.79

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	104/106 (98%)	100 (96%)	4 (4%)	0	100	100
1	F	104/106 (98%)	101 (97%)	3 (3%)	0	100	100
1	H	104/106 (98%)	101 (97%)	3 (3%)	0	100	100
2	A	1016/1116 (91%)	996 (98%)	20 (2%)	0	100	100
2	B	1024/1116 (92%)	1010 (99%)	14 (1%)	0	100	100
2	C	1015/1116 (91%)	994 (98%)	21 (2%)	0	100	100
3	D	116/118 (98%)	114 (98%)	2 (2%)	0	100	100
3	G	116/118 (98%)	115 (99%)	1 (1%)	0	100	100
3	I	116/118 (98%)	115 (99%)	1 (1%)	0	100	100
All	All	3715/4020 (92%)	3646 (98%)	69 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	92/92 (100%)	92 (100%)	0	100	100
1	F	92/92 (100%)	92 (100%)	0	100	100
1	H	92/92 (100%)	92 (100%)	0	100	100
2	A	907/973 (93%)	907 (100%)	0	100	100
2	B	912/973 (94%)	912 (100%)	0	100	100
2	C	907/973 (93%)	907 (100%)	0	100	100
3	D	99/99 (100%)	99 (100%)	0	100	100
3	G	99/99 (100%)	99 (100%)	0	100	100
3	I	99/99 (100%)	99 (100%)	0	100	100
All	All	3299/3492 (94%)	3299 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	89	GLN
1	H	38	GLN
1	F	89	GLN
2	B	1002	GLN
1	F	38	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	J	1	2,4	14,14,15	0.26	0	17,19,21	0.51	0
4	NAG	J	2	4	14,14,15	0.23	0	17,19,21	0.39	0
4	NAG	K	1	2,4	14,14,15	0.21	0	17,19,21	0.49	0
4	NAG	K	2	4	14,14,15	0.21	0	17,19,21	0.41	0
4	NAG	L	1	2,4	14,14,15	0.27	0	17,19,21	0.53	0
4	NAG	L	2	4	14,14,15	0.23	0	17,19,21	0.38	0
4	NAG	M	1	2,4	14,14,15	0.27	0	17,19,21	0.57	0
4	NAG	M	2	4	14,14,15	0.22	0	17,19,21	0.46	0
4	NAG	O	1	2,4	14,14,15	0.28	0	17,19,21	0.49	0
4	NAG	O	2	4	14,14,15	0.24	0	17,19,21	0.38	0
4	NAG	S	1	2,4	14,14,15	0.25	0	17,19,21	0.49	0
4	NAG	S	2	4	14,14,15	0.23	0	17,19,21	0.38	0
4	NAG	U	1	2,4	14,14,15	0.24	0	17,19,21	0.54	0
4	NAG	U	2	4	14,14,15	0.21	0	17,19,21	0.38	0
4	NAG	Z	1	2,4	14,14,15	0.32	0	17,19,21	0.52	0
4	NAG	Z	2	4	14,14,15	0.26	0	17,19,21	0.39	0
4	NAG	d	1	2,4	14,14,15	0.27	0	17,19,21	0.50	0
4	NAG	d	2	4	14,14,15	0.21	0	17,19,21	0.39	0
4	NAG	f	1	2,4	14,14,15	0.23	0	17,19,21	0.48	0
4	NAG	f	2	4	14,14,15	0.21	0	17,19,21	0.41	0
4	NAG	i	1	2,4	14,14,15	1.08	1 (7%)	17,19,21	1.21	1 (5%)
4	NAG	i	2	4	14,14,15	0.24	0	17,19,21	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	J	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	1/6/23/26	0/1/1/1
4	NAG	K	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	2/6/23/26	0/1/1/1
4	NAG	L	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	L	2	4	-	2/6/23/26	0/1/1/1
4	NAG	M	1	2,4	-	1/6/23/26	0/1/1/1
4	NAG	M	2	4	-	2/6/23/26	0/1/1/1
4	NAG	O	1	2,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	O	2	4	-	0/6/23/26	0/1/1/1
4	NAG	S	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	S	2	4	-	2/6/23/26	0/1/1/1
4	NAG	U	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	U	2	4	-	2/6/23/26	0/1/1/1
4	NAG	Z	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	Z	2	4	-	2/6/23/26	0/1/1/1
4	NAG	d	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	d	2	4	-	2/6/23/26	0/1/1/1
4	NAG	f	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	f	2	4	-	2/6/23/26	0/1/1/1
4	NAG	i	1	2,4	-	3/6/23/26	0/1/1/1
4	NAG	i	2	4	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	i	1	NAG	O5-C1	-3.78	1.37	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	i	1	NAG	C3-C4-C5	3.20	115.95	110.24

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	i	1	NAG	C4-C5-C6-O6
4	S	2	NAG	O5-C5-C6-O6
4	Z	2	NAG	O5-C5-C6-O6
4	U	2	NAG	O5-C5-C6-O6
4	Z	2	NAG	C4-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	M	1	NAG	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	J	1	NAG	1	0
4	Z	1	NAG	1	0
4	O	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain O



Oligosaccharide Chain S	
 Bond lengths	 Bond angles
 Torsions	 Rings

Oligosaccharide Chain U



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain Z



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain d



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain f



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain i	
 Bond lengths	 Bond angles
 Torsions	 Rings

Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain K



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain L	
 Bond lengths	 Bond angles
 Torsions	 Rings

Oligosaccharide Chain M



5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	C	1202	2	14,14,15	0.36	0	17,19,21	0.42	0
5	NAG	C	1203	2	14,14,15	0.24	0	17,19,21	0.51	0
5	NAG	B	1204	2	14,14,15	0.22	0	17,19,21	0.32	0
5	NAG	A	1203	2	14,14,15	0.28	0	17,19,21	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	C	1201	2	14,14,15	0.29	0	17,19,21	0.34	0
5	NAG	B	1206	2	14,14,15	0.22	0	17,19,21	0.42	0
5	NAG	B	1202	2	14,14,15	0.31	0	17,19,21	0.36	0
5	NAG	C	1204	2	14,14,15	0.41	0	17,19,21	0.88	0
5	NAG	B	1207	2	14,14,15	0.22	0	17,19,21	0.44	0
5	NAG	A	1201	2	14,14,15	0.23	0	17,19,21	0.34	0
5	NAG	A	1204	2	14,14,15	0.25	0	17,19,21	0.50	0
5	NAG	B	1203	2	14,14,15	0.23	0	17,19,21	0.46	0
5	NAG	B	1205	2	14,14,15	0.26	0	17,19,21	0.49	0
5	NAG	B	1201	2	14,14,15	0.33	0	17,19,21	0.39	0
5	NAG	C	1205	2	14,14,15	0.41	0	17,19,21	0.62	0
5	NAG	A	1202	2	14,14,15	0.44	0	17,19,21	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1202	2	-	4/6/23/26	0/1/1/1
5	NAG	C	1203	2	-	2/6/23/26	0/1/1/1
5	NAG	B	1204	2	-	2/6/23/26	0/1/1/1
5	NAG	A	1203	2	-	2/6/23/26	0/1/1/1
5	NAG	C	1201	2	-	4/6/23/26	0/1/1/1
5	NAG	B	1206	2	-	2/6/23/26	0/1/1/1
5	NAG	B	1202	2	-	4/6/23/26	0/1/1/1
5	NAG	C	1204	2	-	3/6/23/26	0/1/1/1
5	NAG	B	1207	2	-	2/6/23/26	0/1/1/1
5	NAG	A	1201	2	-	4/6/23/26	0/1/1/1
5	NAG	A	1204	2	-	2/6/23/26	0/1/1/1
5	NAG	B	1203	2	-	2/6/23/26	0/1/1/1
5	NAG	B	1205	2	-	2/6/23/26	0/1/1/1
5	NAG	B	1201	2	-	4/6/23/26	0/1/1/1
5	NAG	C	1205	2	-	4/6/23/26	0/1/1/1
5	NAG	A	1202	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1204	NAG	C8-C7-N2-C2
5	C	1204	NAG	O7-C7-N2-C2
5	C	1205	NAG	O7-C7-N2-C2
5	A	1201	NAG	C4-C5-C6-O6
5	C	1205	NAG	C8-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1202	NAG	1	0
5	B	1202	NAG	1	0
5	A	1202	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NAG C 1202	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG C 1203	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1204	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG A 1203	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG C 1201	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1206	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1202	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG C 1204	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1207	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG A 1201	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG A 1204	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1203	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1205	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG B 1201	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG C 1205	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAG A 1202	
 Bond lengths	 Bond angles
 Torsions	 Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-33142. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 200


Y Index: 200


Z Index: 200

6.2.2 Raw map


X Index: 200


Y Index: 200


Z Index: 200

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 195



Y Index: 208



Z Index: 262

6.3.2 Raw map



X Index: 193



Y Index: 209



Z Index: 201

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 204 nm^3; this corresponds to an approximate mass of 184 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.382 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.382 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.62	-	-
Author-provided FSC curve	2.62	2.96	2.68
Unmasked-calculated*	3.17	3.72	3.24

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.17 differs from the reported value 2.62 by more than 10 %

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-33142 and PDB model 7XDB. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

9.4 Atom inclusion (i)



At the recommended contour level, 84% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7770	0.5020
A	0.8827	0.5680
B	0.8963	0.5770
C	0.7261	0.4680
D	0.7594	0.4830
E	0.7400	0.4670
F	0.7152	0.4550
G	0.7704	0.4980
H	0.0037	0.0360
I	0.0088	0.0570
J	0.8214	0.5370
K	0.7857	0.5000
L	0.8214	0.5510
M	0.7143	0.4370
O	0.8571	0.5030
S	0.8929	0.5710
U	0.7500	0.4850
Z	0.7857	0.5150
d	0.8929	0.5470
f	0.7857	0.4580
i	0.7500	0.4010