wwPDB X-ray Structure Validation Summary Report (i)

Aug 15, 2023 – 05:34 PM EDT

PDB ID	:	1XJM
Title	:	Structural mechanism of allosteric substrate specificity in a ribonucleotide reductase: dTTP complex
Authors	:	Larsson, K.-M.; Jordan, A.; Eliasson, R.; Reichard, P.; Logan, D.T.; Nordlund, P.
Deposited on	:	2004-09-23
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	644	4% 74% 18% • •
1	B	644	% 74% 17% • 6%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 10246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ribonucleotide reductase, B12-dependent.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	622	4990	3198	846	926	20	0	0	0
1	B	603	4838	3102	821	895	20	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	SER	TYR	SEE REMARK 999	UNP O33839
B	205	SER	TYR	SEE REMARK 999	UNP O33839

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
2	A	1	1	1	0	0
2	B	1	1	1	0	0

- Molecule 3 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: $C_{10}H_{17}N_2O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	29	10	2	14	3	0	0
			Total	C	N	O	P		
3	A	1	29	10	2	14	3	0	0
			Total	C	N	O	P		
3	B	1	29	10	2	14	3	0	0
			Total	C	N	O	P		
3	B	1	29	10	2	14	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	146	146	146	0	0
			Total	O		
4	B	154	154	154	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ribonucleotide reductase, B12-dependent

Chain A: 4% 74% 18%



• Molecule 1: ribonucleotide reductase, B12-dependent

Chain B: % 74% 17% 6%




D579
L582
D592
N593
S596
M601
P602
Q603
D608
D609
V610
R623
S632
LEU
GLN
THR
GLN
VAL
LEU
THR
LYS
ALA
LEU
LYS
THR

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	118.44Å 123.78Å 106.24Å 90.00° 103.64° 90.00°	Depositor
Resolution (Å)	19.98 – 2.40 19.98 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (19.98-2.40) 99.5 (19.98-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$< I/\sigma(I) >$ [1]	2.43 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.184 , 0.247 0.183 , 0.244	Depositor DCC
R_{free} test set	3813 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	31.8	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 40.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	10246	wwPDB-VP
Average B, all atoms (Å^2)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.79	0/5087	0.82	5/6866 (0.1%)
1	B	0.80	1/4933 (0.0%)	0.82	5/6658 (0.1%)
All	All	0.79	1/10020 (0.0%)	0.82	10/13524 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	464	ASN	CB-CG	5.09	1.62	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	623	ARG	NE-CZ-NH1	9.63	125.12	120.30
1	A	623	ARG	NE-CZ-NH2	-7.34	116.63	120.30
1	A	257	LEU	CA-CB-CG	5.73	128.47	115.30
1	B	243	LEU	CA-CB-CG	5.59	128.17	115.30
1	A	363	ARG	NE-CZ-NH1	5.33	122.96	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	518	LYS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4990	0	5046	94	0
1	B	4838	0	4890	84	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	58	0	26	1	0
3	B	58	0	26	3	0
4	A	146	0	0	4	1
4	B	154	0	0	4	0
All	All	10246	0	9988	175	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:519:GLU:HB2	1:A:520:ASP:HB2	1.23	1.20
1:B:449:TYR:HE1	1:B:473:MET:CE	1.68	1.05
1:A:519:GLU:CB	1:A:520:ASP:HB2	1.91	1.00
1:B:449:TYR:HE1	1:B:473:MET:HE1	1.37	0.89
1:A:62:TYR:CE2	1:A:70:ARG:HG2	2.07	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1145:HOH:O	4:A:1147:HOH:O[2_355]	1.73	0.47

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	614/644 (95%)	578 (94%)	29 (5%)	7 (1%)	14 20
1	B	595/644 (92%)	570 (96%)	22 (4%)	3 (0%)	29 41
All	All	1209/1288 (94%)	1148 (95%)	51 (4%)	10 (1%)	19 29

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	VAL
1	A	520	ASP
1	A	523	LYS
1	B	66	GLU
1	B	538	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	548/566 (97%)	512 (93%)	36 (7%)	16 26
1	B	531/566 (94%)	492 (93%)	39 (7%)	14 22
All	All	1079/1132 (95%)	1004 (93%)	75 (7%)	15 24

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	282	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	547	LYS
1	B	313	HIS
1	B	504	LEU
1	A	504	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	538	ASN
1	A	603	GLN
1	B	345	ASN
1	B	239	ASN
1	B	320	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	TTP	A	1004	-	26,30,30	2.17	7 (26%)	39,47,47	2.44	17 (43%)
3	TTP	A	1001	2	26,30,30	2.15	6 (23%)	39,47,47	2.38	15 (38%)
3	TTP	B	1002	2	26,30,30	2.09	7 (26%)	39,47,47	2.41	11 (28%)
3	TTP	B	1003	-	26,30,30	2.13	6 (23%)	39,47,47	2.64	15 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTP	A	1004	-	-	9/22/34/34	0/2/2/2
3	TTP	A	1001	2	-	2/22/34/34	0/2/2/2
3	TTP	B	1002	2	-	4/22/34/34	0/2/2/2
3	TTP	B	1003	-	-	1/22/34/34	0/2/2/2

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1003	TTP	O4-C4	7.14	1.37	1.23
3	A	1004	TTP	O4-C4	6.79	1.36	1.23
3	A	1001	TTP	O4-C4	6.51	1.36	1.23
3	B	1002	TTP	O4-C4	6.45	1.35	1.23
3	B	1003	TTP	C5M-C5	-4.48	1.39	1.50

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	1002	TTP	C5-C4-N3	7.91	122.06	115.31
3	A	1001	TTP	C5-C4-N3	7.64	121.83	115.31
3	A	1004	TTP	C5-C4-N3	7.12	121.39	115.31
3	B	1003	TTP	C5-C4-N3	6.44	120.81	115.31
3	B	1003	TTP	C2'-C1'-N1	5.26	125.89	113.77

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1004	TTP	C5'-O5'-PA-O1A
3	A	1004	TTP	C5'-O5'-PA-O2A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1004	TTP	C5'-O5'-PA-O3A
3	A	1004	TTP	PB-O3B-PG-O2G
3	A	1004	TTP	C3'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1004	TTP	1	0
3	B	1003	TTP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand TTP A 1004



Bond lengths



Bond angles



Torsions



Rings

Ligand TTP A 1001



Bond lengths



Bond angles



Torsions



Rings

Ligand TTP B 1002



Bond lengths



Bond angles



Torsions



Rings

Ligand TTP B 1003



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	622/644 (96%)	-0.31	25 (4%)	38	37	20, 32, 61, 90	0
1	B	603/644 (93%)	-0.35	9 (1%)	73	72	19, 32, 61, 79	0
All	All	1225/1288 (95%)	-0.33	34 (2%)	53	51	19, 32, 61, 90	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	519	GLU	4.9
1	A	522	THR	4.9
1	A	523	LYS	4.7
1	A	32	LEU	4.7
1	A	516	VAL	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	TTP	B	1003	29/29	0.77	0.29	23,39,48,49	29
3	TTP	A	1004	29/29	0.82	0.25	11,28,40,41	29
3	TTP	B	1002	29/29	0.85	0.17	27,30,68,70	0
2	MG	A	1005	1/1	0.94	0.13	38,38,38,38	0
3	TTP	A	1001	29/29	0.95	0.11	17,31,42,46	0
2	MG	B	1006	1/1	0.97	0.09	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TTP B 1003:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TTP A 1004:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TTP B 1002:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TTP A 1001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.