Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2024 – 07:37 PM EDT

PDB ID	:	5Y39
Title	:	Crystal structure of Ragulator complex (p18 76-145)
Authors	:	Zhang, T.; Ding, J.
Deposited on	:	2017-07-28
Resolution	:	2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	70	10%, 79%, 11%, 6%, •
1	F	70	11%, 39%, •, •, 54%
2	B	125	78%, 14%, 7%
2	G	125	6%, 78%, 15%, •, 6%
3	C	124	85%, 12%, •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	124	6% 82% 10% • 6%
4	D	101	5% 75% 20% • •
4	I	101	20% 53% 6% 41%
5	E	90	83% 13% •
5	J	90	6% 83% 9% • 7%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ragulator complex protein LAMTOR1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	67	534	340	94	99	1	0	0	0
			Total	C	N	O				
1	F	32	227	146	37	44		0	0	0

- Molecule 2 is a protein called Ragulator complex protein LAMTOR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	116	878	552	149	171	6	0	0	0
			Total	C	N	O	S			
2	G	117	872	547	149	170	6	0	0	0

- Molecule 3 is a protein called Ragulator complex protein LAMTOR3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	121	941	606	158	176	1	0	0	0
			Total	C	N	O	S			
3	H	117	893	575	149	168	1	0	0	0

- Molecule 4 is a protein called Ragulator complex protein LAMTOR4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	99	747	465	136	143	3	0	0	0
			Total	C	N	O	S			
4	I	60	380	239	70	70	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	MET	-	expression tag	UNP Q0VGL1
D	0	GLY	-	expression tag	UNP Q0VGL1
I	-1	MET	-	expression tag	UNP Q0VGL1
I	0	GLY	-	expression tag	UNP Q0VGL1

- Molecule 5 is a protein called Ragulator complex protein LAMTOR5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	90	660	403	114	136	7	0	0	0
5	J	84	618	379	106	126	7	0	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	9	9	9	0	0
6	B	15	15	15	0	0
6	C	14	14	14	0	0
6	D	7	7	7	0	0
6	E	6	6	6	0	0
6	G	2	2	2	0	0
6	I	1	1	1	0	0
6	J	4	4	4	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ragulator complex protein LAMTOR1

Chain A:



• Molecule 1: Ragulator complex protein LAMTOR1

Chain F:



• Molecule 2: Ragulator complex protein LAMTOR2

Chain B:



• Molecule 2: Ragulator complex protein LAMTOR2

Chain G:



• Molecule 3: Ragulator complex protein LAMTOR3

Chain C:



• Molecule 3: Ragulator complex protein LAMTOR3

Chain H:



• Molecule 4: Ragulator complex protein LAMTOR4

Chain D:



• Molecule 4: Ragulator complex protein LAMTOR4

Chain I:



• Molecule 5: Ragulator complex protein LAMTOR5

Chain E:



• Molecule 5: Ragulator complex protein LAMTOR5

Chain J:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	110.66Å 117.85Å 157.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.65 47.21 – 2.65	Depositor EDS
% Data completeness (in resolution range)	88.1 (50.01-2.65) 88.1 (47.21-2.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.83 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.216 , 0.263 0.216 , 0.262	Depositor DCC
R_{free} test set	1365 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å^2)	38.4	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 45.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	6808	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.36	0/549	0.57	0/749
1	F	0.35	0/233	0.50	0/321
2	B	0.36	0/888	0.64	0/1202
2	G	0.35	0/881	0.67	0/1195
3	C	0.38	0/958	0.58	0/1300
3	H	0.35	0/910	0.58	0/1238
4	D	0.36	0/756	0.61	0/1020
4	I	0.33	0/376	0.53	0/502
5	E	0.33	0/666	0.56	0/903
5	J	0.31	0/621	0.55	0/839
All	All	0.35	0/6838	0.59	0/9269

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	534	0	531	8	0
1	F	227	0	204	3	0
2	B	878	0	880	4	0
2	G	872	0	869	7	0
3	C	941	0	967	8	0
3	H	893	0	897	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	747	0	756	12	0
4	I	380	0	333	2	0
5	E	660	0	658	9	0
5	J	618	0	615	4	0
6	A	9	0	0	0	0
6	B	15	0	0	0	0
6	C	14	0	0	0	0
6	D	7	0	0	0	0
6	E	6	0	0	0	0
6	G	2	0	0	0	0
6	I	1	0	0	0	0
6	J	4	0	0	0	0
All	All	6808	0	6710	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:5:LEU:HD11	3:C:113:LEU:HD11	1.52	0.90
3:C:9:LEU:HB3	3:C:35:VAL:HG21	1.61	0.81
5:E:139:SER:O	5:E:141:PRO:HD3	1.98	0.64
2:B:77:MET:HB2	3:C:66:SER:HB3	1.82	0.60
2:B:21:SER:HG	2:B:34:TYR:HE1	1.49	0.60
4:I:74:THR:HG22	4:I:88:LYS:HD2	1.87	0.56
1:A:143:LEU:HD13	5:E:164:ILE:HG21	1.88	0.56
5:J:88:GLU:HG3	5:J:116:ARG:HH12	1.73	0.53
1:A:124:ILE:HD11	5:E:150:GLU:HG2	1.93	0.51
2:G:69:LEU:HD21	2:G:72:ILE:HD11	1.92	0.51
1:F:131:GLN:O	1:F:135:ILE:HG22	2.12	0.50
2:B:81:VAL:HG21	3:C:63:LEU:HD13	1.94	0.50
5:E:152:ASP:N	5:E:152:ASP:OD1	2.44	0.49
4:D:43:ILE:HG13	5:E:137:LEU:HD21	1.93	0.49
3:H:16:VAL:HG22	3:H:102:LEU:HD13	1.95	0.49
2:G:14:ALA:HB1	2:G:93:MET:HE3	1.93	0.49
4:D:65:ARG:HD2	5:E:150:GLU:OE2	2.14	0.48
1:F:143:LEU:HD22	5:J:164:ILE:HD12	1.94	0.48
5:J:160:LYS:HA	5:J:165:THR:HA	1.96	0.48
4:D:2:THR:O	4:D:6:THR:HB	2.14	0.48
4:D:43:ILE:HD13	4:D:86:VAL:HG11	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:95:GLU:CD	4:D:95:GLU:H	2.18	0.47
2:G:67:ASP:OD1	3:H:76:ASN:HB2	2.15	0.46
1:A:103:LYS:HG3	3:C:4:ASP:HB3	1.97	0.46
1:F:143:LEU:C	1:F:145:GLN:H	2.20	0.45
1:A:134:ARG:O	1:A:138:TYR:HD1	2.00	0.44
3:H:87:PRO:HD2	3:H:118:ARG:HD2	2.00	0.44
3:C:30:VAL:HA	3:C:31:PRO:HD3	1.85	0.44
2:G:53:TRP:HE1	2:G:85:ARG:NH2	2.15	0.44
4:D:19:LEU:HB3	4:D:86:VAL:HB	1.99	0.44
1:A:126:PHE:HA	1:A:129:LEU:HD12	2.01	0.43
4:D:19:LEU:HB2	4:D:33:LEU:HD23	2.01	0.43
4:I:68:VAL:HB	4:I:75:LEU:HB2	2.01	0.43
2:G:118:PRO:O	2:G:121:GLN:HB2	2.19	0.43
3:H:110:LEU:HB3	3:H:114:PHE:CE2	2.53	0.43
1:A:102:TRP:CE2	3:C:113:LEU:HD13	2.54	0.43
5:E:160:LYS:HG2	5:E:161:HIS:O	2.19	0.42
4:D:16:LEU:HB2	4:D:88:LYS:HD3	2.01	0.42
5:J:95:MET:HG3	5:J:116:ARG:HB3	2.02	0.42
1:A:111:LEU:HG	4:D:63:PHE:O	2.19	0.42
4:D:79:VAL:HG22	4:D:84:VAL:HG22	2.01	0.41
5:E:97:ASN:HA	5:E:98:PRO:HD3	1.94	0.41
2:G:24:LEU:HD11	2:G:89:LEU:HD13	2.01	0.41
4:D:9:LEU:O	4:D:12:ILE:HG12	2.20	0.41
4:D:69:VAL:HA	4:D:74:THR:HB	2.02	0.41
3:H:13:LEU:HA	3:H:19:LEU:HD23	2.02	0.41
1:A:143:LEU:HD11	5:E:113:LEU:HD13	2.03	0.41
2:G:63:ALA:HB1	3:H:44:ALA:HB2	2.03	0.41
2:B:86:VAL:HB	2:B:112:VAL:HG22	2.03	0.40
3:H:66:SER:O	3:H:67:LYS:C	2.59	0.40
3:C:5:LEU:HD11	3:C:113:LEU:CD1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/70 (93%)	61 (94%)	4 (6%)	0	100	100
1	F	30/70 (43%)	28 (93%)	2 (7%)	0	100	100
2	B	114/125 (91%)	110 (96%)	4 (4%)	0	100	100
2	G	115/125 (92%)	108 (94%)	7 (6%)	0	100	100
3	C	119/124 (96%)	117 (98%)	2 (2%)	0	100	100
3	H	115/124 (93%)	111 (96%)	4 (4%)	0	100	100
4	D	97/101 (96%)	93 (96%)	4 (4%)	0	100	100
4	I	46/101 (46%)	42 (91%)	4 (9%)	0	100	100
5	E	88/90 (98%)	85 (97%)	3 (3%)	0	100	100
5	J	78/90 (87%)	74 (95%)	4 (5%)	0	100	100
All	All	867/1020 (85%)	829 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/63 (95%)	53 (88%)	7 (12%)	5	7
1	F	22/63 (35%)	19 (86%)	3 (14%)	3	5
2	B	91/98 (93%)	79 (87%)	12 (13%)	4	5
2	G	89/98 (91%)	80 (90%)	9 (10%)	7	10
3	C	106/108 (98%)	100 (94%)	6 (6%)	20	31
3	H	98/108 (91%)	92 (94%)	6 (6%)	18	29
4	D	81/84 (96%)	72 (89%)	9 (11%)	6	8
4	I	29/84 (34%)	27 (93%)	2 (7%)	15	24
5	E	76/76 (100%)	69 (91%)	7 (9%)	9	13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
5	J	71/76 (93%)	67 (94%)	4 (6%)	21 33
All	All	723/858 (84%)	658 (91%)	65 (9%)	9 14

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	THR
1	A	102	TRP
1	A	104	LYS
1	A	111	LEU
1	A	124	ILE
1	A	143	LEU
1	A	144	SER
2	B	7	LEU
2	B	25	LEU
2	B	31	LEU
2	B	40	THR
2	B	43	ARG
2	B	71	PHE
2	B	85	ARG
2	B	91	LEU
2	B	97	GLU
2	B	109	GLN
2	B	113	GLN
2	B	116	GLU
3	C	11	LYS
3	C	33	ILE
3	C	67	LYS
3	C	102	LEU
3	C	110	LEU
3	C	116	GLU
4	D	6	THR
4	D	15	GLN
4	D	20	VAL
4	D	37	GLU
4	D	46	LEU
4	D	55	LEU
4	D	69	VAL
4	D	74	THR
4	D	94	ARG
5	E	91	LEU
5	E	92	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	116	ARG
5	E	152	ASP
5	E	153	ASN
5	E	161	HIS
5	E	164	ILE
1	F	118	VAL
1	F	131	GLN
1	F	135	ILE
2	G	7	LEU
2	G	24	LEU
2	G	25	LEU
2	G	31	LEU
2	G	68	ASN
2	G	71	PHE
2	G	90	LEU
2	G	91	LEU
2	G	97	GLU
3	H	33	ILE
3	H	102	LEU
3	H	108	LYS
3	H	110	LEU
3	H	117	LEU
3	H	118	ARG
4	I	11	ARG
4	I	69	VAL
5	J	86	THR
5	J	88	GLU
5	J	91	LEU
5	J	143	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	67/70 (95%)	0.35	7 (10%)	6	4	28, 60, 118, 134	0
1	F	32/70 (45%)	1.31	8 (25%)	0	0	68, 103, 198, 206	0
2	B	116/125 (92%)	-0.23	0	100	100	22, 30, 56, 86	0
2	G	117/125 (93%)	0.16	8 (6%)	17	14	27, 40, 111, 133	0
3	C	121/124 (97%)	-0.28	0	100	100	24, 36, 70, 107	0
3	H	117/124 (94%)	0.54	8 (6%)	17	14	34, 76, 112, 135	0
4	D	99/101 (98%)	-0.00	5 (5%)	28	25	25, 45, 109, 151	0
4	I	60/101 (59%)	1.40	20 (33%)	0	0	98, 133, 217, 229	0
5	E	90/90 (100%)	-0.24	0	100	100	24, 37, 74, 87	0
5	J	84/90 (93%)	0.47	5 (5%)	21	18	39, 73, 119, 136	0
All	All	903/1020 (88%)	0.21	61 (6%)	17	14	22, 50, 133, 229	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	65	ASN	6.7
1	A	97	SER	6.6
4	D	97	ILE	5.4
4	D	98	ASP	4.7
2	G	62	GLN	4.7
1	F	118	VAL	4.6
4	D	96	PRO	4.4
4	I	18	TYR	4.3
4	I	84	VAL	4.1
3	H	46	ARG	4.1
4	I	52	GLY	4.0
3	H	43	HIS	4.0
1	F	117	GLN	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	G	59	ASN	3.9
2	G	63	ALA	3.9
3	H	42	GLU	3.7
5	J	96	LYS	3.7
4	I	4	ALA	3.6
1	A	102	TRP	3.6
2	G	64	PHE	3.6
1	F	115	PRO	3.6
4	I	85	PHE	3.5
3	H	41	PRO	3.4
4	I	7	GLN	3.3
3	H	5	LEU	3.2
3	H	33	ILE	3.0
4	I	9	LEU	2.9
2	G	58	ARG	2.9
4	I	66	LEU	2.9
4	I	86	VAL	2.9
4	I	78	THR	2.8
2	G	68	ASN	2.8
1	A	105	LEU	2.7
1	F	121	SER	2.7
4	I	43	ILE	2.7
1	F	116	HIS	2.6
5	J	135	ALA	2.6
4	I	87	VAL	2.6
4	I	65	ARG	2.6
1	F	129	LEU	2.5
4	I	8	GLY	2.5
1	A	103	LYS	2.5
4	I	6	THR	2.4
3	H	50	LEU	2.4
4	D	93	GLY	2.4
4	I	11	ARG	2.4
4	I	27	LEU	2.4
3	H	3	ASP	2.4
5	J	119	LEU	2.3
4	I	22	SER	2.3
5	J	121	ASP	2.3
1	F	120	ALA	2.3
2	G	61	ASN	2.2
4	D	37	GLU	2.2
4	I	79	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	104	LYS	2.1
4	I	10	GLU	2.1
1	A	80	GLU	2.1
5	J	116	ARG	2.0
1	F	124	ILE	2.0
1	A	93	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.