



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2024 – 08:25 pm BST

PDB ID : 2Y6V
Title : Peroxisomal alpha-beta-hydrolase Lpx1 (Yor084w) from *Saccharomyces cerevisiae* (crystal form I)
Authors : Thoms, S.; Niemann, H.H.
Deposited on : 2011-01-26
Resolution : 2.83 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 6% 80% 8% • 11%
1	B	398	 7% 82% 7% • 9%
1	C	398	 4% 85% 8% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEROXISOMAL MEMBRANE PROTEIN LPX1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2855	1809	511	520	15	0	0	0
1	B	363	2915	1844	524	532	15	0	0	0
1	C	377	3023	1909	546	552	16	0	1	0

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	388	ALA	-	expression tag	UNP Q12405
A	389	ALA	-	expression tag	UNP Q12405
A	390	ALA	-	expression tag	UNP Q12405
A	391	LEU	-	expression tag	UNP Q12405
A	392	GLU	-	expression tag	UNP Q12405
A	393	HIS	-	expression tag	UNP Q12405
A	394	HIS	-	expression tag	UNP Q12405
A	395	HIS	-	expression tag	UNP Q12405
A	396	HIS	-	expression tag	UNP Q12405
A	397	HIS	-	expression tag	UNP Q12405
A	398	HIS	-	expression tag	UNP Q12405
B	388	ALA	-	expression tag	UNP Q12405
B	389	ALA	-	expression tag	UNP Q12405
B	390	ALA	-	expression tag	UNP Q12405
B	391	LEU	-	expression tag	UNP Q12405
B	392	GLU	-	expression tag	UNP Q12405
B	393	HIS	-	expression tag	UNP Q12405
B	394	HIS	-	expression tag	UNP Q12405
B	395	HIS	-	expression tag	UNP Q12405
B	396	HIS	-	expression tag	UNP Q12405
B	397	HIS	-	expression tag	UNP Q12405
B	398	HIS	-	expression tag	UNP Q12405
C	388	ALA	-	expression tag	UNP Q12405

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	389	ALA	-	expression tag	UNP Q12405
C	390	ALA	-	expression tag	UNP Q12405
C	391	LEU	-	expression tag	UNP Q12405
C	392	GLU	-	expression tag	UNP Q12405
C	393	HIS	-	expression tag	UNP Q12405
C	394	HIS	-	expression tag	UNP Q12405
C	395	HIS	-	expression tag	UNP Q12405
C	396	HIS	-	expression tag	UNP Q12405
C	397	HIS	-	expression tag	UNP Q12405
C	398	HIS	-	expression tag	UNP Q12405

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	16	Total O 16 16	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	22	Total	O	0	0
			22	22		
3	C	33	Total	O	0	0
			33	33		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PEROXISOMAL MEMBRANE PROTEIN LPX1



- Molecule 1: PEROXISOMAL MEMBRANE PROTEIN LPX1



- Molecule 1: PEROXISOMAL MEMBRANE PROTEIN LPX1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.48Å 87.77Å 125.04Å 90.00° 95.10° 90.00°	Depositor
Resolution (Å)	19.75 – 2.83 19.75 – 2.83	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.75-2.83) 96.0 (19.75-2.83)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.83Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.207 , 0.254 0.206 , 0.247	Depositor DCC
R_{free} test set	1744 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8879	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, PO4, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	2/2872 (0.1%)	0.73	0/3880
1	B	0.76	0/2935	0.74	1/3969 (0.0%)
1	C	0.76	1/3039 (0.0%)	0.79	4/4112 (0.1%)
All	All	0.75	3/8846 (0.0%)	0.76	5/11961 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	343	GLU	CG-CD	5.31	1.59	1.51
1	A	265	CYS	CB-SG	-5.25	1.73	1.81
1	A	335	GLU	CG-CD	5.07	1.59	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	105	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	C	356	PRO	N-CA-C	5.52	126.45	112.10
1	C	206	ASP	CB-CG-OD1	5.38	123.15	118.30
1	B	105	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	C	201	ARG	NE-CZ-NH1	5.36	122.98	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2855	0	2847	19	1
1	B	2915	0	2902	28	1
1	C	3023	0	3019	23	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	16	0	0	1	0
3	B	22	0	0	1	0
3	C	33	0	0	1	0
All	All	8879	0	8768	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:307:GLN:HG3	1:C:317:VAL:HG21	1.51	0.89
1:B:350:LEU:HA	1:B:351:GLN:HB2	1.58	0.85
1:A:350:LEU:HA	1:A:351:GLN:HB2	1.58	0.84
1:B:300:PRO:HG3	1:C:304:LEU:HD12	1.62	0.80
1:C:346:LEU:O	1:C:347:THR:HG23	1.82	0.80
1:B:350:LEU:HA	1:B:351:GLN:CB	2.16	0.75
1:A:350:LEU:HA	1:A:351:GLN:CB	2.17	0.74
1:B:46:ARG:O	1:B:46:ARG:HD3	1.95	0.67
1:A:145:SER:OG	1:A:146:MET:N	2.29	0.66
1:C:122:LEU:O	1:C:126:THR:HB	1.99	0.63
1:A:122:LEU:O	1:A:126:THR:HB	1.98	0.63
1:C:137:ALA:O	1:C:138:LEU:HD23	1.99	0.62
1:A:46:ARG:O	1:A:46:ARG:HD3	2.00	0.62
1:C:95:VAL:HG13	3:C:2011:HOH:O	1.99	0.61
1:B:354:HIS:NE2	1:B:356:PRO:HB3	2.16	0.60
1:B:300:PRO:HB3	1:C:304:LEU:CD1	2.33	0.59
1:B:177:ALA:HB2	1:B:193:PRO:HG3	1.84	0.58
1:B:354:HIS:CE1	1:B:356:PRO:HB3	2.39	0.58
1:B:145:SER:OG	1:B:146:MET:N	2.37	0.58
1:B:300:PRO:HG3	1:C:304:LEU:CD1	2.34	0.58
1:C:145:SER:OG	1:C:146:MET:N	2.38	0.56
1:A:177:ALA:HB2	1:A:193:PRO:HG3	1.87	0.55
1:B:122:LEU:O	1:B:126:THR:HB	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:300:PRO:HA	1:C:304:LEU:HD11	1.91	0.52
1:B:300:PRO:CB	1:C:304:LEU:HD11	2.39	0.52
1:C:304:LEU:O	1:C:308:LYS:HG2	2.10	0.52
1:B:33:LEU:HD11	1:B:120:ASP:HB3	1.90	0.51
1:A:350:LEU:CA	1:A:351:GLN:CB	2.89	0.50
1:B:304:LEU:O	1:B:308:LYS:HG2	2.11	0.49
1:A:304:LEU:O	1:A:308:LYS:HG2	2.13	0.49
1:B:300:PRO:CG	1:C:304:LEU:HD12	2.40	0.49
1:B:300:PRO:CB	1:C:304:LEU:CD1	2.91	0.48
1:B:300:PRO:CA	1:C:304:LEU:HD11	2.44	0.48
1:B:350:LEU:CA	1:B:351:GLN:CB	2.88	0.48
1:A:33:LEU:HD11	1:A:120:ASP:HB3	1.97	0.47
1:C:182:ARG:HB2	1:C:185:LEU:HD12	1.96	0.47
1:C:344:PHE:O	1:C:347:THR:OG1	2.32	0.47
1:B:300:PRO:HB3	1:C:304:LEU:HD11	1.96	0.46
1:C:182:ARG:HB2	1:C:185:LEU:CD1	2.45	0.46
1:A:137:ALA:O	1:A:138:LEU:HD23	2.16	0.46
1:A:341:ILE:O	1:A:345:VAL:HG23	2.16	0.46
1:C:142:ILE:N	1:C:142:ILE:HD12	2.31	0.46
1:B:46:ARG:HD3	1:B:46:ARG:C	2.36	0.45
1:A:46:ARG:HD3	1:A:46:ARG:C	2.36	0.45
1:C:191:GLN:HB2	1:C:270:GLN:NE2	2.32	0.45
1:B:303:GLN:NE2	3:B:2021:HOH:O	2.42	0.44
1:C:47:SER:H	1:C:50:ALA:HB2	1.81	0.44
1:C:180:ALA:HA	1:C:181:GLY:HA3	1.76	0.44
1:B:144:HIS:O	1:B:147:GLY:N	2.50	0.43
1:A:45:ARG:O	1:A:46:ARG:C	2.56	0.43
1:A:142:ILE:HD12	1:A:142:ILE:N	2.34	0.42
1:A:289:HIS:CE1	1:A:303:GLN:HG3	2.55	0.42
1:A:95:VAL:HG13	3:A:2008:HOH:O	2.19	0.42
1:B:175:ARG:NH1	1:B:297:TRP:O	2.53	0.42
1:A:275:PHE:HA	1:B:282:PHE:CZ	2.55	0.42
1:A:267:MET:HB3	1:A:267:MET:HE2	2.01	0.41
1:B:232:LEU:HD22	1:B:232:LEU:O	2.21	0.41
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.86	0.40
1:B:267:MET:HB3	1:B:267:MET:HE2	1.99	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:GLN:O	1:B:369:ARG:NH2[4_645]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/398 (85%)	326 (96%)	11 (3%)	2 (1%)	25	46
1	B	350/398 (88%)	330 (94%)	16 (5%)	4 (1%)	14	30
1	C	368/398 (92%)	350 (95%)	16 (4%)	2 (0%)	29	51
All	All	1057/1194 (88%)	1006 (95%)	43 (4%)	8 (1%)	19	38

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	351	GLN
1	B	351	GLN
1	C	347	THR
1	A	357	GLN
1	B	356	PRO
1	B	357	GLN
1	C	357	GLN
1	B	80	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/342 (91%)	300 (96%)	11 (4%)	36	61
1	B	317/342 (93%)	305 (96%)	12 (4%)	33	59
1	C	327/342 (96%)	311 (95%)	16 (5%)	25	47
All	All	955/1026 (93%)	916 (96%)	39 (4%)	30	56

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	46	ARG
1	A	61	SER
1	A	105	ARG
1	A	126	THR
1	A	131	SER
1	A	145	SER
1	A	176	LYS
1	A	232	LEU
1	A	348	SER
1	A	351	GLN
1	B	3	GLN
1	B	12	THR
1	B	42	GLU
1	B	46	ARG
1	B	61	SER
1	B	105	ARG
1	B	126	THR
1	B	145	SER
1	B	176	LYS
1	B	232	LEU
1	B	348	SER
1	B	351	GLN
1	C	12	THR
1	C	61	SER
1	C	66	VAL
1	C	105	ARG
1	C	126	THR
1	C	131	SER
1	C	145	SER
1	C	178	ILE
1	C	182	ARG
1	C	212	SER
1	C	232	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	346	LEU
1	C	350	LEU
1	C	357	GLN
1	C	372	ASP
1	C	386	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	314	HIS
1	B	4	ASN
1	C	4	ASN
1	C	230	GLN
1	C	314	HIS
1	C	357	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	C	127	1	3,6,7	1.06	0	0,6,8	-	-
1	CME	C	13	1	8,9,10	0.99	1 (12%)	5,9,11	1.69	2 (40%)
1	CME	B	13	1	8,9,10	0.91	0	5,9,11	2.02	2 (40%)
1	CME	A	13	1	8,9,10	0.87	0	5,9,11	2.00	1 (20%)
1	CSO	A	298	1	3,6,7	0.23	0	0,6,8	-	-
1	CSO	B	298	1	3,6,7	0.60	0	0,6,8	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	C	298[B]	1	3,6,7	0.69	0	0,6,8	-	-
1	CSO	A	127	1	3,6,7	0.96	0	0,6,8	-	-
1	CSO	C	26	1	3,6,7	0.50	0	0,6,8	-	-
1	CSO	C	298[A]	1	3,6,7	0.69	0	0,6,8	-	-
1	CSO	A	26	1	3,6,7	0.64	0	0,6,8	-	-
1	CSO	C	205	1	3,6,7	0.69	0	0,6,8	-	-
1	CSO	B	127	1	3,6,7	0.93	0	0,6,8	-	-
1	CSO	A	205	1	3,6,7	0.80	0	0,6,8	-	-
1	CSO	B	26	1	3,6,7	0.34	0	0,6,8	-	-
1	CSO	B	205	1	3,6,7	0.88	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	C	127	1	-	0/1/5/7	-
1	CME	C	13	1	-	3/5/8/10	-
1	CME	B	13	1	-	2/5/8/10	-
1	CME	A	13	1	-	2/5/8/10	-
1	CSO	A	298	1	-	0/1/5/7	-
1	CSO	B	298	1	-	0/1/5/7	-
1	CSO	C	298[B]	1	-	1/1/5/7	-
1	CSO	A	127	1	-	0/1/5/7	-
1	CSO	C	26	1	-	0/1/5/7	-
1	CSO	C	298[A]	1	-	1/1/5/7	-
1	CSO	A	26	1	-	0/1/5/7	-
1	CSO	C	205	1	-	0/1/5/7	-
1	CSO	B	127	1	-	0/1/5/7	-
1	CSO	A	205	1	-	0/1/5/7	-
1	CSO	B	26	1	-	0/1/5/7	-
1	CSO	B	205	1	-	0/1/5/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	13	CME	CB-SG	-2.40	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	13	CME	CB-SG-SD	-3.64	94.39	103.82
1	B	13	CME	CB-SG-SD	-3.45	94.89	103.82
1	C	13	CME	CA-CB-SG	-2.92	102.09	114.55
1	C	13	CME	CB-SG-SD	-2.18	98.18	103.82
1	B	13	CME	CA-CB-SG	-2.13	105.46	114.55

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	298[A]	CSO	N-CA-CB-SG
1	C	298[B]	CSO	N-CA-CB-SG
1	A	13	CME	CA-CB-SG-SD
1	B	13	CME	CA-CB-SG-SD
1	C	13	CME	CA-CB-SG-SD
1	A	13	CME	CZ-CE-SD-SG
1	B	13	CME	CZ-CE-SD-SG
1	C	13	CME	CE-SD-SG-CB
1	C	13	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	B	1382	-	4,4,4	0.84	0	6,6,6	0.76	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	C	1388	-	4,4,4	0.95	0	6,6,6	0.79	0
2	PO4	A	1382	-	4,4,4	0.64	0	6,6,6	0.78	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/398 (88%)	-0.23	25 (7%) 16 10	21, 37, 60, 81	0
1	B	358/398 (89%)	-0.17	26 (7%) 15 9	22, 38, 65, 79	0
1	C	372/398 (93%)	-0.27	15 (4%) 38 30	21, 34, 62, 95	0
All	All	1081/1194 (90%)	-0.22	66 (6%) 21 15	21, 36, 63, 95	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	THR	6.2
1	B	4	ASN	6.0
1	C	177	ALA	5.8
1	A	351	GLN	5.6
1	B	186	PRO	5.5
1	A	178	ILE	5.3
1	C	181	GLY	5.3
1	A	46	ARG	5.2
1	A	45	ARG	5.2
1	A	48	ARG	4.8
1	A	350	LEU	4.7
1	C	178	ILE	4.7
1	A	177	ALA	4.5
1	B	3	GLN	4.4
1	B	187	PRO	4.4
1	B	177	ALA	4.4
1	B	49	THR	4.3
1	B	81	GLU	4.3
1	A	179	GLY	4.1
1	B	46	ARG	4.1
1	A	187	PRO	3.9
1	B	45	ARG	3.9
1	B	178	ILE	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	183	PRO	3.8
1	B	209	ALA	3.7
1	C	180	ALA	3.7
1	A	41	ALA	3.6
1	A	47	SER	3.4
1	A	209	ALA	3.3
1	B	48	ARG	3.2
1	A	207	HIS	3.1
1	C	187	PRO	3.1
1	C	179	GLY	3.1
1	A	352	SER	3.0
1	C	184	GLY	3.0
1	C	384	LYS	3.0
1	C	80	ALA	2.9
1	A	50	ALA	2.9
1	C	185	LEU	2.8
1	B	79	ASP	2.8
1	B	352	SER	2.7
1	C	182	ARG	2.6
1	C	387	LEU	2.6
1	A	356	PRO	2.5
1	B	355	ILE	2.5
1	B	296	ASN	2.5
1	C	382	THR	2.5
1	B	176	LYS	2.5
1	A	353	SER	2.5
1	A	357	GLN	2.5
1	B	353	SER	2.5
1	A	242	LYS	2.5
1	A	296	ASN	2.4
1	C	186	PRO	2.4
1	B	133	ASP	2.3
1	A	176	LYS	2.3
1	B	82	GLY	2.2
1	B	195	ASN	2.2
1	A	186	PRO	2.1
1	A	133	ASP	2.1
1	B	349	PRO	2.1
1	B	350	LEU	2.0
1	B	188	ASP	2.0
1	A	135	HIS	2.0
1	B	354	HIS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	252	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	205	7/8	0.84	0.17	42,43,49,51	0
1	CSO	A	298	7/8	0.87	0.19	38,40,41,44	0
1	CSO	B	298	7/8	0.90	0.26	41,43,45,46	0
1	CSO	B	205	7/8	0.91	0.14	41,42,48,49	0
1	CME	B	13	10/11	0.91	0.17	28,31,35,36	0
1	CME	A	13	10/11	0.94	0.14	27,28,33,33	0
1	CME	C	13	10/11	0.94	0.17	26,27,32,33	0
1	CSO	C	298[A]	7/8	0.94	0.16	31,35,39,39	7
1	CSO	C	298[B]	7/8	0.94	0.16	35,37,38,39	7
1	CSO	C	127	7/8	0.95	0.10	27,28,33,34	0
1	CSO	A	26	7/8	0.95	0.12	33,34,39,40	0
1	CSO	C	26	7/8	0.95	0.11	31,32,36,37	0
1	CSO	B	26	7/8	0.96	0.10	35,36,42,42	0
1	CSO	A	127	7/8	0.96	0.13	28,30,34,35	0
1	CSO	B	127	7/8	0.97	0.13	29,31,36,36	0
1	CSO	C	205	7/8	0.97	0.11	35,36,42,44	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	B	1382	5/5	0.88	0.42	60,60,63,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	C	1388	5/5	0.93	0.45	60,61,62,63	0
2	PO4	A	1382	5/5	0.94	0.34	49,51,53,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.