



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 07:53 PM UTC

PDB ID : 9YXQ / pdb_00009yxq
EMDB ID : EMD-73614
Title : Cryo-EM structure of the *Xenopus laevis* mitotic centromere-associated kinesin (MCAK) bound to the paclitaxel-stabilized microtubule
Authors : Funabiki, H.; Niu, Y.
Deposited on : 2025-10-27
Resolution : 3.39 Å (reported)
Based on initial models : 7SGS, .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14220 (2.89 - 3.89)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	451	 79% 16% 5%
1	3	451	 81% 14% 5%
1	6	451	 82% 13% 5%
1	8	451	 83% 12% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	A	451	 83% 12% 5%
1	a	451	 86% 10% 5%
2	2	445	 81% 15% .
2	4	445	 81% 15% .
2	5	445	 85% 11% .
2	7	445	 86% 10% .
2	B	445	 85% 12% .
2	b	445	 85% 12% .
3	M	739	 45% 9% 45%
3	m	739	 45% 9% 46%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 47551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	1	430	Total	C	N	O	S	0	0
			3372	2137	573	640	22		
1	3	430	Total	C	N	O	S	0	0
			3366	2131	573	640	22		
1	6	430	Total	C	N	O	S	0	0
			3365	2131	573	639	22		
1	8	430	Total	C	N	O	S	0	0
			3365	2131	573	639	22		
1	A	430	Total	C	N	O	S	0	0
			3372	2137	573	640	22		
1	a	430	Total	C	N	O	S	0	0
			3366	2131	573	640	22		

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	2	426	Total	C	N	O	S	0	0
			3351	2106	575	644	26		
2	4	426	Total	C	N	O	S	0	0
			3351	2106	575	644	26		
2	5	426	Total	C	N	O	S	0	0
			3351	2106	575	644	26		
2	7	426	Total	C	N	O	S	0	0
			3351	2106	575	644	26		
2	B	431	Total	C	N	O	S	0	0
			3384	2124	580	654	26		
2	b	432	Total	C	N	O	S	0	0
			3393	2129	581	657	26		

- Molecule 3 is a protein called Kinesin-like protein KIF2C.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	403	Total	C	N	O	S	0	0
			3220	2012	584	604	20		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
3	m	400	Total	C	N	O	S	0	0
			3201	2002	581	598	20		

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	731	SER	-	expression tag	UNP Q91636
M	732	ASN	-	expression tag	UNP Q91636
M	733	SER	-	expression tag	UNP Q91636
M	734	LEU	-	expression tag	UNP Q91636
M	735	GLU	-	expression tag	UNP Q91636
M	736	VAL	-	expression tag	UNP Q91636
M	737	LEU	-	expression tag	UNP Q91636
M	738	PHE	-	expression tag	UNP Q91636
M	739	GLN	-	expression tag	UNP Q91636
m	731	SER	-	expression tag	UNP Q91636
m	732	ASN	-	expression tag	UNP Q91636
m	733	SER	-	expression tag	UNP Q91636
m	734	LEU	-	expression tag	UNP Q91636
m	735	GLU	-	expression tag	UNP Q91636
m	736	VAL	-	expression tag	UNP Q91636
m	737	LEU	-	expression tag	UNP Q91636
m	738	PHE	-	expression tag	UNP Q91636
m	739	GLN	-	expression tag	UNP Q91636

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
4	1	1	Total	Mg	0
			1	1	
4	3	1	Total	Mg	0
			1	1	
4	4	1	Total	Mg	0
			1	1	
4	5	1	Total	Mg	0
			1	1	
4	6	1	Total	Mg	0
			1	1	
4	7	1	Total	Mg	0
			1	1	
4	8	1	Total	Mg	0
			1	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		AltConf
4	A	1	Total	Mg	0
			1	1	
4	B	1	Total	Mg	0
			1	1	
4	a	2	Total	Mg	0
			2	2	

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).




Mol	Chain	Residues	Atoms					AltConf
5	1	1	Total	C	N	O	P	0
			32	10	5	14	3	
5	3	1	Total	C	N	O	P	0
			32	10	5	14	3	
5	6	1	Total	C	N	O	P	0
			32	10	5	14	3	
5	8	1	Total	C	N	O	P	0
			32	10	5	14	3	
5	A	1	Total	C	N	O	P	0
			32	10	5	14	3	
5	a	1	Total	C	N	O	P	0
			32	10	5	14	3	

- Molecule 6 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					AltConf
6	2	1	Total	C	N	O	P	0
			28	10	5	11	2	
6	4	1	Total	C	N	O	P	0
			28	10	5	11	2	
6	5	1	Total	C	N	O	P	0
			28	10	5	11	2	
6	7	1	Total	C	N	O	P	0
			28	10	5	11	2	
6	B	1	Total	C	N	O	P	0
			28	10	5	11	2	
6	b	1	Total	C	N	O	P	0
			28	10	5	11	2	

- Molecule 7 is TAXOL (CCD ID: TA1) (formula: C₄₇H₅₁NO₁₄).




Mol	Chain	Residues	Atoms				AltConf
7	2	1	Total	C	N	O	0
			62	47	1	14	
7	4	1	Total	C	N	O	0
			62	47	1	14	
7	5	1	Total	C	N	O	0
			62	47	1	14	
7	7	1	Total	C	N	O	0
			62	47	1	14	
7	B	1	Total	C	N	O	0
			62	47	1	14	
7	b	1	Total	C	N	O	0
			62	47	1	14	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain






• Molecule 1: Tubulin alpha-1B chain

Chain 8: 83% 12% 5%




• Molecule 1: Tubulin alpha-1B chain

Chain A: 83% 12% 5%




• Molecule 1: Tubulin alpha-1B chain

Chain a: 86% 10% 5%



• Molecule 2: Tubulin beta chain


Chain 2: 81% 15% 5%



ASP
ALA
THR
ALA
ASP
GLU
GLN
GLY
GLU
PHE
GLU
GLU
GLY
ASP
GLU
ALA

• Molecule 2: Tubulin beta chain


Chain 4:  81% 15% .



ASP
GLU
GLN
GLY
GLU
PHE
GLU
GLU
GLY
ASP
GLU
ALA


• Molecule 2: Tubulin beta chain

Chain 5:  85% 11% .




• Molecule 2: Tubulin beta chain

Chain 7:  86% 10% .




• Molecule 2: Tubulin beta chain

Chain B:  85% 12% .




• Molecule 2: Tubulin beta chain


Chain b:  85% 12% .




• Molecule 3: Kinesin-like protein KIF2C



• Molecule 3: Kinesin-like protein KIF2C





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	109829	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	54.6	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.010	Depositor
Minimum map value	-0.005	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.0017	Depositor
Map size ($\text{\AA}$)	371.27997, 371.27997, 371.27997	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.16025, 1.16025, 1.16025	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GTP, TA1, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.11	0/3449	0.27	0/4682
1	3	0.11	0/3442	0.27	0/4673
1	6	0.11	0/3441	0.27	0/4671
1	8	0.10	0/3441	0.27	0/4671
1	A	0.11	0/3449	0.26	0/4682
1	a	0.10	0/3442	0.26	0/4673
2	2	0.11	0/3426	0.28	0/4641
2	4	0.10	0/3426	0.27	0/4641
2	5	0.10	0/3426	0.26	0/4641
2	7	0.10	0/3426	0.25	0/4641
2	B	0.11	0/3459	0.28	0/4687
2	b	0.11	0/3468	0.27	0/4699
3	M	0.28	0/3270	0.51	0/4384
3	m	0.20	0/3251	0.44	0/4357
All	All	0.13	0/47816	0.30	0/64743

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	3372	0	3286	47	0
1	3	3366	0	3279	39	0
1	6	3365	0	3279	35	0
1	8	3365	0	3279	34	0
1	A	3372	0	3286	30	0
1	a	3366	0	3279	23	0
2	2	3351	0	3237	45	0
2	4	3351	0	3237	44	0
2	5	3351	0	3237	32	0
2	7	3351	0	3237	28	0
2	B	3384	0	3262	34	0
2	b	3393	0	3268	36	0
3	M	3220	0	3240	40	0
3	m	3201	0	3231	42	0
4	1	1	0	0	0	0
4	3	1	0	0	0	0
4	4	1	0	0	0	0
4	5	1	0	0	0	0
4	6	1	0	0	0	0
4	7	1	0	0	0	0
4	8	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	a	2	0	0	0	0
5	1	32	0	12	1	0
5	3	32	0	12	0	0
5	6	32	0	12	1	0
5	8	32	0	12	0	0
5	A	32	0	12	1	0
5	a	32	0	12	0	0
6	2	28	0	12	0	0
6	4	28	0	12	0	0
6	5	28	0	12	0	0
6	7	28	0	12	0	0
6	B	28	0	12	0	0
6	b	28	0	12	0	0
7	2	62	0	51	7	0
7	4	62	0	51	8	0
7	5	62	0	51	6	0
7	7	62	0	51	7	0
7	B	62	0	51	5	0
7	b	62	0	51	9	0
All	All	47551	0	46087	513	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 513 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:503:TA1:H193	7:B:503:TA1:H472	1.34	1.07
7:2:502:TA1:H193	7:2:502:TA1:H472	1.42	1.00
7:5:503:TA1:H472	7:5:503:TA1:H193	1.49	0.95
7:7:503:TA1:H472	7:7:503:TA1:H193	1.45	0.94
7:4:503:TA1:H193	7:4:503:TA1:H472	1.48	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	426/451 (94%)	416 (98%)	10 (2%)	0	100	100
1	3	426/451 (94%)	420 (99%)	6 (1%)	0	100	100
1	6	426/451 (94%)	417 (98%)	9 (2%)	0	100	100
1	8	426/451 (94%)	419 (98%)	7 (2%)	0	100	100
1	A	426/451 (94%)	417 (98%)	9 (2%)	0	100	100
1	a	426/451 (94%)	419 (98%)	7 (2%)	0	100	100
2	2	424/445 (95%)	413 (97%)	11 (3%)	0	100	100
2	4	424/445 (95%)	419 (99%)	5 (1%)	0	100	100
2	5	424/445 (95%)	418 (99%)	6 (1%)	0	100	100
2	7	424/445 (95%)	416 (98%)	8 (2%)	0	100	100
2	B	429/445 (96%)	421 (98%)	8 (2%)	0	100	100
2	b	430/445 (97%)	427 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	M	401/739 (54%)	384 (96%)	17 (4%)	0	100	100
3	m	398/739 (54%)	383 (96%)	15 (4%)	0	100	100
All	All	5910/6854 (86%)	5789 (98%)	121 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	364/379 (96%)	363 (100%)	1 (0%)	86	84
1	3	363/379 (96%)	363 (100%)	0	100	100
1	6	363/379 (96%)	363 (100%)	0	100	100
1	8	363/379 (96%)	362 (100%)	1 (0%)	86	84
1	A	364/379 (96%)	364 (100%)	0	100	100
1	a	363/379 (96%)	362 (100%)	1 (0%)	86	84
2	2	367/381 (96%)	367 (100%)	0	100	100
2	4	367/381 (96%)	367 (100%)	0	100	100
2	5	367/381 (96%)	366 (100%)	1 (0%)	86	84
2	7	367/381 (96%)	367 (100%)	0	100	100
2	B	370/381 (97%)	369 (100%)	1 (0%)	86	84
2	b	371/381 (97%)	371 (100%)	0	100	100
3	M	354/657 (54%)	353 (100%)	1 (0%)	86	84
3	m	352/657 (54%)	349 (99%)	3 (1%)	70	76
All	All	5095/5874 (87%)	5086 (100%)	9 (0%)	85	85

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	m	360	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	m	409	TYR
2	B	3	GLU
3	M	239	MET
1	a	367	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
2	7	380	ASN
2	B	258	ASN
1	8	206	ASN
1	A	329	ASN
2	B	349	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 11 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GDP	4	501	-	29,30,30	0.49	0	45,47,47	0.47	0
5	GTP	8	501	4	33,34,34	0.59	0	50,54,54	0.66	0
6	GDP	B	501	-	29,30,30	0.51	0	45,47,47	0.48	0
7	TA1	5	503	-	68,68,68	0.35	0	105,105,105	0.86	2 (1%)
7	TA1	B	503	-	68,68,68	0.37	0	105,105,105	0.84	6 (5%)
5	GTP	1	502	-	33,34,34	0.57	0	50,54,54	0.56	0
7	TA1	b	501	-	68,68,68	0.37	0	105,105,105	0.89	3 (2%)
6	GDP	b	502	-	29,30,30	0.52	0	45,47,47	0.52	0
5	GTP	a	501	4	33,34,34	0.60	0	50,54,54	0.67	0
6	GDP	2	501	-	29,30,30	0.51	0	45,47,47	0.52	0
7	TA1	4	503	-	68,68,68	0.37	0	105,105,105	0.84	3 (2%)
7	TA1	7	503	-	68,68,68	0.35	0	105,105,105	0.87	2 (1%)
6	GDP	7	501	-	29,30,30	0.50	0	45,47,47	0.44	0
7	TA1	2	502	-	68,68,68	0.34	0	105,105,105	0.85	4 (3%)
6	GDP	5	501	-	29,30,30	0.51	0	45,47,47	0.47	0
5	GTP	3	501	4	33,34,34	0.64	0	50,54,54	0.62	0
5	GTP	6	501	-	33,34,34	0.57	0	50,54,54	0.57	0
5	GTP	A	501	4	33,34,34	0.57	0	50,54,54	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GDP	4	501	-	-	0/16/32/32	0/3/3/3
5	GTP	8	501	4	-	2/22/38/38	0/3/3/3
6	GDP	B	501	-	-	0/16/32/32	0/3/3/3
7	TA1	5	503	-	-	2/41/127/127	0/7/7/7
7	TA1	B	503	-	-	2/41/127/127	0/7/7/7
5	GTP	1	502	-	-	3/22/38/38	0/3/3/3
7	TA1	b	501	-	-	2/41/127/127	0/7/7/7
6	GDP	b	502	-	-	1/16/32/32	0/3/3/3
5	GTP	a	501	4	-	3/22/38/38	0/3/3/3
6	GDP	2	501	-	-	0/16/32/32	0/3/3/3
7	TA1	4	503	-	-	11/41/127/127	0/7/7/7
7	TA1	7	503	-	-	10/41/127/127	0/7/7/7
6	GDP	7	501	-	-	0/16/32/32	0/3/3/3
7	TA1	2	502	-	-	8/41/127/127	0/7/7/7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GDP	5	501	-	-	0/16/32/32	0/3/3/3
5	GTP	3	501	4	-	3/22/38/38	0/3/3/3
5	GTP	6	501	-	-	3/22/38/38	0/3/3/3
5	GTP	A	501	4	-	2/22/38/38	0/3/3/3

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	5	503	TA1	C10-C18-C17	4.60	115.28	106.55
7	7	503	TA1	C10-C18-C17	4.53	115.15	106.55
7	2	502	TA1	C10-C18-C17	4.38	114.87	106.55
7	B	503	TA1	C10-C18-C17	4.33	114.77	106.55
7	4	503	TA1	C10-C18-C17	4.24	114.59	106.55

There are no chirality outliers.

5 of 52 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	3	501	GTP	C5'-O5'-PA-O3A
5	3	501	GTP	C5'-O5'-PA-O1A
5	6	501	GTP	C5'-O5'-PA-O1A
5	8	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O3A


There are no ring outliers.


9 monomers are involved in 45 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	5	503	TA1	6	0
7	B	503	TA1	5	0
5	1	502	GTP	1	0
7	b	501	TA1	9	0
7	4	503	TA1	8	0
7	7	503	TA1	7	0
7	2	502	TA1	7	0
5	6	501	GTP	1	0
5	A	501	GTP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand TA1 B 503




Bond lengths




Bond angles





Torsions




Rings









Ligand TA1 4 503




Bond lengths



Bond angles




Torsions




Rings


Ligand TA1 7 503




Bond lengths




Bond angles




Torsions




Rings




Ligand GTP 3 501



Ligand GTP 6 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-73614. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 171




Y Index: 140




Z Index: 167


6.3.2 Raw map



X Index: 171



Y Index: 140




Z Index: 167


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_73614_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 409 nm³; this corresponds to an approximate mass of 370 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.295 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.295 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.39	-	-
Author-provided FSC curve	3.39	3.89	3.43
Unmasked-calculated*	4.14	8.55	4.35

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.14 differs from the reported value 3.39 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-73614 and PDB model 9YXQ. Per-residue inclusion information can be found in [section 3](#) on [page 9](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.0017 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0017) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8910	 0.4980
1	 0.8950	 0.5160
2	 0.9240	 0.5270
3	 0.9150	 0.5340
4	 0.8960	 0.5210
5	 0.9370	 0.4980
6	 0.8650	 0.4790
7	 0.8610	 0.4660
8	 0.8330	 0.4840
A	 0.9500	 0.5300
B	 0.9620	 0.5390
M	 0.7890	 0.3950
a	 0.9520	 0.5410
b	 0.9220	 0.5260
m	 0.7620	 0.4070

