Full wwPDB X-ray Structure Validation Report (i)

Sep 17, 2023 – 03:20 PM EDT

PDB ID	:	4YYM
Title	:	Crystal structure of TAF1 BD2 Bromodomain bound to a butyryllysine peptide
Authors	:	Tang, Y.; Bellon, S.; Cochran, A.G.; Poy, F.
Deposited on	:	2015-03-24
Resolution	:	1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	6% 85% 7% 8%
1	B	144	11% 79% 13% 8%
2	Z	11	18% 55% 9% 36%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2433 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription initiation factor TFIID subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	132	1086	689	175	217	5	0	1	0
1	B	133	1095	694	176	220	5	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1495	GLY	-	expression tag	UNP P21675
A	1496	SER	-	expression tag	UNP P21675
B	1495	GLY	-	expression tag	UNP P21675
B	1496	SER	-	expression tag	UNP P21675

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	Z	7	52	34	9	9	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	105	105	105	0	0
4	B	89	89	89	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	Z	5	5	5	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcription initiation factor TFIID subunit 1

Chain A:



• Molecule 1: Transcription initiation factor TFIID subunit 1

Chain B:

• Molecule 2: Histone H4

Chain Z:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	58.22Å 66.05Å 81.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 1.50 29.70 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.70-1.50) 99.1 (29.70-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.36 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.218 , 0.261 0.216 , 0.258	Depositor DCC
R_{free} test set	2568 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	25.1	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 35.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2433	wwPDB-VP
Average B, all atoms (Å^2)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.30% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BTK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.08	1/1109 (0.1%)	1.16	2/1506 (0.1%)
1	B	1.06	0/1118	1.05	3/1518 (0.2%)
2	Z	0.87	0/22	0.82	0/25
All	All	1.07	1/2249 (0.0%)	1.11	5/3049 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1619	CYS	CB-SG	-5.35	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	1565	ARG	NE-CZ-NH2	-6.19	117.21	120.30
1	B	1570	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	B	1568	PHE	CB-CG-CD2	-5.54	116.92	120.80
1	A	1608	ASP	CB-CG-OD1	5.39	123.15	118.30
1	B	1574	LEU	CA-CB-CG	5.34	127.59	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1086	0	1049	7	0
1	B	1095	0	1055	9	0
2	Z	52	0	59	0	0
3	A	1	0	0	0	0
4	A	105	0	0	1	0
4	B	89	0	0	3	0
4	Z	5	0	0	0	0
All	All	2433	0	2163	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1594[C]:GLN:NE2	1:A:1598:ASN:OD1	1.88	1.05
1:B:1619:CYS:O	1:B:1622:LYS:HG2	1.82	0.80
1:A:1594[C]:GLN:HE21	1:A:1598:ASN:CG	1.87	0.78
1:B:1566:GLU:HG3	4:B:1768:HOH:O	1.92	0.70
1:B:1611:LEU:O	1:B:1615:GLU:HG3	1.95	0.67
1:A:1610:HIS:HE1	4:A:1883:HOH:O	1.84	0.60
1:B:1539:ASP:HB3	1:B:1582:TYR:OH	2.02	0.59
1:B:1530:HIS:HD2	4:B:1708:HOH:O	1.87	0.57
1:A:1504:GLN:NE2	1:A:1563:GLN:HE21	2.03	0.57
1:A:1594[C]:GLN:HE21	1:A:1598:ASN:ND2	2.04	0.55
1:A:1591:LYS:HA	1:A:1594[A]:GLN:NE2	2.24	0.52
1:B:1526:TRP:HA	1:B:1529:HIS:CD2	2.47	0.49
1:B:1606:GLU:HG2	1:B:1607:TYR:CE1	2.49	0.47
1:B:1518:LYS:HE3	4:B:1718:HOH:O	2.17	0.43
1:A:1594[C]:GLN:NE2	1:A:1598:ASN:CG	2.55	0.43
1:B:1513:ASN:O	1:B:1517[A]:GLN:HB2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/144 (91%)	130 (99%)	1 (1%)	0	100	100
1	B	132/144 (92%)	130 (98%)	2 (2%)	0	100	100
2	Z	4/11 (36%)	3 (75%)	1 (25%)	0	100	100
All	All	267/299 (89%)	263 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/134 (92%)	123 (99%)	1 (1%)	81	66
1	B	125/134 (93%)	124 (99%)	1 (1%)	81	66
2	Z	1/3 (33%)	1 (100%)	0	100	100
All	All	250/271 (92%)	248 (99%)	2 (1%)	81	66

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1546	ASN
1	B	1501	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1504	GLN
1	A	1513	ASN
1	A	1546	ASN
1	A	1602	GLN
1	A	1610	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	BTK	Z	5	2	12,13,14	1.07	1 (8%)	9,14,16	1.53	2 (22%)
2	BTK	Z	8	2	12,13,14	0.70	0	9,14,16	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTK	Z	5	2	-	3/12/13/15	-
2	BTK	Z	8	2	-	1/12/13/15	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Z	5	BTK	CE-NZ	-2.55	1.40	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	Z	5	BTK	OAD-CAN-CAJ	-2.95	116.63	122.02
2	Z	5	BTK	CD-CG-CB	-2.57	104.53	113.62

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Z	8	BTK	CA-CB-CG-CD
2	Z	5	BTK	N-CA-CB-CG
2	Z	5	BTK	C-CA-CB-CG
2	Z	5	BTK	CA-CB-CG-CD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	132/144 (91%)	0.42	9 (6%) 17 18	17, 25, 49, 74	0
1	B	133/144 (92%)	0.70	16 (12%) 4 4	19, 30, 48, 67	0
2	Z	5/11 (45%)	1.62	2 (40%) 0 0	39, 44, 58, 59	0
All	All	270/299 (90%)	0.58	27 (10%) 7 7	17, 27, 49, 74	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1500	ASP	8.3
1	A	1629	ALA	6.7
1	B	1631	LEU	5.5
1	B	1629	ALA	4.6
1	A	1630	GLU	4.4
1	A	1631	LEU	4.4
1	B	1632	GLU	4.3
1	A	1626	LEU	4.2
1	B	1620	THR	4.1
1	A	1624	ALA	4.1
1	A	1625	ALA	4.1
1	B	1621	ALA	3.8
1	B	1619	CYS	3.8
1	B	1622	LYS	3.7
1	B	1623	GLU	3.5
1	B	1626	LEU	3.5
1	A	1628	GLU	3.3
1	B	1542	LYS	3.2
2	Z	10	LEU	2.9
1	A	1500	ASP	2.8
1	B	1539	ASP	2.5
1	B	1627	GLU	2.4
2	Z	9	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1622	LYS	2.2
1	B	1625	ALA	2.1
1	B	1618	ILE	2.1
1	B	1526	TRP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BTK	Z	8	14/15	0.83	0.12	32,35,44,46	0
2	BTK	Z	5	14/15	0.90	0.14	21,34,44,44	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CA	A	1701	1/1	0.99	0.03	23,23,23,23	0

6.5 Other polymers (i)

There are no such residues in this entry.