Full wwPDB X-ray Structure Validation Report (i)

Jul 11, 2022 – 07:25 pm BST

PDB ID	:	7Z0W
Title	:	E. coli NfsA bound to NADP+
Authors	:	White, S.A.; Grainger, A.; Parr, R.; Day, M.A.; Jarrom, D.; Graziano, A.; Searle, P.F.; Hyde, E.I.
Deposited on	:	2022-02-23
Resolution	:	2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	94% • ••
1	B	240	96% •
1	C	240	93% 5% ••
1	D	240	97% •
1	E	240	92% 5% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	240	2% 91% 7%
1	G	240	91% 5%
1	H	240	92% 6%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 16522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxygen-insensitive NADPH nitroreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1869	1183	334	344	8	0	0	0
1	B	240	1887	1192	338	349	8	0	0	0
1	C	237	1869	1183	334	344	8	0	0	0
1	D	240	1887	1192	338	349	8	0	0	0
1	E	236	1858	1177	330	343	8	0	0	0
1	F	236	1858	1177	330	343	8	0	0	0
1	G	233	1828	1161	320	339	8	0	0	0
1	H	240	1887	1192	338	349	8	0	0	0

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	E	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	F	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	G	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	H	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is 2'-MONOPHOSPHOADENOSINE-5'-DIPHOSPHATE (three-letter code: ATR) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	B	1	31	10	5	13	3	0	0
			Total	C	N	O	P		
3	D	1	31	10	5	13	3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	B	1	1	1	0	0
			Total	Mg		
4	C	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	179	179	179	0	0
			Total	O		
5	B	151	151	151	0	0
			Total	O		
5	C	189	189	189	0	0
			Total	O		
5	D	182	182	182	0	0
			Total	O		
5	E	125	125	125	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	F	125	125	125	0	0
			Total	O		
5	G	148	148	148	0	0
			Total	O		
5	H	168	168	168	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain A:





• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain B:



• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain C:



• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain D:





• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain E:

• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain F:



• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain G:



• Molecule 1: Oxygen-insensitive NADPH nitroreductase

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	96.77Å 110.76Å 112.25Å 90.00° 103.84° 90.00°	Depositor
Resolution (Å)	81.45 – 2.06 81.45 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.6 (81.45-2.06) 99.6 (81.45-2.06)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.95 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.153 , 0.187 0.162 , 0.192	Depositor DCC
R_{free} test set	6964 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å^2)	31.2	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	16522	wwPDB-VP
Average B, all atoms (Å^2)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6464e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, MG, ATR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.62	0/1907	0.73	0/2586
1	B	0.58	0/1926	0.72	1/2613 (0.0%)
1	C	0.59	0/1907	0.70	2/2586 (0.1%)
1	D	0.63	0/1926	0.75	1/2613 (0.0%)
1	E	0.54	0/1896	0.68	0/2572
1	F	0.57	0/1896	0.69	0/2572
1	G	0.57	0/1866	0.71	0/2533
1	H	0.57	0/1926	0.71	2/2613 (0.1%)
All	All	0.59	0/15250	0.71	6/20688 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	218	ARG	NE-CZ-NH2	5.77	123.19	120.30
1	H	97	LEU	CA-CB-CG	-5.67	102.27	115.30
1	H	103	LEU	CB-CG-CD1	-5.28	102.03	111.00
1	C	103	LEU	CA-CB-CG	5.17	127.19	115.30
1	C	185	ASP	CB-CG-OD1	5.13	122.92	118.30
1	D	218	ARG	NE-CZ-NH2	5.11	122.85	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	207	ASN	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1869	0	1877	12	0
1	B	1887	0	1892	8	0
1	C	1869	0	1877	8	0
1	D	1887	0	1892	4	0
1	E	1858	0	1864	12	0
1	F	1858	0	1864	11	0
1	G	1828	0	1832	10	0
1	H	1887	0	1892	14	0
2	A	31	0	19	0	0
2	B	31	0	19	0	0
2	C	31	0	18	3	0
2	D	31	0	19	0	0
2	E	31	0	19	0	0
2	F	31	0	19	3	0
2	G	31	0	19	0	0
2	H	31	0	17	1	0
3	B	31	0	11	0	0
3	D	31	0	11	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	179	0	0	3	0
5	B	151	0	0	0	0
5	C	189	0	0	2	0
5	D	182	0	0	0	0
5	E	125	0	0	5	0
5	F	125	0	0	4	0
5	G	148	0	0	2	0
5	H	168	0	0	5	0
All	All	16522	0	15161	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:207:ASN:N	5:E:401:HOH:O	1.89	1.04
1:H:90:CYS:SG	5:H:409:HOH:O	2.14	1.04
1:C:208:ARG:NH2	5:C:401:HOH:O	1.93	1.01
1:G:214:ASP:OD1	1:G:217:ARG:NH2	1.97	0.97
1:F:72:GLN:OE1	5:F:401:HOH:O	1.88	0.90
1:A:214:ASP:OD1	5:A:401:HOH:O	1.98	0.81
1:G:134:ASN:ND2	5:G:401:HOH:O	2.15	0.77
1:A:24:GLU:OE1	5:A:402:HOH:O	2.05	0.74
2:F:301:FMN:H5'2	5:F:434:HOH:O	1.88	0.74
1:A:117:ILE:HD11	1:B:114:ASN:OD1	1.87	0.73
1:H:186:LYS:O	5:H:401:HOH:O	2.06	0.72
1:E:195:GLN:NE2	5:E:406:HOH:O	2.17	0.71
1:E:19:ASP:OD2	5:E:402:HOH:O	2.07	0.71
1:H:195:GLN:OE1	5:H:402:HOH:O	2.08	0.71
1:C:117:ILE:HD11	1:D:114:ASN:OD1	1.91	0.70
1:E:20:GLU:OE1	5:E:403:HOH:O	2.10	0.69
1:F:72:GLN:OE1	5:F:403:HOH:O	2.14	0.66
1:F:201:LEU:O	1:F:207:ASN:ND2	2.27	0.66
1:E:223:GLU:OE1	5:E:404:HOH:O	2.14	0.65
1:A:72:GLN:NE2	5:A:403:HOH:O	2.28	0.63
1:G:117:ILE:HD11	1:H:114:ASN:OD1	1.97	0.63
1:G:114:ASN:OD1	1:H:117:ILE:HD11	1.99	0.62
1:G:212:TRP:CZ2	1:G:216:ILE:HD11	2.35	0.62
1:B:24:GLU:OE1	1:B:27:ARG:NH2	2.33	0.61
1:E:114:ASN:OD1	1:F:117:ILE:HD11	1.99	0.61
1:C:114:ASN:OD1	1:D:117:ILE:HD11	2.02	0.59
1:G:216:ILE:HD12	1:H:86:HIS:CD2	2.41	0.56
1:H:12:ARG:NH2	5:H:404:HOH:O	2.32	0.56
1:G:195:GLN:OE1	5:G:402:HOH:O	2.17	0.55
1:A:201:LEU:HD12	1:A:207:ASN:HA	1.89	0.55
1:A:12:ARG:NH1	1:A:163:ASN:OD1	2.41	0.53
1:G:12:ARG:HD2	1:G:164:PRO:O	2.08	0.53
2:C:301:FMN:O3P	2:C:301:FMN:C4'	2.57	0.52
1:H:12:ARG:HD2	1:H:164:PRO:O	2.10	0.52
1:E:209:ARG:O	1:E:210:ASP:CB	2.58	0.52
1:C:12:ARG:HD2	1:C:164:PRO:O	2.10	0.51
1:A:12:ARG:HD2	1:A:164:PRO:O	2.11	0.51
1:H:199:TYR:OH	5:H:403:HOH:O	2.15	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ILE:HD13	1:B:117:ILE:HD13	1.94	0.50
1:A:217:ARG:HG2	1:B:90:CYS:HB2	1.95	0.49
1:F:136:ILE:HG22	1:F:227:PHE:HE2	1.77	0.49
1:E:12:ARG:HD2	1:E:164:PRO:O	2.13	0.48
2:C:301:FMN:H3'	1:D:39:SER:O	2.14	0.48
1:H:40:SER:HB2	1:H:103:LEU:HD21	1.95	0.48
2:H:301:FMN:O2'	2:H:301:FMN:H5'1	2.12	0.48
1:E:117:ILE:HD13	1:F:117:ILE:HD13	1.96	0.47
1:E:117:ILE:HD11	1:F:114:ASN:OD1	2.13	0.47
1:E:208:ARG:HE	1:E:210:ASP:HB2	1.80	0.47
1:F:167:LYS:CE	2:F:301:FMN:H5'1	2.45	0.47
2:C:301:FMN:O3P	2:C:301:FMN:H4'	2.15	0.47
1:F:12:ARG:NH1	5:F:407:HOH:O	2.48	0.47
1:C:217:ARG:NH1	5:C:404:HOH:O	2.46	0.47
1:A:207:ASN:OD1	1:A:208:ARG:N	2.48	0.46
1:A:114:ASN:OD1	1:B:117:ILE:HD11	2.16	0.45
1:B:12:ARG:HD2	1:B:164:PRO:O	2.16	0.45
1:B:140:THR:HG23	1:B:145:LEU:HB2	1.97	0.45
1:A:117:ILE:HD13	1:B:117:ILE:CD1	2.47	0.45
2:F:301:FMN:O4'	2:F:301:FMN:O2'	2.23	0.43
1:H:16:HIS:HB2	1:H:162:ASP:HB2	2.01	0.43
1:F:231:TYR:CZ	1:F:235:GLN:HG3	2.54	0.42
1:C:137:GLU:OE1	1:C:234:LYS:HE2	2.19	0.42
1:C:117:ILE:HD13	1:D:117:ILE:HD13	2.02	0.42
1:H:94:GLN:HB3	1:H:240:ARG:NH2	2.35	0.42
1:C:12:ARG:NH1	1:C:163:ASN:OD1	2.53	0.42
1:E:224:SER:HB2	1:F:97:LEU:HD13	2.03	0.41
1:H:12:ARG:NH1	1:H:163:ASN:OD1	2.55	0.40
1:G:137:GLU:OE2	1:G:234:LYS:HE2	2.21	0.40
1:G:117:ILE:HD13	1:H:117:ILE:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/240 (97%)	224 (96%)	9 (4%)	0	100	100
1	B	238/240 (99%)	228 (96%)	10 (4%)	0	100	100
1	C	233/240 (97%)	223 (96%)	10 (4%)	0	100	100
1	D	238/240 (99%)	231 (97%)	7 (3%)	0	100	100
1	E	232/240 (97%)	225 (97%)	6 (3%)	1 (0%)	34	25
1	F	232/240 (97%)	224 (97%)	8 (3%)	0	100	100
1	G	229/240 (95%)	224 (98%)	5 (2%)	0	100	100
1	H	238/240 (99%)	227 (95%)	11 (5%)	0	100	100
All	All	1873/1920 (98%)	1806 (96%)	66 (4%)	1 (0%)	51	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	210	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/202 (99%)	199 (100%)	1 (0%)	88	89
1	B	202/202 (100%)	201 (100%)	1 (0%)	88	89
1	C	200/202 (99%)	195 (98%)	5 (2%)	47	41
1	D	202/202 (100%)	199 (98%)	3 (2%)	65	62
1	E	199/202 (98%)	196 (98%)	3 (2%)	65	62
1	F	199/202 (98%)	192 (96%)	7 (4%)	36	29
1	G	196/202 (97%)	193 (98%)	3 (2%)	65	62
1	H	202/202 (100%)	197 (98%)	5 (2%)	47	41
All	All	1600/1616 (99%)	1572 (98%)	28 (2%)	59	55

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ARG
1	B	12	ARG
1	C	12	ARG
1	C	24	GLU
1	C	203	ARG
1	C	208	ARG
1	C	218	ARG
1	D	12	ARG
1	D	24	GLU
1	D	234	LYS
1	E	12	ARG
1	E	202	THR
1	E	208	ARG
1	F	12	ARG
1	F	141	LYS
1	F	202	THR
1	F	208	ARG
1	F	209	ARG
1	F	224	SER
1	F	240	ARG
1	G	12	ARG
1	G	201	LEU
1	G	202	THR
1	H	12	ARG
1	H	97	LEU
1	H	141	LYS
1	H	186	LYS
1	H	198	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	113	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	FMN	E	301	-	33,33,33	0.70	0	48,50,50	0.79	0
2	FMN	D	301	-	33,33,33	0.80	1 (3%)	48,50,50	0.84	1 (2%)
2	FMN	B	301	-	33,33,33	0.58	0	48,50,50	0.81	1 (2%)
2	FMN	F	301	-	33,33,33	0.75	0	48,50,50	1.23	4 (8%)
3	ATR	D	302	-	27,33,33	0.79	1 (3%)	35,52,52	0.75	0
3	ATR	B	302	-	27,33,33	0.73	0	35,52,52	0.71	0
2	FMN	C	301	-	33,33,33	1.38	4 (12%)	48,50,50	2.08	8 (16%)
2	FMN	G	301	-	33,33,33	0.59	0	48,50,50	0.75	1 (2%)
2	FMN	H	301	-	33,33,33	1.53	5 (15%)	48,50,50	2.81	20 (41%)
2	FMN	A	301	-	33,33,33	0.72	0	48,50,50	0.83	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	E	301	-	-	1/18/18/18	0/3/3/3
2	FMN	D	301	-	-	1/18/18/18	0/3/3/3
2	FMN	B	301	-	-	2/18/18/18	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	F	301	-	-	6/18/18/18	0/3/3/3
3	ATR	D	302	-	-	4/17/37/37	0/3/3/3
3	ATR	B	302	-	-	3/17/37/37	0/3/3/3
2	FMN	C	301	-	-	10/18/18/18	0/3/3/3
2	FMN	G	301	-	-	0/18/18/18	0/3/3/3
2	FMN	H	301	-	-	8/18/18/18	0/3/3/3
2	FMN	A	301	-	-	0/18/18/18	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	301	FMN	C1'-C2'	4.24	1.58	1.52
2	C	301	FMN	O2'-C2'	-4.16	1.34	1.43
2	C	301	FMN	C1'-C2'	3.52	1.57	1.52
2	H	301	FMN	C10-N1	3.36	1.40	1.33
2	H	301	FMN	C4A-N5	3.14	1.36	1.30
2	C	301	FMN	C4'-C3'	-2.75	1.48	1.53
2	C	301	FMN	O3'-C3'	2.67	1.49	1.43
2	H	301	FMN	C10-N10	2.31	1.42	1.37
3	D	302	ATR	P2'-O2'	2.26	1.63	1.59
2	H	301	FMN	C1'-N10	2.20	1.53	1.48
2	D	301	FMN	P-O1P	2.03	1.57	1.50

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	H	301	FMN	O3'-C3'-C4'	8.57	129.51	108.81
2	H	301	FMN	C5'-C4'-C3'	7.30	126.31	112.20
2	C	301	FMN	C1'-C2'-C3'	7.25	130.04	109.79
2	H	301	FMN	O2'-C2'-C1'	6.51	125.55	109.80
2	C	301	FMN	O3'-C3'-C2'	5.98	123.26	108.81
2	C	301	FMN	O4'-C4'-C5'	5.14	121.47	109.92
2	H	301	FMN	O3'-C3'-C2'	-4.85	97.09	108.81
2	C	301	FMN	C5'-C4'-C3'	-4.65	103.22	112.20
2	C	301	FMN	O3'-C3'-C4'	-4.44	98.09	108.81
2	H	301	FMN	C4A-C10-N1	-4.29	114.76	124.73
2	H	301	FMN	C4A-C10-N10	4.19	122.61	116.48
2	H	301	FMN	C9A-N10-C10	-4.02	114.50	120.77
2	H	301	FMN	C5A-C9A-N10	3.97	122.05	117.95
2	H	301	FMN	C4-C4A-C10	3.91	123.36	116.79
2	C	301	FMN	O2'-C2'-C3'	-3.85	99.74	109.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	F	301	FMN	O2P-P-O5'	3.69	116.55	106.73
2	H	301	FMN	C10-C4A-N5	-3.28	117.91	124.86
2	F	301	FMN	O4'-C4'-C5'	-3.24	102.63	109.92
2	H	301	FMN	O2P-P-O5'	3.15	115.11	106.73
2	F	301	FMN	O5'-C5'-C4'	2.92	117.17	109.36
2	H	301	FMN	C4-N3-C2	-2.88	120.32	125.64
2	F	301	FMN	O3P-P-O5'	-2.74	99.45	106.73
2	H	301	FMN	C1'-N10-C9A	2.70	125.02	120.51
2	H	301	FMN	O4'-C4'-C3'	-2.67	102.62	109.10
2	H	301	FMN	C10-N1-C2	2.53	121.95	116.90
2	H	301	FMN	C9A-C5A-N5	-2.48	119.74	122.43
2	D	301	FMN	O2P-P-O5'	-2.39	100.36	106.73
2	C	301	FMN	C4-N3-C2	-2.33	121.34	125.64
2	G	301	FMN	O3P-P-O2P	2.26	116.27	107.64
2	H	301	FMN	C9A-C9-C8	2.26	123.84	119.30
2	C	301	FMN	O4'-C4'-C3'	-2.24	103.64	109.10
2	B	301	FMN	C4-N3-C2	-2.19	121.60	125.64
2	H	301	FMN	C4A-C4-N3	2.18	118.72	113.19
2	H	301	FMN	C9-C9A-C5A	-2.15	116.04	120.11
2	A	301	FMN	O4'-C4'-C5'	-2.12	105.16	109.92
2	A	301	FMN	C4-N3-C2	-2.10	121.77	125.64
2	H	301	FMN	O5'-C5'-C4'	2.08	114.92	109.36

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	301	FMN	C1'-C2'-C3'-O3'
2	C	301	FMN	C1'-C2'-C3'-C4'
2	C	301	FMN	O2'-C2'-C3'-O3'
2	C	301	FMN	O2'-C2'-C3'-C4'
2	C	301	FMN	C3'-C4'-C5'-O5'
2	C	301	FMN	O4'-C4'-C5'-O5'
2	C	301	FMN	C4'-C5'-O5'-P
2	F	301	FMN	C2'-C3'-C4'-C5'
2	H	301	FMN	C2'-C1'-N10-C10
2	H	301	FMN	C1'-C2'-C3'-O3'
2	H	301	FMN	C2'-C3'-C4'-O4'
2	H	301	FMN	C2'-C3'-C4'-C5'
2	H	301	FMN	O3'-C3'-C4'-O4'
2	H	301	FMN	C3'-C4'-C5'-O5'
2	H	301	FMN	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	302	ATR	C5'-O5'-PA-O1A
3	D	302	ATR	C5'-O5'-PA-O2A
3	D	302	ATR	C5'-O5'-PA-O3A
2	F	301	FMN	O3'-C3'-C4'-C5'
2	H	301	FMN	O3'-C3'-C4'-C5'
2	F	301	FMN	C2'-C3'-C4'-O4'
2	F	301	FMN	C5'-O5'-P-O1P
2	F	301	FMN	C4'-C5'-O5'-P
3	B	302	ATR	PB-O3A-PA-O5'
2	C	301	FMN	C5'-O5'-P-O2P
2	B	301	FMN	C5'-O5'-P-O1P
2	C	301	FMN	C5'-O5'-P-O1P
2	E	301	FMN	C4'-C5'-O5'-P
2	F	301	FMN	O3'-C3'-C4'-O4'
2	B	301	FMN	C4'-C5'-O5'-P
2	C	301	FMN	C5'-O5'-P-O3P
3	B	302	ATR	C2'-O2'-P2'-O1P
3	D	302	ATR	PA-O3A-PB-O2B
3	B	302	ATR	C2'-O2'-P2'-O3P
2	D	301	FMN	C4'-C5'-O5'-P

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	FMN	3	0
2	C	301	FMN	3	0
2	H	301	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FMN E 301



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN D 301



Ligand FMN B 301



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN F 301



Ligand ATR D 302



Bond lengths



Bond angles



Torsions



Rings

Ligand ATR B 302



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN C 301



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN G 301



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN H 301



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN A 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	237/240 (98%)	-0.48	1 (0%)	92	93	20, 31, 58, 80	0
1	B	240/240 (100%)	-0.47	0	100	100	22, 36, 58, 78	0
1	C	237/240 (98%)	-0.49	0	100	100	20, 31, 57, 107	0
1	D	240/240 (100%)	-0.57	0	100	100	19, 30, 49, 68	0
1	E	236/240 (98%)	-0.38	1 (0%)	92	93	24, 39, 73, 105	0
1	F	236/240 (98%)	-0.26	4 (1%)	70	71	22, 40, 75, 134	0
1	G	233/240 (97%)	-0.52	0	100	100	22, 32, 59, 87	0
1	H	240/240 (100%)	-0.49	0	100	100	21, 34, 69, 78	0
All	All	1899/1920 (98%)	-0.46	6 (0%)	94	94	19, 34, 63, 134	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	207	ASN	3.5
1	F	208	ARG	3.2
1	A	207	ASN	3.1
1	E	208	ARG	2.6
1	F	200	TYR	2.5
1	F	209	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MG	C	302	1/1	0.75	0.07	32,32,32,32	0
3	ATR	D	302	31/31	0.91	0.15	33,50,111,148	0
3	ATR	B	302	31/31	0.91	0.14	39,55,148,174	0
2	FMN	F	301	31/31	0.95	0.10	24,31,38,41	0
2	FMN	C	301	31/31	0.95	0.10	19,25,34,40	0
2	FMN	H	301	31/31	0.96	0.10	21,27,40,44	0
2	FMN	A	301	31/31	0.97	0.10	19,24,28,30	0
2	FMN	E	301	31/31	0.97	0.09	22,25,29,31	0
2	FMN	B	301	31/31	0.97	0.09	21,27,34,37	0
2	FMN	G	301	31/31	0.97	0.09	19,23,30,32	0
2	FMN	D	301	31/31	0.98	0.08	19,22,24,29	0
4	MG	B	303	1/1	0.99	0.05	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATR D 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around ATR B 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN F 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN C 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN H 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN E 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN G 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN D 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.