



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:30 AM UTC

PDB ID : 9Z2I / pdb\_00009z2i  
Title : Crystal Structure of the Poly(Hexamethylene Adipamide) (Nylon66) Hydro-  
lase Nyl50 in its Apo form at Room Temperature  
Authors : Capra, N.; Meilleur, F.  
Deposited on : 2025-11-05  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>7% ..</div> </div> </div>
1	B	305	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>9% .</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly(hexamethylene adipamide) hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	4	0
			2188	1363	384	424	17			
1	B	303	Total	C	N	O	S	0	2	0
			2181	1359	384	421	17			

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	82	Total	O	0	0
			82	82		
3	B	90	Total	O	0	1
			91	91		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly(hexamethylene adipamide) hydrolase



- Molecule 1: Poly(hexamethylene adipamide) hydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.76Å 96.40Å 104.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.27 – 2.20 28.27 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.7 (28.27-2.20) 97.6 (28.27-2.20)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.29 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, $R_{free}$	0.148 , 0.196 0.150 , 0.190	Depositor DCC
$R_{free}$ test set	1462 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.3	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 34.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4544	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	5/2223 (0.2%)	1.23	5/3013 (0.2%)
1	B	1.09	2/2213 (0.1%)	1.26	5/2999 (0.2%)
All	All	1.10	7/4436 (0.2%)	1.24	10/6012 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	ARG	NE-CZ	6.85	1.40	1.33
1	A	193	VAL	C-O	-6.45	1.17	1.24
1	A	122	ARG	CZ-NH1	5.64	1.40	1.32
1	A	256	ARG	CZ-NH1	5.54	1.40	1.32
1	A	38	ARG	NE-CZ	-5.18	1.27	1.33
1	B	256	ARG	NE-CZ	5.18	1.38	1.33
1	B	96	ILE	C-O	-5.11	1.18	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	115	ARG	O-C-N	-8.51	116.26	121.71
1	A	256	ARG	CD-NE-CZ	7.44	134.82	124.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	122	ARG	NE-CZ-NH2	-7.39	112.55	119.20
1	A	277	LYS	CB-CA-C	-5.94	101.17	110.74
1	A	84	GLU	CB-CA-C	-5.70	100.08	110.63
1	B	91	VAL	N-CA-CB	-5.64	104.78	112.28
1	B	185	LEU	N-CA-CB	-5.60	101.78	110.57
1	B	256	ARG	CD-NE-CZ	5.17	131.64	124.40
1	A	84	GLU	N-CA-CB	5.08	118.05	110.22
1	B	279	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	123	ARG	Sidechain
1	A	153	ARG	Sidechain
1	A	17	ARG	Sidechain
1	B	137	ARG	Sidechain
1	B	191	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	0	2209	13	0
1	B	2181	0	2204	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	82	0	0	2	0
3	B	91	0	0	2	0
All	All	4544	0	4413	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:TYR:CZ	1:A:211:LYS:HD2	2.31	0.66
1:B:207:TYR:CZ	1:B:211:LYS:HD2	2.35	0.61
1:A:219:ALA:HB1	1:B:91:VAL:HG22	1.85	0.58
1:A:17:ARG:NH2	1:A:111:ASP:OD2	2.37	0.58
1:A:137:ARG:HG3	1:A:137:ARG:HH11	1.71	0.55
1:B:122:ARG:HD3	3:B:501:HOH:O	2.08	0.54
1:A:137:ARG:HD3	3:A:572:HOH:O	2.11	0.51
1:A:137:ARG:CD	3:A:572:HOH:O	2.57	0.51
1:B:129:LYS:HB2	1:B:130:PRO:HD2	1.93	0.51
1:B:249:LEU:HD12	1:B:288:GLY:HA3	1.95	0.49
1:B:172:LYS:HE3	3:B:560:HOH:O	2.13	0.49
1:A:123:ARG:NH1	1:A:126:GLU:CD	2.72	0.47
1:B:251:HIS:ND1	1:B:262:HIS:HE1	2.14	0.46
1:A:123:ARG:HH11	1:A:126:GLU:CD	2.24	0.45
1:A:249:LEU:HD12	1:A:288:GLY:HA3	1.99	0.44
1:B:61:GLY:O	1:B:99:VAL:HA	2.18	0.44
1:B:232:VAL:O	1:B:272:ALA:HA	2.16	0.44
1:A:177:ILE:HD13	1:A:230:GLY:HA3	2.01	0.43
1:A:300:LEU:O	1:A:304:MET:HG2	2.19	0.42
1:A:212:LEU:HD12	1:A:212:LEU:HA	1.92	0.42
1:B:145:THR:O	1:B:180:ASN:HA	2.20	0.42
1:B:5:ILE:HB	1:B:8:ILE:HD12	2.01	0.41
1:A:232:VAL:O	1:A:272:ALA:HA	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/305 (99%)	292 (97%)	9 (3%)	0	100	100
1	B	300/305 (98%)	291 (97%)	9 (3%)	0	100	100
All	All	601/610 (98%)	583 (97%)	18 (3%)	0	100	100



There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/222 (100%)	218 (98%)	5 (2%)	45	61
1	B	221/222 (100%)	220 (100%)	1 (0%)	81	90
All	All	444/444 (100%)	438 (99%)	6 (1%)	59	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	ARG
1	A	123	ARG
1	A	153	ARG
1	A	212	LEU
1	A	305	GLN
1	B	153	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	262	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	302/305 (99%)	-0.09	14 (4%) 37 34	6, 14, 38, 92	3 (0%)
1	B	303/305 (99%)	-0.13	13 (4%) 40 36	6, 13, 44, 87	1 (0%)
All	All	605/610 (99%)	-0.11	27 (4%) 38 35	6, 14, 42, 92	4 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	189	ALA	7.5
1	B	224	GLY	5.1
1	A	89	TYR	4.9
1	A	123	ARG	4.6
1	B	188	THR	4.5
1	A	153	ARG	4.0
1	B	190	GLY	3.4
1	A	223	MET	3.4
1	B	223	MET	3.3
1	A	222	ARG	2.9
1	A	215	LYS	2.8
1	A	17	ARG	2.8
1	B	222	ARG	2.8
1	A	189	ALA	2.7
1	B	304	MET	2.7
1	A	305	GLN	2.5
1	A	200	GLY	2.4
1	B	219	ALA	2.3
1	A	201	GLY	2.3
1	A	199	GLU	2.3
1	B	89	TYR	2.2
1	B	279	ASP	2.2
1	B	221	ALA	2.2
1	A	218	SER	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	191	ARG	2.1
1	A	277	LYS	2.0
1	B	220	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NA	B	401	1/1	0.97	0.09	6,6,6,6	0
2	NA	A	401	1/1	0.98	0.07	5,5,5,5	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.