



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:32 AM UTC

PDB ID : 9Z2L / pdb\_00009z2l  
Title : The ubiquitin-associated domain of human thirty-eight negative kinase-1 rigidly fused to the 1TEL crystallization chaperone  
Authors : Wilson, E.W.; Bradford, M.J.; Averett, J.C.; Averett, B.J.; Anderson, E.; Anderson, A.; Doukov, T.; Moody, J.D.  
Deposited on : 2025-11-05  
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	<div> <div>16%</div> <div> <div></div> <div>89%</div> <div>5%</div> <div>6%</div> </div> </div>
1	B	164	<div> <div>21%</div> <div> <div></div> <div>86%</div> <div>6%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription factor ETV6,Non-receptor tyrosine-protein kinase TNK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	0	0
			1172	743	209	218	2			
1	B	151	Total	C	N	O	S	0	1	0
			1131	725	202	202	2			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP P41212
A	3	HIS	-	expression tag	UNP P41212
A	4	HIS	-	expression tag	UNP P41212
A	5	HIS	-	expression tag	UNP P41212
A	6	HIS	-	expression tag	UNP P41212
A	7	HIS	-	expression tag	UNP P41212
A	8	HIS	-	expression tag	UNP P41212
A	9	HIS	-	expression tag	UNP P41212
A	10	HIS	-	expression tag	UNP P41212
A	11	HIS	-	expression tag	UNP P41212
A	12	HIS	-	expression tag	UNP P41212
A	46	SER	ARG	engineered mutation	UNP P41212
A	78	GLU	VAL	engineered mutation	UNP P41212
A	90	VAL	LEU	engineered mutation	UNP Q13470
A	109	ALA	CYS	engineered mutation	UNP Q13470
A	143	ALA	CYS	engineered mutation	UNP Q13470
B	2	MET	-	initiating methionine	UNP P41212
B	3	HIS	-	expression tag	UNP P41212
B	4	HIS	-	expression tag	UNP P41212
B	5	HIS	-	expression tag	UNP P41212
B	6	HIS	-	expression tag	UNP P41212
B	7	HIS	-	expression tag	UNP P41212
B	8	HIS	-	expression tag	UNP P41212
B	9	HIS	-	expression tag	UNP P41212

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	HIS	-	expression tag	UNP P41212
B	11	HIS	-	expression tag	UNP P41212
B	12	HIS	-	expression tag	UNP P41212
B	46	SER	ARG	engineered mutation	UNP P41212
B	78	GLU	VAL	engineered mutation	UNP P41212
B	90	VAL	LEU	engineered mutation	UNP Q13470
B	109	ALA	CYS	engineered mutation	UNP Q13470
B	143	ALA	CYS	engineered mutation	UNP Q13470

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Cl	0	0
			3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	62	Total 62	O 62	0	0
5	B	42	Total 42	O 42	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.00Å 77.85Å 119.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.50 – 2.02 59.50 – 2.02	Depositor EDS
% Data completeness (in resolution range)	75.0 (59.50-2.02) 73.0 (59.50-2.02)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.26 (at 2.00Å)	Xtriage
Refinement program	PDB-REDO, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.231 , 0.274 0.244 , 0.282	Depositor DCC
$R_{free}$ test set	908 reflections (3.68%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.3	Xtriage
Anisotropy	0.486	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2416	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/1200	0.32	0/1636
1	B	0.11	0/1161	0.21	0/1584
All	All	0.17	0/2361	0.27	0/3220

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1172	0	1066	4	0
1	B	1131	0	1024	5	0
2	A	5	0	0	0	0
3	A	1	0	0	0	0
4	B	3	0	0	0	0
5	A	62	0	0	0	0
5	B	42	0	0	0	0
All	All	2416	0	2090	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:VAL:HG21	1:A:144:TRP:CE3	2.51	0.45
1:B:95:MET:O	1:B:99:LEU:HG	2.17	0.45
1:A:132:LEU:HD22	1:A:147:LEU:HG	1.99	0.44
1:B:129:VAL:HG21	1:B:144:TRP:CE3	2.53	0.43
1:A:89:GLU:HG3	1:A:92:ARG:HH21	1.84	0.43
1:A:17:PRO:O	1:A:21:ARG:HG3	2.19	0.43
1:B:80:TYR:O	1:B:84:GLN:HG2	2.19	0.42
1:B:60:LEU:HD12	1:B:63:LEU:HD12	2.02	0.41
1:B:24:PRO:O	1:B:57:GLY:HA3	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/164 (93%)	150 (99%)	2 (1%)	0	100	100
1	B	150/164 (92%)	148 (99%)	2 (1%)	0	100	100
All	All	302/328 (92%)	298 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/145 (74%)	108 (100%)	0	100	100
1	B	97/145 (67%)	97 (100%)	0	100	100
All	All	205/290 (71%)	205 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	B	91	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	201	-	4,4,4	0.24	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	154/164 (93%)	1.14	26 (16%)	4 4	26, 43, 84, 101	0
1	B	151/164 (92%)	1.28	35 (23%)	2 2	21, 47, 91, 124	1 (0%)
All	All	305/328 (92%)	1.21	61 (20%)	3 2	21, 44, 88, 124	1 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	20	LEU	9.9
1	B	20	LEU	6.0
1	A	14	ILE	5.7
1	A	19	HIS	5.1
1	A	18	ALA	5.0
1	A	51	ASN	4.7
1	A	12	HIS	4.2
1	B	61	LEU	4.2
1	B	49	ASP	4.0
1	B	22	LEU	4.0
1	A	26	TYR	3.9
1	A	22	LEU	3.9
1	B	62	LEU	3.9
1	B	51	ASN	3.8
1	B	50	SER	3.8
1	B	14	ILE	3.8
1	A	23	GLN	3.6
1	B	18	ALA	3.6
1	A	13	SER	3.5
1	A	135	LEU	3.4
1	B	31	ASP	3.3
1	B	48	ILE	3.3
1	A	25	ILE	3.2
1	A	136	SER	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	16	LEU	3.0
1	B	19	HIS	3.0
1	B	53	PHE	3.0
1	B	41	ASN	2.9
1	A	106	HIS	2.7
1	A	24	PRO	2.7
1	A	133	PHE	2.7
1	B	70	TYR	2.6
1	B	63	LEU	2.6
1	B	46	SER	2.6
1	B	26	TYR	2.6
1	A	50	SER	2.6
1	B	25	ILE	2.5
1	B	161	VAL	2.5
1	A	137	SER	2.5
1	A	134	HIS	2.4
1	B	86	ILE	2.4
1	B	28	SER	2.4
1	B	57	GLY	2.4
1	A	49	ASP	2.4
1	B	135	LEU	2.3
1	A	17	PRO	2.3
1	B	16	LEU	2.3
1	B	32	VAL	2.2
1	B	47	PRO	2.2
1	B	90	VAL	2.2
1	B	23	GLN	2.2
1	B	85	HIS	2.2
1	B	52	THR	2.2
1	B	67	ASP	2.2
1	B	104	VAL	2.1
1	B	73	PRO	2.1
1	B	137	SER	2.1
1	A	27	TRP	2.1
1	A	48	ILE	2.0
1	A	53	PHE	2.0
1	A	62	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CL	B	203	1/1	0.74	0.18	78,78,78,78	0
3	NA	A	202	1/1	0.84	0.15	53,53,53,53	0
4	CL	B	201	1/1	0.92	0.09	59,59,59,59	0
4	CL	B	202	1/1	0.93	0.08	62,62,62,62	0
2	SO4	A	201	5/5	0.96	0.06	50,50,52,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.