



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2026 – 05:26 PM EDT

PDB ID : 9Z4R / pdb_00009z4r
Title : Crystal structure of MAIT A-F7 TCR-MR1-compound 6 complex
Authors : Awad, W.; Rossjohn, R.
Deposited on : 2025-11-11
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	C	271	
2	B	100	
2	F	100	
3	D	204	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	204	
4	E	246	
4	H	246	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	ACT	H	302	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 14918 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	14	0
			2253	1445	387	408	13			
1	C	268	Total	C	N	O	S	0	8	0
			2253	1446	387	409	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	conflict	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	conflict	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	98	Total	C	N	O	S	0	0	0
			782	502	135	142	3			
2	F	100	Total	C	N	O	S	0	0	0
			818	523	139	152	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
F	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called TCRa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	192	Total	C	N	O	S	0	8	0
			1504	959	239	296	10			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	201	Total	C	N	O	S	0	9	0
			1614	1023	258	323	10			

- Molecule 4 is a protein called TCRb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	242	Total	C	N	O	S	0	6	0
			1908	1203	327	368	10			
4	H	244	Total	C	N	O	S	0	11	0
			1962	1239	337	373	13			

- Molecule 5 is 1-[(6-acetyl-2-oxo-2,3-dihydropteridin-4-yl)amino]-1-deoxy-D-ribose (CCD ID: A1C0R) (formula: C₁₃H₁₇N₅O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			23	13	5	5		
5	C	1	Total	C	N	O	0	0
			23	13	5	5		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		
6	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	F	1	Total	Na	0	0
			1	1		
7	H	1	Total	Na	0	0
			1	1		

- Molecule 8 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	F	1	Total	C	O	0	0
			4	2	2		
8	H	1	Total	C	O	0	0
			4	2	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	280	Total	O	0	0
			280	280		
9	B	79	Total	O	0	0
			79	79		
9	C	315	Total	O	0	0
			315	315		
9	D	159	Total	O	0	0
			159	159		
9	E	164	Total	O	0	0
			164	164		
9	F	134	Total	O	0	0
			134	134		
9	G	270	Total	O	0	0
			270	270		
9	H	331	Total	O	0	0
			331	331		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein





- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 3: TCRa




- Molecule 3: TCRa



- Molecule 4: TCRb



- Molecule 4: TCRb



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	216.43Å 69.90Å 142.60Å 90.00° 104.21° 90.00°	Depositor
Resolution (Å)	39.80 – 2.08 39.80 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.80-2.08) 88.8 (39.80-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.94 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.181 , 0.215 0.182 , 0.214	Depositor DCC
R_{free} test set	1998 reflections (1.61%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.165	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14918	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, A1C0R, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2360	0.33	0/3205
1	C	0.15	0/2339	0.34	0/3180
2	B	0.13	0/805	0.32	0/1097
2	F	0.13	0/841	0.36	0/1141
3	D	0.13	0/1557	0.33	0/2112
3	G	0.15	0/1673	0.39	0/2265
4	E	0.13	0/1971	0.34	0/2687
4	H	0.15	0/2044	0.36	0/2781
All	All	0.14	0/13590	0.35	0/18468

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2253	0	2157	15	0
1	C	2253	0	2141	22	0
2	B	782	0	726	1	0
2	F	818	0	773	2	0
3	D	1504	0	1408	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1614	0	1556	10	0
4	E	1908	0	1777	13	0
4	H	1962	0	1866	22	0
5	A	23	0	0	0	0
5	C	23	0	0	0	0
6	A	24	0	32	0	0
6	C	6	0	8	1	0
6	F	6	0	8	0	0
7	F	1	0	0	0	0
7	H	1	0	0	0	0
8	F	4	0	3	0	0
8	H	4	0	3	2	0
9	A	280	0	0	0	0
9	B	79	0	0	1	0
9	C	315	0	0	1	0
9	D	159	0	0	0	0
9	E	164	0	0	1	0
9	F	134	0	0	0	0
9	G	270	0	0	0	0
9	H	331	0	0	2	0
All	All	14918	0	12458	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:2:GLN:HE21	3:D:27[A]:SER:H	1.30	0.79
3:D:2:GLN:HE21	3:D:27[B]:SER:H	1.31	0.79
3:G:2:GLN:HE21	3:G:27:SER:H	1.35	0.75
1:A:8:PHE:CE2	1:A:95[B]:MET:HG3	2.23	0.73
4:H:19[B]:MET:HE1	4:H:112:LEU:HD21	1.75	0.69
4:H:210:ARG:HE	4:H:212:GLN:NE2	1.98	0.62
1:C:210:ILE:O	4:H:206[A]:ARG:HG2	2.00	0.61
1:C:222:VAL:HA	4:H:200[A]:THR:HG21	1.84	0.60
4:E:155:HIS:HB3	4:E:216:TYR:HB2	1.83	0.60
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.37	0.60
3:G:147[A]:LYS:HD2	3:G:188:ASN:HD21	1.67	0.60
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.84	0.59
3:G:28:GLY:HA3	3:G:93[A]:SER:OG	2.02	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149[A]:GLU:OE2	4:H:99:GLU:HB2	2.03	0.58
3:G:150:ASP:HB2	3:G:177:LYS:HD2	1.86	0.57
8:H:302:ACT:H3	9:H:406:HOH:O	2.06	0.55
1:C:54:LEU:HD13	1:C:58[A]:HIS:CD2	2.41	0.55
3:G:2:GLN:NE2	3:G:27:SER:H	2.05	0.54
4:H:210:ARG:HE	4:H:212:GLN:HE21	1.54	0.54
1:A:95[B]:MET:HE1	1:A:109:PHE:CD1	2.42	0.54
4:H:180:GLU:OE2	8:H:302:ACT:H2	2.08	0.54
4:E:95:VAL:HG12	4:E:96:TRP:CD1	2.43	0.54
1:C:230:ILE:HD11	4:H:206[B]:ARG:HD3	1.89	0.53
3:D:164:MET:HE1	4:E:198:SER:HB3	1.92	0.51
3:D:36:GLN:HB2	3:D:46:LEU:HD11	1.93	0.50
3:D:2:GLN:NE2	3:D:27[A]:SER:H	2.06	0.50
3:D:2:GLN:NE2	3:D:27[B]:SER:H	2.06	0.50
3:D:28:GLY:HA3	3:D:93[A]:SER:OG	2.11	0.50
1:A:54:LEU:HD13	1:A:58[B]:HIS:CE1	2.47	0.50
1:C:146:ASN:CG	1:C:149:GLU:HB2	2.38	0.49
3:D:159:LEU:HD21	4:E:196:ARG:HB2	1.93	0.49
1:C:58[B]:HIS:CE1	3:D:94:ASN:HD21	2.31	0.49
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.47	0.48
4:H:128:VAL:HG23	4:H:238:ALA:HB3	1.94	0.48
1:C:211[A]:TYR:HB2	1:C:259:GLU:HB3	1.95	0.48
1:C:211[B]:TYR:HB3	1:C:259:GLU:HB3	1.95	0.48
4:E:96:TRP:HB3	9:E:301:HOH:O	2.12	0.48
1:C:28:VAL:HG23	1:C:33:ILE:HD13	1.93	0.48
3:G:2:GLN:HE21	3:G:27:SER:N	2.07	0.47
4:H:19[B]:MET:CE	4:H:112:LEU:HD21	2.43	0.47
1:C:152:TYR:CD1	4:E:100:GLY:HA3	2.49	0.47
1:C:210:ILE:O	4:H:206[B]:ARG:HD2	2.15	0.47
1:A:19:VAL:HG11	9:B:134:HOH:O	2.15	0.47
4:H:120:ASN:HB2	9:H:472:HOH:O	2.14	0.47
1:A:43:LYS:HE2	1:A:58[B]:HIS:HE1	1.80	0.46
4:E:19[A]:MET:HB3	4:E:19[A]:MET:HE2	1.49	0.46
1:A:151:LEU:HD22	3:G:51:LEU:HD12	1.96	0.46
4:E:128:VAL:HG23	4:E:238:ALA:HB3	1.97	0.46
4:E:19[A]:MET:HE1	4:E:112:LEU:HD21	1.97	0.46
1:C:241:TRP:HE1	6:C:302:GOL:C3	2.29	0.46
1:C:151:LEU:HD22	3:D:51:LEU:HD12	1.98	0.45
4:H:210:ARG:HH21	4:H:212:GLN:HE22	1.63	0.45
1:C:1:ARG:HD2	9:C:559:HOH:O	2.16	0.45
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:221:ILE:HD11	1:C:225:ILE:HG13	1.98	0.45
1:A:152:TYR:CD1	4:H:100:GLY:HA3	2.52	0.44
1:C:156:TRP:CZ3	1:C:161:CYS:HB2	2.52	0.44
3:D:54:LEU:HD11	3:D:61:SER:HB3	1.98	0.44
3:G:61[A]:SER:OG	3:G:74:LEU:HB3	2.17	0.44
3:D:18:VAL:HG12	3:D:78:LEU:HD21	1.98	0.44
4:H:19[B]:MET:HB3	4:H:19[B]:MET:HE2	1.41	0.44
4:H:208:HIS:HD2	4:H:241:TRP:CE2	2.36	0.44
1:A:8:PHE:CZ	1:A:95[B]:MET:HG3	2.54	0.43
1:C:111:GLN:HG2	1:C:121:ILE:HG23	1.99	0.43
1:C:169[A]:LEU:HD23	1:C:176:LEU:HD13	2.00	0.43
4:H:8:PRO:HD2	4:H:21[A]:LEU:HD22	2.00	0.43
3:D:134:PHE:HB2	3:D:186:PHE:CE2	2.54	0.43
4:E:19[A]:MET:HE1	4:E:112:LEU:CD2	2.49	0.43
1:A:43:LYS:HE2	1:A:58[B]:HIS:CE1	2.53	0.43
1:A:4:SER:HB3	1:A:99:GLU:HG2	2.02	0.42
4:H:21[B]:LEU:HD13	4:H:76:LEU:HD23	2.01	0.42
1:A:215:MET:HG3	1:A:257:HIS:CD2	2.54	0.42
4:E:79:GLU:H	4:E:79:GLU:HG2	1.58	0.42
4:H:27[A]:MET:HE1	4:H:105:PHE:CD2	2.55	0.42
4:H:208:HIS:HB2	4:H:241:TRP:CZ3	2.55	0.42
4:E:131:PRO:HG2	4:E:142:ALA:HB1	2.02	0.41
1:C:46:ARG:HD3	1:C:46:ARG:HA	1.93	0.41
1:C:45:PRO:HB3	1:C:59:TRP:CH2	2.56	0.41
3:G:152:TYR:O	3:G:173:ALA:HA	2.21	0.41
1:A:214:TRP:H	1:A:225:ILE:HD13	1.85	0.40
4:E:152:TYR:HB2	4:E:188:ARG:HG2	2.04	0.40
3:G:159[B]:LEU:HB2	4:H:172:CYS:HB2	2.03	0.40
4:H:125:GLU:O	4:H:148:ALA:HA	2.21	0.40
2:F:0:MET:HE2	2:F:2:GLN:HE21	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/271 (102%)	270 (98%)	5 (2%)	0	100	100
1	C	272/271 (100%)	266 (98%)	6 (2%)	0	100	100
2	B	96/100 (96%)	95 (99%)	1 (1%)	0	100	100
2	F	98/100 (98%)	98 (100%)	0	0	100	100
3	D	194/204 (95%)	190 (98%)	4 (2%)	0	100	100
3	G	208/204 (102%)	203 (98%)	5 (2%)	0	100	100
4	E	246/246 (100%)	243 (99%)	3 (1%)	0	100	100
4	H	253/246 (103%)	250 (99%)	3 (1%)	0	100	100
All	All	1642/1642 (100%)	1615 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/241 (101%)	239 (98%)	5 (2%)	48	54
1	C	240/241 (100%)	231 (96%)	9 (4%)	29	30
2	B	83/95 (87%)	81 (98%)	2 (2%)	43	47
2	F	90/95 (95%)	89 (99%)	1 (1%)	65	73
3	D	163/181 (90%)	160 (98%)	3 (2%)	51	58
3	G	185/181 (102%)	179 (97%)	6 (3%)	34	36
4	E	204/212 (96%)	199 (98%)	5 (2%)	42	46
4	H	214/212 (101%)	211 (99%)	3 (1%)	59	66
All	All	1423/1458 (98%)	1389 (98%)	34 (2%)	47	47

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	LYS
1	A	187[A]	ASN
1	A	187[B]	ASN
1	A	221	ILE
1	A	253	LEU
2	B	70	PHE
2	B	97	ARG
1	C	1	ARG
1	C	16	ILE
1	C	73	PHE
1	C	149	GLU
1	C	215	MET
1	C	222	VAL
1	C	224	GLU
1	C	246	LEU
1	C	247	ASP
3	D	27[A]	SER
3	D	27[B]	SER
3	D	131	VAL
4	E	19[A]	MET
4	E	19[B]	MET
4	E	79	GLU
4	E	97	THR
4	E	140	GLN
2	F	40	LEU
3	G	27	SER
3	G	96	GLN
3	G	140	GLN
3	G	146	SER
3	G	159[A]	LEU
3	G	159[B]	LEU
4	H	99	GLU
4	H	206[A]	ARG
4	H	206[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	70	GLN
1	A	203	HIS
2	B	2	GLN
1	C	111	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	153	GLN
1	C	177	GLN
3	D	2	GLN
4	E	138	HIS
4	E	168	HIS
2	F	2	GLN
2	F	13	HIS
2	F	17	ASN
3	G	2	GLN
3	G	3	ASN
3	G	38	HIS
3	G	94	ASN
3	G	145	GLN
3	G	188	ASN
4	H	65	ASN
4	H	181	GLN
4	H	212	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	ACT	F	102	-	3,3,3	1.19	0	3,3,3	1.15	0
6	GOL	A	304	-	5,5,5	0.30	0	5,5,5	0.48	0
6	GOL	C	302	-	5,5,5	0.32	0	5,5,5	0.40	0
5	A1C0R	A	301	1	24,24,25	1.14	2 (8%)	24,33,35	0.81	1 (4%)
6	GOL	F	103	-	5,5,5	0.32	0	5,5,5	0.35	0
5	A1C0R	C	301	1	24,24,25	1.13	2 (8%)	24,33,35	0.89	2 (8%)
6	GOL	A	305	-	5,5,5	0.32	0	5,5,5	0.51	0
8	ACT	H	302	-	3,3,3	1.08	0	3,3,3	1.30	0
6	GOL	A	302	-	5,5,5	0.32	0	5,5,5	0.41	0
6	GOL	A	303	-	5,5,5	0.35	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	304	-	-	2/4/4/4	-
6	GOL	C	302	-	-	4/4/4/4	-
5	A1C0R	A	301	1	-	4/17/17/19	0/2/2/2
6	GOL	F	103	-	-	0/4/4/4	-
5	A1C0R	C	301	1	-	4/17/17/19	0/2/2/2
6	GOL	A	305	-	-	2/4/4/4	-
6	GOL	A	302	-	-	0/4/4/4	-
6	GOL	A	303	-	-	3/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	301	A1C0R	C7-C23	3.95	1.47	1.41
5	C	301	A1C0R	C7-C23	3.82	1.47	1.41
5	C	301	A1C0R	C5-C4	2.63	1.44	1.39
5	A	301	A1C0R	C5-C4	2.58	1.43	1.39

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	301	A1C0R	C4-N24-C23	-2.53	115.69	118.45
5	A	301	A1C0R	C4-N24-C23	-2.50	115.73	118.45
5	C	301	A1C0R	C14-N13-C12	-2.34	121.75	126.69

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	301	A1C0R	N11-C12-N13-C14
5	A	301	A1C0R	C1-C2-C4-C5
5	A	301	A1C0R	C1-C2-C4-N24
5	C	301	A1C0R	N11-C12-N13-C14
5	C	301	A1C0R	C1-C2-C4-C5
5	C	301	A1C0R	C1-C2-C4-N24
6	A	304	GOL	O1-C1-C2-C3
6	A	303	GOL	O1-C1-C2-C3
6	A	305	GOL	O1-C1-C2-C3
6	C	302	GOL	O1-C1-C2-C3
6	C	302	GOL	C1-C2-C3-O3
6	A	304	GOL	O1-C1-C2-O2
6	A	303	GOL	O1-C1-C2-O2
6	A	303	GOL	O2-C2-C3-O3
6	C	302	GOL	O1-C1-C2-O2
6	A	305	GOL	O1-C1-C2-O2
6	C	302	GOL	O2-C2-C3-O3
5	A	301	A1C0R	N13-C14-C15-C17
5	C	301	A1C0R	N13-C14-C15-C17


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	302	GOL	1	0
8	H	302	ACT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1C0R A 301





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/271 (97%)	0.03	11 (4%) 40 42	17, 33, 68, 92	18 (6%)
1	C	268/271 (98%)	-0.12	7 (2%) 57 60	16, 30, 64, 95	11 (4%)
2	B	98/100 (98%)	0.56	1 (1%) 79 82	28, 52, 73, 79	0
2	F	100/100 (100%)	-0.09	2 (2%) 65 67	24, 35, 55, 69	1 (1%)
3	D	192/204 (94%)	0.53	19 (9%) 13 13	13, 45, 85, 109	11 (5%)
3	G	201/204 (98%)	-0.22	3 (1%) 72 74	11, 28, 52, 70	15 (7%)
4	E	242/246 (98%)	0.46	11 (4%) 38 40	20, 44, 77, 85	10 (4%)
4	H	244/246 (99%)	-0.25	2 (0%) 82 84	14, 29, 45, 77	16 (6%)
All	All	1610/1642 (98%)	0.08	56 (3%) 47 49	11, 34, 71, 109	82 (5%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	150	ASP	4.1
1	A	221	ILE	3.7
1	C	221	ILE	3.5
3	D	195	THR	3.4
3	D	159	LEU	3.2
4	E	96	TRP	3.2
3	D	149	SER	3.2
1	A	52	GLU	3.1
3	D	180	PHE	3.1
1	A	19	VAL	3.0
4	E	241	TRP	3.0
2	F	99	MET	3.0
3	D	196	PHE	2.9
4	E	205	PRO	2.8
3	D	197	PHE	2.8
3	G	201	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	E	221	ASN	2.6
1	A	17	HIS	2.6
1	C	246	LEU	2.6
4	E	243	ARG	2.6
3	D	188	ASN	2.6
1	A	222	VAL	2.5
3	D	177	LYS	2.5
3	D	117	ALA	2.5
3	D	124	SER	2.5
3	G	202	SER	2.5
4	E	2	ALA	2.4
4	E	97	THR	2.4
3	D	147	LYS	2.4
4	E	178	LEU	2.4
3	D	191	ILE	2.4
3	G	163	SER	2.4
1	A	211	TYR	2.3
1	C	219	GLU	2.3
1	A	253	LEU	2.3
3	D	130	SER	2.3
4	E	207	ASN	2.3
2	F	0	MET	2.3
1	C	270	PRO	2.3
4	E	99[A]	GLU	2.2
4	E	227	ASP	2.2
1	A	55	ALA	2.2
1	A	270	PRO	2.2
3	D	190	ILE	2.2
3	D	192	PRO	2.2
1	C	16	ILE	2.1
4	H	244	ALA	2.1
1	A	248	PRO	2.1
3	D	146	SER	2.1
2	B	1	ILE	2.1
3	D	118	VAL	2.1
1	C	17	HIS	2.1
1	C	211[A]	TYR	2.1
3	D	121	LEU	2.0
4	H	1	ASN	2.0
1	A	196	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	A	303	6/6	0.62	0.21	51,57,67,68	0
6	GOL	C	302	6/6	0.82	0.15	48,53,53,62	0
8	ACT	H	302	4/4	0.84	0.12	22,49,56,56	0
6	GOL	A	305	6/6	0.87	0.15	44,52,60,62	0
6	GOL	A	304	6/6	0.88	0.14	32,43,48,52	0
8	ACT	F	102	4/4	0.89	0.10	28,33,36,51	0
6	GOL	A	302	6/6	0.92	0.13	35,43,51,54	0
6	GOL	F	103	6/6	0.95	0.09	32,35,38,42	0
5	A1C0R	A	301	23/24	0.96	0.06	23,25,28,30	0
5	A1C0R	C	301	23/24	0.96	0.06	22,25,29,29	0
7	NA	F	101	1/1	0.97	0.09	39,39,39,39	0
7	NA	H	301	1/1	0.97	0.04	35,35,35,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1C0R A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1C0R C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.