



Full wwPDB EM Validation Report ⓘ

Oct 28, 2025 – 06:10 PM JST

PDB ID : 8ZCZ / pdb_00008zcz
EMDB ID : EMD-39942
Title : Cryo-EM structure of eSaCas9_NNG-guide RNA-target DNA complex in an interrogation state
Authors : Omura, S.N.; Nakagawa, R.; Yamashita, K.; Nishimasu, H.; Nureki, O.
Deposited on : 2024-04-30
Resolution : 2.90 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev129
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	13054 (2.40 - 3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1056	
2	B	99	
3	C	43	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	43	<div><div><div></div><div></div><div></div></div><div>5%21%14%65%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18215 atoms, of which 8333 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR-associated endonuclease Cas9.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	830	Total	C	H	N	O	S	1	0
			13823	4377	6922	1228	1283	13		

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP J7RUA5
A	-1	SER	-	expression tag	UNP J7RUA5
A	0	HIS	-	expression tag	UNP J7RUA5
A	413	ALA	ASN	conflict	UNP J7RUA5
A	420	ALA	ARG	conflict	UNP J7RUA5
A	782	LYS	GLU	conflict	UNP J7RUA5
A	800	ARG	LEU	conflict	UNP J7RUA5
A	927	LYS	THR	conflict	UNP J7RUA5
A	929	ASN	LYS	conflict	UNP J7RUA5
A	968	ARG	ASN	conflict	UNP J7RUA5
A	985	ALA	ASN	conflict	UNP J7RUA5
A	986	SER	ASN	conflict	UNP J7RUA5
A	991	ALA	ARG	conflict	UNP J7RUA5
A	1017	PHE	ILE	conflict	UNP J7RUA5
A	1021	SER	ALA	conflict	UNP J7RUA5

- Molecule 2 is a RNA chain called sgRNA.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	B	96	Total	C	H	N	O	P	0	0
			2948	905	922	357	668	96		


- Molecule 3 is a DNA chain called Target DNA strand.

Mol	Chain	Residues	Atoms							AltConf	Trace
3	C	32	Total	C	H	N	O	P	S	0	0
			989	313	336	122	180	31	7		


- Molecule 4 is a DNA chain called Non-target DNA strand.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
4	D	15	455	146	153	49	92	15	0	0

• Molecule 3: Target DNA strand



• Molecule 4: Non-target DNA strand



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	156154	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50.0	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.788	Depositor
Minimum map value	-0.463	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.06	Depositor
Map size (Å)	170.4318, 170.4318, 170.4318	wwPDB
Map dimensions	154, 154, 154	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1067, 1.1067, 1.1067	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AS, GS, SC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/7022	1.08	5/9406 (0.1%)
2	B	0.55	0/2265	1.03	10/3527 (0.3%)
3	C	0.35	0/571	1.20	6/877 (0.7%)
4	D	0.40	0/336	1.43	6/515 (1.2%)
All	All	0.51	0/10194	1.09	27/14325 (0.2%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	44	U	O3'-P-O5'	-11.56	86.65	104.00
2	B	18	G	O3'-P-O5'	-11.40	86.89	104.00
4	D	3	DG	O3'-P-O5'	-8.72	90.92	104.00
3	C	7	DG	O3'-P-O5'	-8.47	91.30	104.00
2	B	19	G	O3'-P-O5'	-8.36	91.46	104.00
2	B	45	C	O3'-P-O5'	-7.60	92.60	104.00
1	A	690	PHE	CA-CB-CG	6.19	119.99	113.80
3	C	6	DA	O3'-P-O5'	-6.18	94.72	104.00
4	D	3	DG	C2'-C3'-O3'	-6.12	102.32	111.50
4	D	7	DA	O3'-P-O5'	-6.11	94.84	104.00
4	D	0	DC	O3'-P-O5'	-5.96	95.06	104.00
4	D	1	DT	O3'-P-O5'	-5.89	95.17	104.00
2	B	46	U	O3'-P-O5'	-5.53	95.71	104.00
3	C	9	DC	C2'-C3'-O3'	-5.50	103.25	111.50
1	A	486	ASP	CA-CB-CG	5.45	118.05	112.60
2	B	18	G	C4'-C3'-O3'	-5.44	104.84	113.00
3	C	9	DC	O3'-P-O5'	-5.43	95.86	104.00
2	B	19	G	C1'-C2'-O2'	5.41	116.51	108.40
4	D	6	DT	O3'-P-O5'	-5.31	96.03	104.00
2	B	89	U	O3'-P-O5'	-5.28	96.07	104.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	7	DG	C2'-C3'-O3'	-5.24	103.64	111.50
2	B	14	G	O3'-P-O5'	-5.24	96.14	104.00
3	C	6	DA	C2'-C3'-O3'	-5.18	103.73	111.50
1	A	622	LYS	CB-CA-C	-5.17	102.77	110.88
1	A	1014	PRO	N-CA-CB	-5.08	99.41	102.81
2	B	56	A	O3'-P-O5'	-5.07	96.40	104.00
1	A	704	ASP	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6901	6922	7071	29	0
2	B	2026	922	1017	0	0
3	C	653	336	364	2	0
4	D	302	153	172	1	0
All	All	9882	8333	8624	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:534:CYS:SG	1:A:576:LYS:HG3	2.10	0.91
1:A:582:LYS:O	1:A:586:ARG:NH2	2.12	0.81
1:A:918:LEU:HB2	1:A:963:ILE:HD11	1.77	0.67
1:A:534:CYS:SG	1:A:541:ILE:HD11	2.42	0.60
1:A:564:SER:HA	1:A:641:PHE:HB3	1.85	0.58
1:A:570:ASN:O	1:A:622:LYS:HG2	2.04	0.58
1:A:530:GLN:OE1	1:A:573:VAL:HG23	2.05	0.57
1:A:536:TYR:CE2	1:A:575:VAL:HG11	2.42	0.55
1:A:541:ILE:HG23	1:A:553:TYR:CE1	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:916:VAL:HG12	1:A:963:ILE:HD12	1.94	0.49
1:A:608:ILE:HD12	1:A:626:LEU:HD13	1.95	0.48
1:A:514:LYS:HE2	1:A:548:ASN:OD1	2.12	0.48
1:A:587:THR:HG23	1:A:632:ILE:HD12	1.95	0.48
1:A:560:PRO:HD2	1:A:632:ILE:HD11	1.95	0.48
1:A:564:SER:O	1:A:645:ASN:ND2	2.46	0.47
1:A:558:ILE:HD13	1:A:626:LEU:HG	1.97	0.47
3:C:21:DA:H2'	3:C:22:DC:C6	2.50	0.46
3:C:9:DC:H2''	3:C:10:DA:C8	2.51	0.46
1:A:752:GLU:HA	1:A:755:ILE:HG23	1.96	0.46
1:A:534:CYS:O	1:A:538:LEU:N	2.47	0.46
1:A:557:HIS:HA	1:A:571:ASN:O	2.18	0.44
1:A:527:HIS:CD2	1:A:532:GLY:HA2	2.52	0.44
1:A:710:ASN:ND2	1:A:763:ILE:HD11	2.34	0.43
1:A:927:LYS:NZ	4:D:4:DC:OP2	2.51	0.43
1:A:573:VAL:HG22	1:A:575:VAL:HG13	2.00	0.43
1:A:718:TRP:CD2	1:A:753:ILE:HG23	2.54	0.43
1:A:506:GLU:HA	1:A:509:ILE:HG12	2.01	0.41
1:A:26:ARG:HG2	1:A:26:ARG:O	2.21	0.41
1:A:916:VAL:HG12	1:A:963:ILE:CD1	2.51	0.40
1:A:986:SER:OG	1:A:989:LEU:HB2	2.20	0.40
1:A:592:LEU:HA	1:A:597:SER:OG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	825/1056 (78%)	791 (96%)	33 (4%)	1 (0%)	48 77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	758	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	765/971 (79%)	751 (98%)	14 (2%)	54 82

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	SER
1	A	116	ARG
1	A	174	SER
1	A	539	GLU
1	A	567	ASN
1	A	573	VAL
1	A	578	GLU
1	A	611	LEU
1	A	628	GLU
1	A	632	ILE
1	A	639	LYS
1	A	644	ARG
1	A	646	LEU
1	A	829	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	HIS
1	A	570	ASN
1	A	607	HIS
1	A	633	ASN
1	A	701	HIS
1	A	758	HIS
1	A	759	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	831	HIS
1	A	835	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	94/99 (94%)	10 (10%)	0

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	9	G
2	B	20	C
2	B	45	C
2	B	46	U
2	B	55	A
2	B	63	A
2	B	65	G
2	B	74	A
2	B	87	G
2	B	90	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GS	C	17	3,2	18,24,25	0.98	1 (5%)	19,35,38	0.75	1 (5%)
3	AS	C	15	3,2	18,23,24	0.75	0	17,33,36	1.01	1 (5%)
3	AS	C	16	3,2	18,23,24	0.69	0	17,33,36	0.82	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SC	C	18	3,2	17,20,21	0.40	0	24,28,31	0.55	0
3	GS	C	19	3,2	18,24,25	1.06	2 (11%)	19,35,38	0.67	0
3	SC	C	13	3,2	17,20,21	0.31	0	24,28,31	0.43	0
3	SC	C	14	3,2	17,20,21	0.31	0	24,28,31	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GS	C	17	3,2	-	2/2/21/22	0/3/3/3
3	AS	C	15	3,2	-	2/3/21/22	0/3/3/3
3	AS	C	16	3,2	-	2/3/21/22	0/3/3/3
3	SC	C	18	3,2	-	1/7/21/22	0/2/2/2
3	GS	C	19	3,2	-	0/2/21/22	0/3/3/3
3	SC	C	13	3,2	-	2/7/21/22	0/2/2/2
3	SC	C	14	3,2	-	0/7/21/22	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	19	GS	C5-C6	-2.97	1.41	1.47
3	C	17	GS	C5-C6	-2.50	1.42	1.47
3	C	19	GS	C8-N7	-2.12	1.31	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	15	AS	C4'-O4'-C1'	-2.74	102.84	109.45
3	C	17	GS	O6-C6-C5	2.21	128.69	124.37
3	C	16	AS	C5-C6-N6	2.18	123.66	120.35

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	15	AS	C3'-C4'-C5'-O5'
3	C	16	AS	C3'-C4'-C5'-O5'
3	C	17	GS	C3'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	15	AS	O4'-C4'-C5'-O5'
3	C	16	AS	O4'-C4'-C5'-O5'
3	C	17	GS	O4'-C4'-C5'-O5'
3	C	13	SC	O4'-C4'-C5'-O5'
3	C	13	SC	C3'-C4'-C5'-O5'
3	C	18	SC	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-39942. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 77




Y Index: 77




Z Index: 77


6.2.2 Raw map



X Index: 77



Y Index: 77




Z Index: 77


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 81




Y Index: 64




Z Index: 76


6.3.2 Raw map



X Index: 73



Y Index: 71




Z Index: 50


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.06. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_39942_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 77 nm^3 ; this corresponds to an approximate mass of 69 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.90	-	-
Author-provided FSC curve	2.90	3.19	2.94
Unmasked-calculated*	3.16	3.54	3.19

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-39942 and PDB model 8ZCZ. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.06 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.06).

9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.06) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8240	<div></div> 0.4840
A	<div></div> 0.7980	<div></div> 0.4750
B	<div></div> 0.9170	<div></div> 0.5300
C	<div></div> 0.7700	<div></div> 0.4490
D	<div></div> 0.7290	<div></div> 0.4410

