



wwPDB EM Validation Summary Report ⓘ

Mar 30, 2026 – 02:06 PM UTC

PDB ID : 9ZWE / pdb_00009zwe
EMDB ID : EMD-74907
Title : Soluble ectodomain of Herpes simplex virus 2 (HSV-2) glycoprotein B (gB) in the prefusion conformation in complex with 2c and D48 Fabs
Authors : Sponholtz, M.R.; Johnson, N.V.; McLellan, J.S.
Deposited on : 2026-01-01
Resolution : 2.75 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10570 (2.25 - 3.25)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	738	
1	B	738	
1	G	738	
2	C	228	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	228	
2	L	228	
3	D	214	
3	I	214	
3	M	214	
4	E	226	
4	J	226	
4	N	226	
5	F	219	
5	K	219	
5	O	219	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 48423 atoms, of which 23889 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	B	568	Total	C	H	N	O	S	0	0
			8990	2883	4420	803	860	24		
1	A	568	Total	C	H	N	O	S	0	0
			8990	2883	4420	803	860	24		
1	G	568	Total	C	H	N	O	S	0	0
			8990	2883	4420	803	860	24		

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	172	ASN	HIS	conflict	UNP P08666
B	174	THR	TYR	conflict	UNP P08666
B	198	ALA	THR	conflict	UNP P08666
B	238	CYS	THR	conflict	UNP P08666
B	665	PRO	SER	conflict	UNP P08666
B	666	PRO	THR	conflict	UNP P08666
B	676	CYS	GLU	conflict	UNP P08666
A	172	ASN	HIS	conflict	UNP P08666
A	174	THR	TYR	conflict	UNP P08666
A	198	ALA	THR	conflict	UNP P08666
A	238	CYS	THR	conflict	UNP P08666
A	665	PRO	SER	conflict	UNP P08666
A	666	PRO	THR	conflict	UNP P08666
A	676	CYS	GLU	conflict	UNP P08666
G	172	ASN	HIS	conflict	UNP P08666
G	174	THR	TYR	conflict	UNP P08666
G	198	ALA	THR	conflict	UNP P08666
G	238	CYS	THR	conflict	UNP P08666
G	665	PRO	SER	conflict	UNP P08666
G	666	PRO	THR	conflict	UNP P08666
G	676	CYS	GLU	conflict	UNP P08666

- Molecule 2 is a protein called Antibody D48 Heavy Chain.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	H	123	Total	C	H	N	O	S	0	0
			1887	603	933	169	176	6		
2	C	123	Total	C	H	N	O	S	0	0
			1887	603	933	169	176	6		
2	L	123	Total	C	H	N	O	S	0	0
			1887	603	933	169	176	6		

- Molecule 3 is a protein called Antibody D48 Light Chain.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	I	108	Total	C	H	N	O	S	0	0
			1653	527	819	142	161	4		
3	D	108	Total	C	H	N	O	S	0	0
			1653	527	819	142	161	4		
3	M	108	Total	C	H	N	O	S	0	0
			1653	527	819	142	161	4		

- Molecule 4 is a protein called Antibody 2c Heavy Chain.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	J	122	Total	C	H	N	O	S	0	0
			1902	617	945	157	179	4		
4	E	122	Total	C	H	N	O	S	0	0
			1902	617	945	157	179	4		
4	N	122	Total	C	H	N	O	S	0	0
			1902	617	945	157	179	4		


- Molecule 5 is a protein called Antibody 2c Light Chain.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	K	112	Total	C	H	N	O	S	0	0
			1709	548	846	145	167	3		
5	F	112	Total	C	H	N	O	S	0	0
			1709	548	846	145	167	3		
5	O	112	Total	C	H	N	O	S	0	0
			1709	548	846	145	167	3		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Envelope glycoprotein B




• Molecule 1: Envelope glycoprotein B





ASN
THR
LYS
VAL
ASP
LYS
LYS
VAL
GLU
PRO
LYS
SER
CYS
ASP


• Molecule 4: Antibody 2c Heavy Chain



VAL
SER
TRP
ASN
GLY
SER
GLY
ALA
LEU
THR
THR
SER
GLY
VAL
HIS
THR
PHE
PRO
ALA
VAL
LEU
GLN
SER
TYR
SER
SER
GLY
SER
VAL
VAL
PHE
PRO
VAL
SER
ALA
SER
PRO
SER
SER
SER
LEU
GLY
THR
GLN
THR
TYR
ILE
CYS
ASN
VAL
VAL
SER
ASN
HIS
GLY
LYS
CYS
PRO
VAL
SER
ASN
THR
LYS
VAL
PHE
PRO
GLU
LYS
VAL
THR

GLU
PRO
LYS
SER
CYS
ASP


• Molecule 5: Antibody 2c Light Chain



PRO
ARG
GLU
ALA
VAL
VAL
GLN
TRP
LYS
THR
VAL
ASP
ASN
ALA
LEU
GLN
SER
GLY
ASN
SER
GLN
GLU
VAL
THR
GLU
GLN
ASP
SER
SER
LYS
VAL
SER
THR
ALA
PRO
SER
VAL
SER
PHE
ILE
PHE
THR
PRO
THR
LEU
SER
SER
LYS
ALA
ASP
TYR
GLU
LYS
SER
GLY
HIS
LYS
HIS
VAL
THR
SER
VAL
VAL
CYS
CYS
GLU
VAL
LEU
LEU
ASN
THR
HIS
GLN
PHE
TYR
PRO
GLY

LEU
SER
SER
PRO
VAL
THR
LYS
SER
PHE
ASN
ARG
GLY
GLU
CYS


• Molecule 5: Antibody 2c Light Chain



ARG
GLU
LYS
VAL
GLN
TRP
LYS
VAL
ASP
ASN
ALA
LEU
GLN
SER
GLY
ASN
SER
GLN
GLU
SER
VAL
THR
GLU
GLN
ASP
SER
SER
LYS
VAL
SER
THR
ALA
PRO
SER
SER
VAL
SER
PHE
ILE
PHE
THR
LEU
THR
LEU
SER
SER
LYS
ALA
ASP
TYR
GLU
LYS
HIS
LYS
HIS
VAL
TYR
ALA
CYS
CYS
GLU
VAL
THR
HIS
GLN
PHE
TYR
LEU
SER

SER
SER
PRO
VAL
THR
LYS
SER
PHE
ASN
ARG
GLY
GLU
CYS

• Molecule 5: Antibody 2c Light Chain



GLU
ALA
VAL
VAL
TRP
LYS
VAL
ASP
ASN
ALA
LEU
GLN
SER
GLY
ASN
SER
GLN
GLU
SER
VAL
THR
GLU
GLN
ASP
SER
LYS
ASP
SER
THR
TYR
SER
LEU
SER
SER
SER
PHE
ILE
PHE
PRO
THR
LEU
SER
SER
LYS
ALA
ASP
TYR
GLU
LYS
HIS
LYS
VAL
TYR
ALA
CYS
CYS
GLU
VAL
THR
HIS
GLN
GLY
PHE
TYR
PRO
SER

SER
PRO
VAL
THR
LYS
SER
PHE
ASN
ARG
GLY
GLU
CYS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	227517	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	28.490	Depositor
Minimum map value	-0.262	Depositor
Average map value	-0.008	Depositor
Map value standard deviation	0.530	Depositor
Recommended contour level	5.3	Depositor
Map size ($\text{\AA}$)	359.94238, 359.94238, 359.94238	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.8332, 0.8332, 0.8332	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/4678	0.40	0/6355
1	B	0.15	0/4678	0.40	0/6355
1	G	0.15	0/4678	0.40	0/6355
2	C	0.12	0/976	0.36	0/1324
2	H	0.12	0/976	0.36	0/1324
2	L	0.12	0/976	0.36	0/1324
3	D	0.10	0/854	0.32	0/1161
3	I	0.10	0/854	0.32	0/1161
3	M	0.10	0/854	0.32	0/1161
4	E	0.11	0/984	0.35	0/1338
4	J	0.12	0/984	0.35	0/1338
4	N	0.11	0/984	0.35	0/1338
5	F	0.11	0/884	0.29	0/1197
5	K	0.11	0/884	0.30	0/1197
5	O	0.11	0/884	0.29	0/1197
All	All	0.13	0/25128	0.37	0/34125

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	G	0	1
3	D	0	1
3	I	0	1
3	M	0	1
All	All	0	6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	401	ARG	Sidechain
1	B	401	ARG	Sidechain
3	D	53	ARG	Sidechain
1	G	401	ARG	Sidechain
3	I	53	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4570	4420	4416	82	0
1	B	4570	4420	4416	79	0
1	G	4570	4420	4416	88	0
2	C	954	933	932	14	0
2	H	954	933	932	13	0
2	L	954	933	932	14	0
3	D	834	819	819	11	0
3	I	834	819	819	10	0
3	M	834	819	819	9	0
4	E	957	945	945	11	0
4	J	957	945	945	11	0
4	N	957	945	945	8	0
5	F	863	846	846	7	0
5	K	863	846	846	6	0
5	O	863	846	846	6	0
All	All	24534	23889	23874	318	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 318 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:613:GLN:OE1	1:G:621:ARG:NH2	2.12	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:613:GLN:OE1	1:A:621:ARG:NH2	2.14	0.81
1:B:613:GLN:OE1	1:B:621:ARG:NH2	2.15	0.80
1:B:126:ARG:NH1	1:B:389:THR:O	2.15	0.79
1:A:126:ARG:NH1	1:A:389:THR:O	2.15	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	560/738 (76%)	533 (95%)	27 (5%)	0	100	100
1	B	560/738 (76%)	532 (95%)	28 (5%)	0	100	100
1	G	560/738 (76%)	533 (95%)	27 (5%)	0	100	100
2	C	121/228 (53%)	117 (97%)	4 (3%)	0	100	100
2	H	121/228 (53%)	118 (98%)	3 (2%)	0	100	100
2	L	121/228 (53%)	117 (97%)	4 (3%)	0	100	100
3	D	106/214 (50%)	99 (93%)	7 (7%)	0	100	100
3	I	106/214 (50%)	99 (93%)	7 (7%)	0	100	100
3	M	106/214 (50%)	99 (93%)	7 (7%)	0	100	100
4	E	120/226 (53%)	115 (96%)	5 (4%)	0	100	100
4	J	120/226 (53%)	115 (96%)	5 (4%)	0	100	100
4	N	120/226 (53%)	115 (96%)	5 (4%)	0	100	100
5	F	110/219 (50%)	106 (96%)	4 (4%)	0	100	100
5	K	110/219 (50%)	106 (96%)	4 (4%)	0	100	100
5	O	110/219 (50%)	106 (96%)	4 (4%)	0	100	100
All	All	3051/4875 (63%)	2910 (95%)	141 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/623 (79%)	493 (100%)	0	100	100
1	B	493/623 (79%)	493 (100%)	0	100	100
1	G	493/623 (79%)	493 (100%)	0	100	100
2	C	104/195 (53%)	104 (100%)	0	100	100
2	H	104/195 (53%)	104 (100%)	0	100	100
2	L	104/195 (53%)	104 (100%)	0	100	100
3	D	90/185 (49%)	90 (100%)	0	100	100
3	I	90/185 (49%)	90 (100%)	0	100	100
3	M	90/185 (49%)	90 (100%)	0	100	100
4	E	105/195 (54%)	105 (100%)	0	100	100
4	J	105/195 (54%)	105 (100%)	0	100	100
4	N	105/195 (54%)	105 (100%)	0	100	100
5	F	98/194 (50%)	98 (100%)	0	100	100
5	K	98/194 (50%)	98 (100%)	0	100	100
5	O	98/194 (50%)	98 (100%)	0	100	100
All	All	2670/4176 (64%)	2670 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	55	ASN
3	M	79	GLN
1	A	197	ASN
1	A	124	GLN
5	O	6	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-74907. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 216




Y Index: 216




Z Index: 216


6.2.2 Raw map



X Index: 216



Y Index: 216




Z Index: 216


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 234




Y Index: 209




Z Index: 234


6.3.2 Raw map



X Index: 234



Y Index: 209




Z Index: 233


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 5.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 102 nm^3 ; this corresponds to an approximate mass of 93 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.364 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.364 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.75	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.19	3.54	3.21

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.19 differs from the reported value 2.75 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-74907 and PDB model 9ZWE. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 5.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (5.3).

9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 77% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (5.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7710	0.6050
A	0.8900	0.6540
B	0.8890	0.6540
C	0.3310	0.4030
D	0.4860	0.4950
E	0.8820	0.6490
F	0.8330	0.6310
G	0.8890	0.6530
H	0.3220	0.4020
I	0.4800	0.4980
J	0.8800	0.6490
K	0.8300	0.6320
L	0.2990	0.3970
M	0.4780	0.4960
N	0.8680	0.6500
O	0.8300	0.6320

1.0

0.0

<0.0